RESEARCH ARTICLE

Cholinergic blockade reveals a role for human hippocampal theta in memory encoding but not retrieval

Tamara Gedankien[1], Jennifer Kriegel[2], Erfan Zabeh[1], David McDonagh[3], Bradley C Lega[2*†], Joshua Jacobs[1,4*†]

[1]Department of Biomedical Engineering, Columbia University, New York, United States; [2]Department of Neurological Surgery, University of Texas Southwestern, Dallas, United States; [3]Department of Anesthesiology, University of Texas Southwestern, Dallas, United States; [4]Department of Neurological Surgery, Columbia University, Dallas, United States

eLife Assessment

This **fundamental** work advances our understanding of the role of human hippocampal theta oscillations in memory encoding and retrieval. The evidence supporting the conclusions is **convincing**, using both scopolamine administration and intracranial EEG recordings. This work will be of broad interest to neuroscientists and has translational implications.

***For correspondence:** bradley.lega@utsouthwestern.edu (BCL); joshuajacobs@gmail.com (JJ)

†These authors contributed equally to this work

Competing interest: The authors declare that no competing interests exist.

Funding: See page 16

Preprint posted 22 August 2025
Sent for Review 05 September 2025
Reviewed preprint v1 04 December 2025
Reviewed preprint v2 27 February 2026

Reviewing Editor: Gui Xue, Beijing Normal University, China

© Copyright Gedankien *et al.* This article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use and redistribution provided that the original author and source are credited.

Abstract Cholinergic dysfunction is a hallmark of Alzheimer's disease and other memory disorders. Yet, the neurophysiological mechanisms linking cholinergic signaling to memory remain poorly understood. In this study, we administered scopolamine, a muscarinic cholinergic antagonist, to neurosurgical patients with intracranial electrodes as they performed an associative recognition memory task. When scopolamine was present at encoding, we observed disruptions to hippocampal slow theta oscillations (2–4 Hz), with selective impairments to recollection-based memory. However, when scopolamine was present during retrieval alone, we observed disruptions to slow theta without impaired memory performance. These disruptions included dose-dependent reductions in theta power, theta phase reset, and encoding–retrieval pattern reinstatement. Together, our results challenge the notion that theta oscillations are necessary for memory retrieval, and instead suggest that theta universally reflects an encoding-related neural state. These findings motivate updates to current models of acetylcholine's role in memory and may inform future therapies targeting rhythmic biomarkers of memory dysfunction.

Introduction

Decades of research have established a strong link between memory and the cholinergic system, which comprises the network of neurons that produce, release, and respond to acetylcholine. Evidence from neuroimaging (***Kilimann et al., 2014***; ***Roy et al., 2016***; ***Aghourian et al., 2017***; ***Sabri et al., 2018***), post-mortem studies (***Bowen et al., 1976***; ***Whitehouse et al., 1981***), and genetic research ***Weng et al., 2013*** demonstrate that Alzheimer's disease (AD) and other memory disorders are marked by widespread loss of cholinergic pathways. Established therapies for these conditions include cholinergic agonists to mitigate memory deficits (***Summers et al., 1986***; ***Hampel et al., 2018***). In experimental settings, cholinergic antagonists impair memory in both animals (***Hamburg, 1967***; ***Tang et al., 1997***; ***McGaughy et al., 2005***) and humans (***Drachman and Leavitt, 1974***; ***Ghoneim and Mewaldt,***

1975; *Petersen, 1977*; *Grober et al., 1989*; *Atri et al., 2004*; *Gedankien et al., 2023*). However, the neurophysiological mechanisms through which the cholinergic system influences memory during encoding and retrieval remain largely unknown, especially in humans. This knowledge gap creates a barrier to the development of new treatments for AD and other memory disorders associated with cholinergic imbalances.

In a previous study (*Gedankien et al., 2023*), we administered scopolamine—a cholinergic blocker targeting muscarinic acetylcholine receptors—to neurosurgical patients during a memory task and found that memory impairment was linked to disruptions in the amplitude and timing of hippocampal theta oscillations (2–10 Hz) during encoding. Theta power in the human hippocampus has been previously linked to successful episodic memory encoding (*Sederberg et al., 2003*; *Lega et al., 2012*), and our findings provided evidence that the power and phase of hippocampal theta are cholinergic-sensitive and, as such, potential therapeutic targets for memory disorders like AD.

However, in that initial study, our task design did not allow us to distinguish whether scopolamine's effects on memory and theta activity were specific to encoding, retrieval, or both. This distinction is critical, as deficits in memory encoding and retrieval are both hallmarks of advanced AD and other cholinergic-linked disorders (*Grober et al., 1987*; *Hodges et al., 1992*; *Perri et al., 2005*). Notably, prior behavioral studies suggest that cholinergic blockade during retrieval alone is not sufficient to cause memory impairments (*Mewaldt and Ghoneim, 1979*; *Beatty et al., 1986b*; *Tröster et al., 1989*; *Atri et al., 2004*; *Huang et al., 2011*). Yet, computational models posit that cholinergic tone influences theta oscillations across both encoding and retrieval (*Hasselmo and Bower, 1993*; *Hasselmo and Schnell, 1994*; *Hasselmo and Giocomo, 2006*). These models propose that acetylcholine dynamically gates hippocampal network states: promoting synaptic plasticity and theta rhythm alignment during encoding, while facilitating reactivation of stored information during retrieval through reduced cholinergic tone. Thus, cholinergic modulation was proposed as a mechanism for toggling the hippocampus between encoding and retrieval states (*Hasselmo et al., 2002*; *Kunec et al., 2005*; *Hasselmo, 2005*). To date, there has been no empirical evidence probing the neurophysiological effects of cholinergic modulation at retrieval in humans to test these models.

To address this question, we conducted an experiment combining human intracranial recordings with cholinergic modulation to isolate the effects of scopolamine at encoding versus retrieval as well as to investigate the specificity of modulation for recollection versus familiarity- and novelty-based memory. In a double-blind, within-subject design, we administered a single dose of scopolamine (or saline, in a placebo condition) to 12 neurosurgical patients as they performed an associative recognition memory task. Replicating prior behavioral findings (*Mewaldt and Ghoneim, 1979*; *Beatty et al., 1986b*; *Tröster et al., 1989*; *Atri et al., 2004*; *Huang et al., 2011*), we found no changes in memory performance when scopolamine was present during retrieval alone, but there were impairments to recollection-based memory when the drug was present prior to encoding. We then examined how scopolamine influenced three key neurophysiological markers of memory: theta power, theta phase reset, and encoding–retrieval pattern reinstatement. Despite a lack of behavioral impairments when the drug was present at retrieval alone, we found significant electrophysiological changes at retrieval, including disruptions to theta power, theta phase reset, and reinstatement of encoding-related spectral patterns. These results show that theta oscillations that appear during the recall period of a task are not essential to core memory retrieval processes, but instead could reflect other processes such as memory updating and re-encoding. We examine how these findings may promote updated theories of cholinergic tuning of the episodic memory system, with potential implications for novel therapeutic strategies.

Results

Cholinergic blockade during encoding, but not retrieval, impairs recollection memory

To investigate the effects of cholinergic modulation on retrieval, we administered scopolamine, a muscarinic blocker, to 12 neurosurgical patients with implanted depth electrodes while they performed an associative recognition (AR) memory task (*Figure 1a*). All participants provided written informed consent, and the study protocol was approved by the Institutional Review Board (IRB) of Human Subjects Research of the University of Texas Southwestern Medical Center (UTSW) prior to data



Figure 1. Cholinergic blockade during encoding, but not retrieval, impairs recollection memory. (**a**) Overview of methods. In order to study the neural mechanisms behind the effects of cholinergic blockade on human memory, we delivered scopolamine, a cholinergic blocker, to patients with intracranial encephalographic (iEEG) electrodes during an associative recognition memory task. (**b**) Diagram of the drug paradigm. In each session (drug or placebo), patients completed four blocks of the memory task. The blocks had the same task structure, starting with encoding followed by a math distractor and retrieval. In Block 2, subjects received a substance injection (scopolamine or saline), after encoding but prior to the retrieval period. Block 2's paradigm allowed us to examine whether scopolamine impacts retrieval independently from encoding. (**c**) Diagram of the task structure. Briefly, during the encoding period of the task, subjects are presented with a sequence of word pairs. Then, after completing a math distractor consisting of simple algebra equations, subjects initiate retrieval where they are asked to indicate whether word pairs presented are intact, rearranged, or new in relation to the word pairs presented during encoding. (**d**) Bar plot showing recall probabilities for encoding (placebo = all blocks, scopolamine = Blocks 3 and/or 4) and retrieval (placebo = all blocks, scopolamine = Block 2) across all trials. We found a significant decrease in memory performance when scopolamine is present at encoding, but not when it is present during retrieval alone (encoding Block 3 only: t=−42.0, p=9.77×10^{-3}; encoding Block 4 only: t=41.0, p=1.37×10^{-2}; encoding Blocks 3 and 4: t=41.0, p=3.91×10^{-3}; retrieval: t=32.0, p=0.15; Wilcoxon rank sum tests). (**e**) Bar plot showing change in memory as a function of block for all trials ($t(9) = -2.42$, p=1.5×10^{-2}, LME model), and separately for recollection, familiarity, and novelty trials (recollection:,$t(9) = -2.72$, p=6.5×10^{-3}; familiarity: $t(9) = -1.01$, p=0.31; novelty trials: $t(9) = -0.50$, p=0.61; LME models). (**f**) Correlation between percent change in memory (with respect to the placebo session) and dosage during recollection trials for Blocks 3 and 4 combined ($r = -0.80$, p=0.01, two-tail Pearson's correlation).

The online version of this article includes the following figure supplement(s) for figure 1:

Figure supplement 1. Scopolamine dosage correlates with changes in memory during recollection trials.

Figure supplement 2. Changes in memory following scopolamine for subjects with medium and high doses only.

Figure supplement 3. Bar plot showing change in memory across blocks for recollection trials.

collection. For each subject, we recorded intracranial electroencephalographic (iEEG) signals from the hippocampus and entorhinal cortex during two randomized double-blind sessions: an experimental condition in the presence of the drug (scopolamine) and a control condition with a placebo (saline). During the encoding phase of the AR task, subjects were presented with a series of word pairs. Following a math distractor, subjects began the retrieval phase in which they were cued with another series of word pairs. For each retrieval cue word pair, subjects indicated how it related to the encoding items, in particular whether it was intact, rearranged, or new with respect to encoding (***Figure 1c***). This paradigm allows us to dissociate neural correlates of recollection-based memory from recall based on familiarity and novelty detection.

By using a drug paradigm with intra-task scopolamine injection, we were able to probe the effects of cholinergic blockade on encoding and retrieval separately. In each session, subjects completed four blocks of the AR task (***Figure 1b***). In Block 1 (baseline), subjects completed encoding and retrieval without substance injection. In Block 2, subjects received a single injection of scopolamine (or saline, in the placebo condition) after encoding, but prior to retrieval. Thus, Block 2 allowed us to examine whether cholinergic blockade impacts retrieval independently from encoding, as individuals had encoded items in the absence of the drug but retrieved them after the injection. Lastly, subjects continued into Blocks 3 and 4, where both encoding and retrieval were affected by the substance administered in the earlier block. To assess how cholinergic blockade affected memory, we compared subjects' accuracy between blocks and conditions (drug versus placebo), separately analyzing performance for items that required memory processes related to recollection (intact pairs), familiarity (rearranged pairs), and novelty (new pairs).

First, as expected, we found that the presence of scopolamine during encoding significantly impaired memory performance overall (placebo = all blocks, scopolamine = Blocks 3 and 4; t=44.0, $p=3.91\times10^{-3}$; Wilcoxon rank sum test; ***Figure 1***, ***Figure 1—figure supplement 3***). The effect was also significant when including Block 3 and Block 4 individually (scopolamine = Block 3 only: t=42.0, $p=9.77\times10^{-3}$; scopolamine = Block 4 only: t=41.0, $p=1.37\times10^{-2}$; Wilcoxon rank sum tests). However, we found no significant changes in memory when cholinergic blockade was restricted to the retrieval stage (placebo = all blocks, scopolamine = Block 2; t=32.0, p=0.15, Wilcoxon rank sum test; ***Figure 1d***). This result is consistent with previous behavioral findings showing that cholinergic blockade during retrieval alone is not sufficient to elicit memory deficits (***Mewaldt and Ghoneim, 1979***; ***Beatty et al., 1986b***; ***Tröster et al., 1989***; ***Atri et al., 2004***; ***Huang et al., 2011***), and it was present even when excluding subjects with the lowest scopolamine doses (<400 mcg; ***Figure 1—figure supplement 2***). Notably, these effects were exclusive to items recovered on the basis of recollection rather than familiarity or novelty (recollection: $t(9) = -2.72$, $p=6.5\times10^{-3}$; familiarity trials: $t(9) = -1.01$, p=0.31; novelty: $t(9) = -0.50$, p=0.61; LME models) (***Figure 1e***). Lastly, we found that changes in memory correlated with scopolamine dose during Blocks 3 and 4 ($r = -0.80$, p=0.01, two-tail Pearson's correlation; ***Figure 1f***). Dosage effects were specific for items recalled on the basis of recollection, which highlights the selective impact of cholinergic blockade on memory (***Figure 1—figure supplement 1***).

Overall, our results indicate that the behavioral effects of scopolamine on associative memory are selective in two ways: first, by selectively disrupting episodic memory when present during encoding but not during retrieval alone; and second, by selectively affecting hippocampal-dependent encoding processes (i.e. recollection-based memory).

Cholinergic blockade disrupts theta power during retrieval

We next examined whether scopolamine affected electrophysiological signals at retrieval, even as memory performance remained unchanged. To test the effects of scopolamine on the amplitude of theta oscillations during retrieval, we computed changes in normalized power following the retrieval cue. For each electrode, we measured power during the entire 4-s-long retrieval period, and normalized the measured power relative to the mean and SD of the baseline period 0.5 s before stimulus onset. We then compared retrieval-related shifts in slow theta power (2–4 Hz) between drug and placebo blocks.

To test whether cholinergic blockade during retrieval alone altered oscillatory activity, we analyzed data from Block 2, where subjects encoded normally but retrieved in the presence of scopolamine. We found a significant disruption in slow theta power during retrieval in the scopolamine condition compared to placebo in Block 2 (t=−2.9, $p=3.37\times10^{-3}$, LME model; ***Figure 2c***). This result shows



Figure 2. Cholinergic blockade disrupts theta power during retrieval. (**a**) Anatomical locations of all hippocampal electrodes analyzed in the study. (**b**) Retrieval power effects for an example electrode during Block 4. Left: line plot displaying the raw iEEG trace (black) and corresponding slow theta (2–4 Hz) filtered signal (blue) from an example trial. Middle: time-frequency spectrograms displaying mean normalized power across all trials for the example electrode during the placebo and scopolamine sessions. Time 0 s denotes the onset of the retrieval cue, and contrast denotes changes in normalized power. Right: mean slow theta power over time for placebo (blue) and scopolamine (red). Shading denotes ± SEM. (**c**) Group-level mean slow theta power across all trials for Blocks 1, 2, 3, and 4 for placebo (blue) and scopolamine (red). Significant disruptions to slow theta power following scopolamine are present in Blocks 2 through 4 (p's < 0.05, LME models, multiple-comparison corrected). Shading denotes ± SEM. (**d**) Correlations between scopolamine dosage and change in slow theta power for all blocks (Block 1: $r = 0.19$, p=0.94; Block 2: $r = -0.17$, p=0.17; Block 3: $r = -0.29$, p=0.02; Block 4: $r = -0.55$, p=1.80×10^{-6}; one-tail Pearson's correlations). (**e**) Scatter plot showing change in slow theta power for all electrodes across blocks (t=−3.20, p=1.52×10^{-3}, one-sided t-test). (**f**) Change in power across blocks for different frequency bands. Only the slow theta frequency band shows a significant decrease in power across blocks. Dashed line denotes 95% significance level. (**g**) Bar plot showing changes in mean slow theta power as a function of subregion (Blocks 3 and 4: anterior hippocampus (AH): t=−2.9, p=0.02; posterior hippocampus (PH): t=−2.9, p=0.02; and entorhinal cortex (EC): t=−1.46, p=0.14; LME model, multiple-comparison corrected). Dashed line denotes 95% significance level. (h) Bar plot showing changes in mean slow theta power as a function of trial category (Blocks 3 and 4: recollection: t=−3.35, p=8.07×10^{-4}; familiarity: t=−3.26, p=1.10×10^{-3}; novelty: $t = -3.40$, p=6.77×10^{-4}; LME model, multiple-comparison corrected). Dashed line denotes 95% significance level.

The online version of this article includes the following figure supplement(s) for figure 2:

Figure supplement 1. Cholinergic blockade disrupts slow theta power during encoding.

Figure supplement 2. Disruptions to retrieval slow theta power in remembered trials only.

that cholinergic blockade during retrieval alone leads to theta amplitude disruption. When subjects encoded and retrieved in the setting of cholinergic blockade (Blocks 3 and 4), we also found a significant theta power disruption during retrieval in the scopolamine condition compared to placebo (t=−2.58, p=3.46×10^{-4}, LME model, multiple-comparison corrected), mirroring effects observed during encoding (***Gedankien et al., 2023***; ***Figure 2—figure supplement 1***). This effect can be seen on individual electrodes, as shown in ***Figure 2b***, as well as at the group level, as seen in ***Figure 2c***. Notably, the magnitude of these effects correlated with the absolute dosage of scopolamine (p's <

0.05, one-tail Pearson's correlations; ***Figure 2d***) and were present when including remembered trials only (***Figure 2—figure supplement 2***).

Next, to better understand how power changed as a function of drug-injection paradigm, we computed the correlation between mean changes in power for each electrode across blocks. We found a significant negative correlation for changes in slow theta power across blocks (t=−3.20, p=1.52×10^{-3}, one-sided t-test; ***Figure 2e***). This correlation was only significant for the slow theta band, highlighting the particular sensitivity of the slow theta rhythm to cholinergic modulation compared to other higher-frequency oscillations (***Figure 2f***).

To assess the robustness of this pattern and explore regional differences, we implemented a linear mixed-effects (LME) model that tested for differences in time-averaged theta power during retrieval. We found that power was significantly lower in the scopolamine condition compared to placebo, and this effect was driven by the anterior and posterior hippocampus (Blocks 3 and 4: anterior hippocampus (AH): t=−2.32, p=0.02; posterior hippocampus (PH): t=−2.36, p=0.02; and entorhinal cortex (EC): t=−1.46, p=0.14; LME models, multiple-comparison corrected; ***Figure 2g***). Lastly, we also found that changes in slow theta power during retrieval were maintained across recollection, familiarity, and novelty trials (Blocks 3 and 4: all p's < 0.002, LME models, multiple-comparison corrected; ***Figure 2h***).

In sum, these findings indicate that cholinergic blockade changes iEEG activity during retrieval by disrupting the amplitude of slow theta oscillations, mirroring effects previously observed during encoding (***Gedankien et al., 2023***). Importantly, despite the robust disruption to theta power during retrieval, we did not observe accompanying behavioral impairments when scopolamine was present during retrieval alone (Block 2). These results suggest the possibility that theta oscillations measured during the retrieval stage of a task may serve a function other than supporting recall processes.

Dimensionality reduction of spectral power enables decoding of drug condition

To uncover latent structure in the data, we applied principal component analysis (PCA) to the power spectra. This unsupervised method allows us to characterize the correlational structure of retrieval-related oscillations across frequencies and brain regions, and identify features associated with pharmacological condition (***Figure 3a and b***). In doing so, this data-driven analysis, which integrates data across multiple frequencies, complements our theta-focused results shown above.

We applied PCA to power data filtered in predefined frequency bands to identify low-dimensional electrophysiological patterns that may appear across multiple frequencies (***Figure 3a***). Each resulting principal component (PC) consists of a weighted sum of power across frequency bands, capturing variability between trials. Across all electrodes, we found that the first PC accounted on average for nearly 80% of the total variance (***Figure 3c and d***). The substantial variance captured by a single component suggests that, despite the complexity of the underlying neural circuits, the activity patterns relevant to memory retrieval are governed by coherent, large-scale fluctuations across frequencies and electrodes (***Miller et al., 2009***).

To evaluate whether the PCA space could be used to decode pharmacological condition, we trained two linear classifiers. In the first model, we performed decoding at the electrode level, using the first three PCs as features of spectral subspace. We found that decoding accuracy was significantly above chance across all electrodes (p<0.001, Wilcoxon signed-rank test) (***Figure 3e***), demonstrating that the compressed PCA representation (even when using only the first PC, ***Figure 3—figure supplement 1***) retained sufficient information to distinguish between the placebo and scopolamine conditions. In the second model, we modified the structure and trained frequency-band-specific decoders, using electrode identity as the feature space. We found that decoding accuracy was highest for the slow theta band, exceeding that of both fast theta and slow gamma bands (***Figure 3f***). Furthermore, we observed a positive relationship between decoding accuracy and the number of electrodes included in the model (***Figure 3f***), suggesting that different subregions within the hippocampus contribute distinct electrophysiological signatures to the classification.

Together, these results demonstrate that scopolamine-related LFP activity can be accurately decoded from a low-dimensional feature space. Moreover, our results suggest that cholinergic blockade modulates neural dynamics along a restricted set of spectral patterns, which can enhance decoding and reflect selective engagement of dominant circuit activity, as seen in other neuronal modalities (***Kaufman et al., 2014***; ***Gallego et al., 2017***). Importantly, the strongest decoding



Figure 3. Dimensionality reduction of spectral power enables decoding of drug condition. (**a**) Schematic panel outlining the analysis pipeline, including frequency-band data processing, principal component analysis (PCA) decomposition, and classification methods. This unsupervised method allows us to characterize the correlational structure of retrieval-related oscillations across frequencies and brain regions, and identify features associated with pharmacological condition. (**b**) Schematic representation of low- and high-dimensional PCA decompositions. The separation of trajectories illustrates distinct neural representations across conditions. In low-dimensional problems, one or a few principal components (PCs) capture most of the variance and may be sufficient to support accurate decoding. In contrast, high-dimensional problems typically require many components to capture relevant variance, making decoding more complex. (**c**) Bar plot showing variance explained by the first three PCs, emphasizing the dominant contribution of PC1. Error bars denote ± SEM across electrodes. (**d**) Histogram displaying mean cumulative variance (average of the first three PCs) across electrodes. The mean cumulative variance for all individual electrodes is greater than 0.9, indicating low-dimensionality in the data decomposition. (**e**) Left: PCA visualization for an example electrode, showing neural trajectories from the placebo (blue) and scopolamine (red) conditions in the first three principal components. We performed dimensionality reduction for each electrode and decoded condition (placebo or scopolamine) using the resulting PCs. Right: histogram of decoding accuracies across electrodes with a significant decoding accuracy peak at 0.6 ($p<0.001$, Wilcoxon signed-rank test). Dashed line indicates chance-level decoding (0.5). (**f**) Left: schematic of frequency-band-level decoder. We decoded condition (placebo or scopolamine) using frequency-band-filtered data across electrodes. Right: line plot comparing decoding accuracies (AUC) using slow theta, fast theta, and slow gamma frequency-band-filtered data with increasing number of electrodes. Decoding accuracy increased with number of electrodes and was highest for the slow theta band, exceeding that of both fast theta and slow gamma bands.

The online version of this article includes the following figure supplement(s) for figure 3:

Figure supplement 1. Dimensionality of spectral decoding subspace.

performance emerged from the slow theta band, aligning with our hypothesis-driven findings that identified disruptions in slow theta oscillations following scopolamine administration (***Figure 2***). This convergence across analytical approaches reinforces the conclusion that cholinergic modulation critically shapes slow theta dynamics during memory processing.

Cholinergic blockade impairs theta phase alignment during retrieval

Building on evidence that the phase of ongoing neuronal rhythms facilitates memory functions (***Jutras et al., 2013***; ***Ter Wal et al., 2021***) and earlier work showing disruptions to theta phase reset during encoding following scopolamine (***Gedankien et al., 2023***), we examined whether cholinergic blockade altered phase dynamics of theta during retrieval. We examined whether cholinergic blockade leads to changes in theta phase alignment by computing the inter-trial phase coherence (ITPC, or phase reset) at the time of the retrieval cue.



Figure 4. Cholinergic blockade impairs theta phase alignment during retrieval. (**a**) Retrieval inter-trial phase coherence (ITPC, or phase reset) effects for an example electrode during Block 2. Time 0 s denotes onset of retrieval cue, and contrast denotes strength of phase reset. During the placebo session, the electrode shows increased slow theta (2–4 Hz) phase reset following cue onset. This effect is absent during the scopolamine session. (**b**) Polar plots showing phase distribution for a 3 Hz oscillation at 0.3 s for the same example electrode. During the placebo session, the electrode exhibits significant phase clustering during Blocks 2, 3, and 4 (Block 2: $z = 14.9$, p=2.27×10^{-7} Block 3: $z = 9.16$, p=9.15×10^{-5}; Block 4: $z = 4.63$, p=9.54×10^{-3}; Rayleigh tests), but that effect is absent during the scopolamine session (Block 2: $z = 0.23$, p=0.79 Block 3: $z = 0.51$, p=0.60, Block 4: $z = 2.78$, p=0.06; Rayleigh tests). (**c**) Group-level mean slow theta phase reset is significantly disrupted following scopolamine in Blocks 2 through 4 ($p's < 0.05$, LME models, multiple-comparison corrected) across trials. Shading denotes ± SEM. (**d**) Correlations between scopolamine dosage and change in slow theta phase reset for all blocks (Block 1: $r = 0.29$, p=0.98; Block 2: $r = -0.27$, p=0.02; Block 3: $r = -0.30$, p=0.01; Block 4:,$r = -0.10$, p=0.24; one-tail Pearson's correlations). (**e**) Group-level phase reset effects at encoding. (**f**) Bar plot showing mean phase clustering across all electrodes during placebo and scopolamine for the encoding and retrieval periods. Scopolamine disrupts slow theta phase reset during both encoding and retrieval periods (encoding: ,t=−2.07, p=0.04; retrieval: t=−2.69,p=0.01, paired t-tests). (**g**) Phase reset effects for an example electrode during placebo. Increased 2–4 Hz ITPC is present at both encoding and retrieval. (**h**) Example electrode effects for a 3 Hz oscillation at 0.5 s. Top: mean phase. Bottom: corresponding phase difference. Phase difference between encoding and retrieval is constant and nearly zero between 0 and 1 s. (**i**) Group-level consistency of slow theta phase differences between encoding and retrieval during placebo. Dashed line denotes Rayleigh test significance (95% confidence level). (**j**) Group-level distribution of slow theta phase differences at 0.5 s ($z = 8.05$,,p=2.98×10^{-4} Rayleigh test).

The online version of this article includes the following figure supplement(s) for figure 4:

Figure supplement 1. Disruptions in retrieval phase reset following cholinergic blockade are driven by the anterior and posterior hippocampus.

Figure supplement 2. Mean preferred phase at encoding is recapitulated at retrieval.

In the placebo session, we found that theta oscillations exhibited significant phase reset following the retrieval cue (***Figure 4a–c***). However, the magnitude of theta phase reset was significantly reduced in Blocks 2 through 4, when the drug was present at recall (all $p's < 0.05$, LME models, multiple-comparison corrected). This can be seen in individual electrodes as shown in ***Figure 4a and b***, and

at the group level in ***Figure 4c***, where the effects were driven by the anterior and posterior hippocampus (***Figure 4—figure supplement 1***). Furthermore, we found significant correlations between changes in slow theta phase reset and scopolamine dosage ($p's < 0.05$, one-tail Pearson's correlations) (***Figure 4d***).

Since scopolamine disrupted theta phase in both the encoding and retrieval stages of the task (encoding: t=−2.07, p=0.04; retrieval: t=−2.69, p=0.01; paired t-tests) (***Figure 4e and f***), we next examined the similarity in theta phase dynamics between encoding and retrieval under drug-free conditions. For each electrode during the placebo session, we computed the differences between the instantaneous phase at matched time points relative to the encoding and retrieval cues (***Figure 4g and h***). We found significant matches in the phases between encoding and retrieval between 0 and 0.75 s of cue onset (***Figure 4i***), with strongest clustering at 0.5 s post stimulus onset ($z = 16.6$, p=1.49×10^{-8}, Rayleigh test) (***Figure 4j***). This result indicates that theta phase alignment during encoding is recapitulated during retrieval. Additionally, between 0 and 0.75 s, the preferred theta phase at retrieval closely matched that of encoding, with a mean phase difference of 8.83° at 0.5 s (***Figure 4j***, ***Figure 4—figure supplement 2***).

Taken together, our findings indicate that cholinergic blockade disrupts theta phase reset at retrieval in a manner that closely mirrors previously observed disruptions during encoding (***Gedankien et al., 2023***). Rather than supporting separate phase regimes for different memory stages (***Hasselmo and Bower, 1993***; ***Hasselmo and Schnell, 1994***; ***Hasselmo and McGaughy, 2004***), our results suggest that theta phase reset at retrieval reflects a reinstantiation of the encoding state, which has implications for our understanding of the role of theta in memory recall (see *Discussion*).



Figure 5. Cholinergic blockade prevents encoding–retrieval pattern reinstatement. (**a**) Schematic panel outlining methods for spectral pattern reinstatement analysis. First, we identified matching encoding and retrieval periods for all recollection trials. We then computed the correlation of 4–128 Hz power between encoding and retrieval across 250ms time bins during the initial 3 s following the encoding or retrieval cue. The resulting similarity matrix quantifies spectral reinstatement by showing whether spectral patterns at different time points during encoding are reinstated at similar, later, or earlier time points during retrieval. (**b**) Similarity matrix for example electrode. The 'no drug' plot reflects the mean across all blocks within the placebo session, whereas the 'scopolamine' plot shows combined data from Blocks 3 and 4. The contrast denotes the Pearson's correlation coefficient (PCC) of 4–128 Hz power data between encoding and retrieval. Black outlines denote pixels exhibiting greater-than-chance reinstatement compared to surrogate data (95% confidence level). Surrogate data for electrode-level analyses involved shuffling trial labels. (**c**) Bar plot showing mean reinstatement across conditions. Retrieval reinstatement during the no-drug condition is significantly greater than during the scopolamine conditions (no-drug versus scopolamine Block 2: t=−5.04, p=1.05×10^{-6}; no-drug versus scopolamine Blocks 3 and 4:,t=5.06, p=9.68×10^{-7}; paired t-tests). (**d**) Group-level mean reinstatement matrices for no-drug condition, scopolamine Block 2, and scopolamine Blocks 3 and 4 combined. Black outlines denote pixels exhibiting greater-than-chance reinstatement compared to surrogate data (95% confidence level). Surrogate data for group-level analyses involved shuffling block labels.

Cholinergic blockade prevents encoding–retrieval pattern reinstatement

Previous studies, including noninvasive (***Polyn et al., 2005***; ***Griffiths et al., 2019***) and intracranial EEG experiments (***Jacobs and Kahana, 2009***; ***Kunz et al., 2024***), have shown that stimulus-specific spectral representations emerge during memory encoding and are reactivated during retrieval—a phenomenon known as reinstatement. We therefore tested whether scopolamine disrupts spectral power reinstatement in the hippocampus to evaluate how cholinergic modulation shapes the neural conditions that support memory retrieval.

To assess whether scopolamine altered neural pattern reinstatement, we focused on recalled trials from the recollection category. We computed the correlation of spectral power (4–128 Hz) between the corresponding encoding and retrieval periods in 250ms time bins (***Figure 5a***). We chose a broadband frequency range of 4–128 Hz since reinstatement patterns have been observed at a wide range of frequency bands (***Manning et al., 2011***; ***Staresina et al., 2016***). However, since we report changes in slow theta power between conditions, we excluded the slow theta frequency range (2–4 Hz) from the reinstatement correlations to avoid confounding effects. The resulting correlation matrix for an example electrode is shown in ***Figure 5a***: the color contrast indicates the strength of the spectral power correlation between encoding and retrieval for each combination of time-bin pairs. This similarity matrix allows us to quantify whether spectral patterns from encoding are reinstated during matched, earlier, or later time points during retrieval.

We found that the strength of overall reinstatement was significantly greater under drug-free conditions than under scopolamine conditions. This can be seen at individual electrodes (***Figure 5b***), where reinstatement is diminished during the scopolamine condition ($z = 1.99$, $p<0.05$, two-proportion z-test). Similar effects appear at the group level (***Figure 5c and d***), where reinstatement during no-drug conditions is significantly greater than during the scopolamine conditions (no-drug versus Block 2: $t=5.04$, $p=1.05\times10^{-6}$; no-drug versus Blocks 3 and 4: $t=5.06$, $p=9.68\times10^{-7}$; paired t-tests) (***Figure 5c***). In particular, the no-drug condition shows a mean reinstatement pattern that is clustered in the matrix's diagonal, corresponding to matched time points between encoding and retrieval. This reinstatement pattern is absent during Block 2 and Blocks 3 and 4 in the scopolamine conditions (***Figure 5d***). This result indicates that cholinergic blockade disrupts both the magnitude and the timing of encoding–retrieval spectral pattern reinstatement.

Overall, these findings demonstrate that scopolamine disrupts the reinstatement of encoding-related LFP spectral patterns during retrieval, even in the absence of behavioral impairments. This suggests that while cholinergic tone may not be strictly necessary for retrieval, it may support the reactivation of an encoding-like mesoscopic neural state.

Discussion

Although there is ample evidence that the cholinergic system is crucial for episodic memory, the specific neurophysiological mechanisms by which it modulates human memory circuits remain largely unknown. In this study, we used rare human intracranial recordings coupled with scopolamine administration to investigate how cholinergic blockade influences neuronal activity, describing neurophysiological effects during episodic memory retrieval for the first time. Replicating existing behavioral findings (***Mewaldt and Ghoneim, 1979***; ***Beatty et al., 1986b***; ***Tröster et al., 1989***; ***Atri et al., 2004***; ***Huang et al., 2011***), we showed that memory was not disrupted when scopolamine was present during retrieval alone (***Figure 1***). However, when scopolamine was present during encoding, we found that it specifically impaired memory for items recovered on the basis of recollection, without impact on familiarity-based recovery or the identification of novel items. These findings are consistent with the proposed role of cholinergic circuits in mediating associative processes within the medial temporal lobe (MTL; ***Hasselmo, 2006***; ***Teles-Grilo Ruivo and Mellor, 2013***). We then found that scopolamine induced dose-dependent disruptions to the amplitude (***Figures 2 and 3***) and phase synchronization (***Figure 4***) of slow theta oscillations during retrieval, including trials when the drug was present during retrieval alone, where no behavioral impairments occurred. Finally, we also showed that cholinergic blockade prevented the reinstatement of spectral patterns from encoding during retrieval (***Figure 5***). Our unique results present a puzzle: why is there a disruption in an oscillatory process that supports associative memory but no commensurate change in behavior? Below, we examine how these data

stimulate updated models of cholinergic influence on episodic memory processing as well as models of theta oscillations in the human MTL.

Our behavioral results extend a body of research that explored how cholinergic blockers affect behavior in both animals and humans (***Hamburg, 1967***; ***Tang et al., 1997***; ***McGaughy et al., 2005***; ***Drachman and Leavitt, 1974***; ***Ghoneim and Mewaldt, 1975***; ***Petersen, 1977***; ***Grober et al., 1989***; ***Gedankien et al., 2023***). In humans, cholinergic blockade has been shown to impair memory when present during encoding but not during retrieval alone (***Mewaldt and Ghoneim, 1979***; ***Beatty and Bierley, 1986a***; ***Tröster et al., 1989***; ***Atri et al., 2004***; ***Huang et al., 2011***). Our results replicate these findings, as we did not find significant differences between Block 1 (the drug-free baseline) and Block 2 (when the drug was present during retrieval only). Consistent with previous work (***Gedankien et al., 2023***), memory was significantly impaired in Blocks 3 and 4 (when the drug was present during encoding and retrieval). Together, these results support the idea that cholinergic blockade primarily disrupts encoding but spares retrieval processes. Our behavioral results expand earlier findings by demonstrating that memory disruptions are exclusive to recollection-based memory (***Figure 2d***). These findings—identified using a version of the AR paradigm that is able to tease apart recollection, familiarity, and novelty-based mnemonic processes—provide unique evidence for the importance of cholinergic signaling to generate item/context associations at encoding. By highlighting a selective disruption to recollection-based memory, our results shed light on potential constraints of cholinergic function in mnemonic processes. To devise treatments that address the full spectrum of memory deficiencies observed in patients with cholinergic imbalances, it will be essential to further qualify the impact of cholinergic blockade on different types of memory.

Further, our experiments directly explored how cholinergic modulation alters neurophysiological processes at retrieval. Based on previous theoretical models (***Hasselmo and Bower, 1993***; ***Hasselmo and Schnell, 1994***; ***Hasselmo and McGaughy, 2004***) and the lack of memory impairment following cholinergic blockade during retrieval alone (***Mewaldt and Ghoneim, 1979***; ***Beatty et al., 1986b***; ***Tröster et al., 1989***; ***Atri et al., 2004***; ***Huang et al., 2011***; ***Figure 1***), we hypothesized that scopolamine would not disrupt hippocampal theta oscillatory dynamics during retrieval. On the contrary, we found that cholinergic blockade universally disrupted slow theta power (***Figures 2 and 3***) and phase reset (***Figure 4***) during retrieval, including during trials when the drug was present during retrieval alone and encoding occurred under drug-free conditions (Block 2). Together, these observations challenge the notion that theta oscillations directly support neural processes necessary for retrieval. A key function of theta oscillations in animals (***O'Keefe and Recce, 1993***) and humans (***Rutishauser et al., 2010***; ***Yoo et al., 2021***; ***Qasim et al., 2021***) is to support the coordinated firing of neuronal ensembles on precise time scales, establishing sequences through fundamental mechanisms like spike-timing-dependent plasticity. During encoding, the importance of this process is clear: theta oscillations provide a temporal scaffold for facilitating plasticity and forming sparse neural representations, and our results are fully consistent with this model. As for retrieval, our data suggest that theta oscillations are not critical for the re-representation of those ensembles at the time of recall—because theta disruption at retrieval was not coupled with behavioral deficits. One possible synthesis of our findings with existing models is that the observed theta activity at retrieval may reflect a mechanism supporting memory updating processes that imbue flexibility and integration to representations. This potential interpretation extends earlier models (i.e. SPEAR ***Hasselmo and Bower, 1993***; ***Hasselmo and Schnell, 1994***; ***Hasselmo and McGaughy, 2004***) that posit distinct theta phases for encoding and retrieval, by emphasizing that even during retrieval, the presence of theta reflects cholinergic-dependent encoding operations. This interpretation is also consistent with modern conceptions of episodic representations, which note that memory is inherently dynamic. As such, episodic memory retrieval during real-world behavior is not a passive playback of stored information, but an active, constructive process that often involves integrating the retrieval cue with ongoing contextual input.

Moreover, our PCA and decoding analyses provided new insight into how cholinergic modulation shapes memory-related neural dynamics. By showing that retrieval-related hippocampal LFP activity can be captured by a small number of PCs, we demonstrate that cholinergic disruption alters coherent, large-scale spectral patterns despite the complexity of the underlying circuits. This low-dimensional structure was sufficient to decode pharmacological condition and aligned with our hypothesis-driven results, particularly in highlighting the role of slow theta oscillations. Notably, decoding accuracy increased with the number of electrodes included, and frequency-band-specific models revealed

that different electrodes contributed distinct, meaningful features to the classification. These findings suggest that theta is not a spatially uniform signal but rather exhibits regional specificity across the hippocampus, with subregions expressing unique electrophysiological signatures in response to cholinergic disruption. The superior decoding performance of the slow theta band, combined with its spatial heterogeneity, reinforces its selective vulnerability to cholinergic modulation. The convergence of unsupervised and supervised analyses reinforces the conclusion that cholinergic tone modulates hippocampal theta dynamics and highlights how local circuit activity can be leveraged to decode behavioral or pharmacological states.

Our results also include the unique finding that cholinergic blockade impairs reinstatement of oscillatory patterns at the time of retrieval. Our proposed interpretation of theta as an encoding-related signal also invites a reconsideration of the nature and functional significance of encoding–retrieval reinstatement measured at the LFP-level. Traditionally, reinstatement refers to the reactivation of neural patterns present during encoding at the time of retrieval—a mechanism thought to support successful memory recall. This property, extensively described using intracranial EEG and noninvasive imaging (***Polyn et al., 2005***; ***Manning et al., 2011***; ***Staresina et al., 2016***), implies that theta synchronization, at least during encoding, supports the ability of the hippocampus to regenerate activity patterns at cue. However, if theta at retrieval reflects flexible updating of retrieved information, then recapitulation of oscillatory activity may not reflect the original memory trace in isolation, but the similarity of these theta-dependent processes between encoding and retrieval. We propose that the reinstatement signal we measure is composed of two overlapping components: one reflecting the reactivation of specific memory representations (e.g. single neurons that encoded the original experience), and another reflecting the oscillatory mechanisms that support the modification or reconsolidation of those representations. Our LFP-based reinstatement analyses lack the resolution to isolate single-neuron activity, and instead capture the broader oscillatory dynamics that support re-encoding. This distinction suggests that LFP-based reinstatement is not a direct proxy for the reinstatement of content, but rather an index of the brain's entry into an encoding-like state that facilitates representational integration. The observed disruption to LFP reinstatement under scopolamine, then, may not reflect a failure to retrieve, but rather a failure to initiate the processes necessary for memory updating. This perspective reframes LFP reinstatement as an approximation of encoding-state reinstatement, rather than content-specific replay, and further supports our proposal that theta oscillations serve a unifying role in encoding and re-encoding across memory operations.

An alternative theory to our interpretation of theta as an encoding-related neural state is that theta plays a role in retrieval, but cued retrieval mobilizes several partly redundant representational systems, so that disruption of hippocampal theta power and phase degrades only one route to the past without fully abolishing recall capabilities. Extra-hippocampal structures such as the parahippocampal and retrosplenial cortices, posterior parietal cortex, and medial prefrontal networks have all been shown to sustain successful recall (***Ranganath and Ritchey, 2012***; ***Rugg and Vilberg, 2013***; ***Ritchey et al., 2015***). This redundancy is consistent with distributed frameworks in which the hippocampus orchestrates, but does not monopolize, episodic memory reconstruction (***McClelland et al., 1995***; ***Moscovitch et al., 2016***). Selective disruption of theta phase reset may therefore impair temporal sequencing while parallel extra-hippocampal pathways preserve overall retrieval performance. Even under this alternative view, our findings show that hippocampal theta is not critically required for retrieval in the same way it is for encoding.

In sum, our findings provide converging behavioral and electrophysiological evidence that cholinergic modulation plays a critical role in memory by selectively supporting hippocampal encoding processes. By showing that theta oscillations are disrupted by scopolamine during retrieval—even in the absence of behavioral impairments—we challenge the classical view that theta reflects retrieval-specific computations. Instead, we argue that theta activity that shows up at retrieval reflects an encoding-related neural mode, re-engaged during retrieval to support memory updating, contextual integration, and re-encoding. This interpretation suggests that our LFP-level encoding–retrieval reinstatement reflects the reactivation of an encoding state rather than of content alone, with implications for how we conceptualize retrieval in dynamic, reconstructive terms. More broadly, these findings prompt a reassessment of theta's role in organizing and generating neural sequences during retrieval and underscore the importance of distinguishing between mechanisms that support memory access and those that enable memory transformation. Future models may need to account for a role of theta

oscillations in dynamically modifying previously encoded information during retrieval. Understanding how neuromodulatory systems like cholinergic pathways gate these processes will be essential for advancing both theoretical frameworks of memory and clinical strategies for memory disorders.

Methods

Participants

Twelve subjects participated in the study (***Supplementary file 1***). All participants were epilepsy patients who received iEEG implants as part of their treatment plan at the UTSW epilepsy surgery program.

Ethics and safety

All participants provided written informed consent to participate in the study and for the publication of the results. The study protocol was approved by the IRB of Human Subjects Research of the UTSW prior to data collection. All procedures were conducted in accordance with relevant institutional guidelines and regulations. Intravenous scopolamine was administered by the participant's attending nurse, and a board-certified anesthesiologist was present at the time of injection and remained available throughout the duration of the experiment to treat any potential adverse reactions. Physiological monitoring of blood pressure, electrocardiogram, oxygen saturation, and mental status was performed just before and for 1 hr after the injection of scopolamine. Low doses of scopolamine (such as 500 mcg) are commonly administered in clinical settings and generally do not present major risk factors (***Renner et al., 2005***). There were no reports of adverse events related to substance administration under drug or placebo conditions.

Scopolamine injection

Participants received a single dose of scopolamine (or saline, in the placebo session) via an intravenous line. The dosage of scopolamine for each participant was adjusted by the subject's weight in an attempt to normalize effect sizes across subjects. The drug has a half-life of 69 min and starts to take effect approximately 5 min after administration (***Renner et al., 2005***). The second session using the alternate agent (scopolamine or saline) took place 1–3 days after the first session. The drug/placebo randomization was unknown to the test administrators and participants.

Drug paradigm

Subjects completed two sessions (drug and placebo), and each session consisted of four blocks of a memory task (***Figure 1b***). The blocks followed the same behavioral paradigm, but different drug paradigms. In Block 1, subjects completed the task without any substance injections. In Block 2, subjects received an injection of scopolamine or saline after encoding but prior to the retrieval period. In Blocks 3 and 4, subjects completed the task under the effects of the scopolamine (or saline) injection from Block 2 but without new interventions (***Figure 1b***). This paradigm enables us to probe the effects of scopolamine on encoding and retrieval separately. In Block 2, the substance injection was administered 16±2 min after the end of encoding, and the retrieval period commenced at 15±0.2 min after injection (i.e. the total time between encoding and retrieval during Block 2 was approximately 31 min). The total session duration was 2 hr and 26 min ±23 min, and there were no time delays between blocks beyond the substance injection procedure described above.

Memory task

In each block, subjects completed an AR memory task (***de Chastelaine et al., 2011***; ***Figure 1c***). During encoding, participants were shown word pairs, and later during retrieval participants were cued with word pairs that were either intact (the same as encoding), rearranged (mixed with other previously shown word pairs), or new (previously unseen words). For each retrieval cue, subjects were asked to indicate whether the cued word pair was intact, rearranged, or new. The encoding period included 90 word pairs, and each pair was displayed in capital letters for 3 s. To enhance and control for attention during the encoding period, after each word pair subjects were asked to indicate which of the objects in the word pair was more likely 'to fit into the other'. After the encoding period, subjects completed a brief math distractor task, where they had to indicate if simple arithmetic problems in the form A+B

+ C=D were true or false. Next, participants began the retrieval period, which included 120 trials (60 intact, 30 rearranged, and 30 new word pairs), with cues presented for 4 s. During both encoding and retrieval, word pair cues were followed by an 0.8–1.2 s blank interstimulus interval (ISI) to decorrelate physiological responses from consecutive item presentations. Word pairs were randomized without repetitions across blocks.

Intracranial recordings

Patients were implanted with either Ad-Tech Medical or PMT depth electrodes with 0.8 mm in outer diameter, forming 10–14 recording contacts arrayed at 4–5 mm intervals along the shaft of the electrode (Ad-Tech: 10 electrodes with variable spacing; PMT: 10–14 electrodes with uniform spacing, depending upon the overall depth of insertion). Intracranial EEG was sampled at 1 kHz on a Nihon–Kohden 2100 clinical system under a bipolar montage with adjacent electrodes as reference. A kurtosis threshold of 4 was applied for each trial to exclude abnormal events and interictal activity. Subject 3 was excluded from the electrophysiological analyses due to an issue with the trigger pulses. Subjects 8, 10, and 11 were excluded from the behavioral and electrophysiological analyses due to logfile errors. The electrode coverage included contacts in the posterior and anterior hippocampus, and entorhinal cortex (***Supplementary file 1***).

Anatomical localization

The anatomical locations of the depth electrodes were chosen by the neurology team based on each patient's clinical needs. Electrodes were laterally inserted into the specified regions with robotic assistance. Final electrode localization within hippocampal subregions was determined by post-operative neuroradiology review.

Spectral power analysis

For the power analysis, we extracted spectral power using a continuous Morlet wavelet transform (wave number = 6) over 49 logarithmically spaced frequencies ranging from 2 to 128 Hz. The analysis encompassed the full 4 s retrieval periods following cue onset, with an additional 3 s buffer to minimize edge artifacts. Power values for each session, electrode, and frequency were normalized against the mean and standard deviation of the baseline period, defined as the 0.5 s preceding cue onset. To identify significant differences between conditions, we applied an LME model, using time-averaged power during the 4 s retrieval period as the dependent variable and condition (scopolamine/placebo) as a fixed effect, with electrodes treated as a random effect.

Principal component analysis

To perform the PCA, frequency-band power data for each electrode were represented as a matrix $X \in \mathbb{R}^{T \times F}$, where T is the number of time points, and F is the number of spectral features. The covariance matrix of X was computed as:

$$C = \frac{1}{T} X^{\top} X \tag{1}$$

where $C \in \mathbb{R}^{F \times F}$ represents the pairwise covariance between features. The eigenvectors $V \in \mathbb{R}^{F \times F}$ and eigenvalues $\lambda \in \mathbb{R}^{F}$ of C were then computed, where the eigenvectors define the PCs, and the eigenvalues represent the variance explained by each component. The data were projected onto the first three PCs for visualization $X_{\mathrm{PCA}} = XV_{1:3}$, with $V_{1:3}$ containing the eigenvectors corresponding to the top three eigenvalues. The proportion of variance explained by each component was calculated as:

$$\text{Variance Explained} = \frac{\lambda_i}{\sum_{j=1}^{F} \lambda_j} \tag{2}$$

where λ_i is the eigenvalue of the i-th PC.

Decoding analysis

The first decoding analysis involved an electrode-level logistic regression classifier. The input to the classifier was the power of specific frequency bands projected onto the PCs. For each trial t, the input features were represented as $X_t \in \mathbb{R}^F$, and the binary output label $y_t \in \{0, 1\}$ indicated the drug condition. The logistic regression model was defined as:

$$P(y_t = 1|X_t) = \sigma(w^\top X_t + b) \tag{3}$$

where $w \in \mathbb{R}^F$ is the weight vector, b is the bias term, and $\sigma(z)$ is the sigmoid function $\sigma(z) = \frac{1}{1+e^{-z}}$. The model was trained using a cross-entropy loss function:

$$\mathcal{L} = -\frac{1}{N}\sum_{t=1}^{N}\left[y_t \log(P(y_t)) + (1 - y_t)\log(1 - P(y_t))\right] \tag{4}$$

where N is the total number of trials. The data were divided into training (80%) and testing (20%) sets using stratified random sampling to ensure balanced class distributions. Decoding accuracies were aggregated across electrodes, and significance was determined using permutation testing with 1000 label shuffles.

In the second decoding analysis, we performed classification using trial-wise power features from slow theta (2–4 Hz), fast theta (4–8 Hz), and slow gamma (30–60 Hz) bands to compare the discriminative power of individual frequency bands and to assess whether decoding accuracy scales with spatial coverage. For each patient, we incrementally increased the number of electrodes used in the decoder, from 1 up to the total available. At each step, we randomly sub-sampled that number of electrodes 100 times and trained a linear support vector machine to classify trial conditions. A fixed regularization parameter (C=10) was used to prevent overfitting and dampen the influence of potentially noisy or uninformative electrodes (***Pereira et al., 2009***). Trial labels were balanced across conditions. All preprocessing steps, including normalization and electrode subsampling, were performed within the training data in each fold. Performance was estimated using stratified fivefold cross-validation, and accuracy was averaged across folds and repetitions. Analyses were performed independently for each patient, and final results were averaged across patients.

Phase reset analysis

We extracted phase measures using a continuous Morlet wavelet transformation (wave number = 6) across 49 logarithmically spaced frequencies from 2 to 32 Hz. We then computed the phase reset of oscillations at the retrieval cue onset using the inter-trial phase coherence (ITPC) measure (***Tallon-Baudry et al., 1996***; ***van Diepen and Mazaheri, 2018***), which quantifies the consistency of instantaneous phase measures across trials as follows:

$$ITPC(f, t) = \frac{1}{N}\left|\sum_{k=1}^{N} e^{i\varphi_k(f,t)}\right| \tag{5}$$

where f is the frequency for a given time t, N is the number of trials, and $e^{i\varphi^k}$ is the polar representation of the phase angle φ. Since ITPC calculations are susceptible to the number of observations, we excluded Subject 5, who had an incomplete number of retrieval trials (<120) in the scopolamine condition. The ITPC was computed separately for individual electrodes and averaged at the subject-level for the group analysis (***Gedankien et al., 2023***).

Reinstatement analysis

To quantify encoding–retrieval pattern reinstatement (***Nyberg et al., 2000***; ***Manning et al., 2011***), we first extracted 4–128 Hz spectral power in the –0.5–3 s surrounding the encoding and retrieval cues. We excluded the slow theta band (2–4 Hz), as we reported significant differences in this frequency range between conditions. Next, we binned the power data into 250 ms time bins. For each trial pair, we then computed the Pearson's correlation of power measures between encoding and retrieval. The resulting matrix of Pearson's correlation coefficients indicates the strength and direction of spectral similarity between encoding and retrieval for different time-bin combinations. Reinstatement matrices were computed separately for individual trials and averaged at the electrode- and subject levels for

the group analysis. To identify reinstatement patterns that are significantly greater than chance (95% confidence level), we compared the true distributions to surrogate data comprised of 1000 shuffle distributions. At the electrode level, surrogate data involved shuffling trial labels. At the group level, surrogate data involved shuffling block labels.

Acknowledgements

We are thankful to the patients who participated in this study. This project received support from National Institutes of Health grants R01-MH104606 (JJ), R01-NS125250 (BL), and R01-NS107357 (BL).

Additional information

Funding

Funder	Grant reference number	Author
National Institutes of Health	R01-MH104606	Joshua Jacobs
National Institutes of Health	R01-NS125250	Bradley C Lega
National Institutes of Health	R01-NS107357	Bradley C Lega

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

Tamara Gedankien, Data curation, Software, Formal analysis, Investigation, Visualization, Methodology, Writing – original draft, Writing – review and editing; Jennifer Kriegel, Project administration; Erfan Zabeh, Software, Formal analysis, Writing – original draft, Writing – review and editing; David McDonagh, Conceptualization, Investigation, Methodology; Bradley C Lega, Conceptualization, Resources, Funding acquisition, Investigation, Writing – original draft, Writing – review and editing; Joshua Jacobs, Supervision, Funding acquisition, Investigation, Writing – original draft, Writing – review and editing

Author ORCIDs

Tamara Gedankien https://orcid.org/0000-0002-6007-3928
Bradley C Lega https://orcid.org/0000-0003-3634-406X

Ethics

All participants provided their informed written consent, and the study protocol was approved by the Institutional Review Board (IRB) of Human Subjects Research of the UTSW prior to data collection. Intravenous scopolamine was administered by the participant's attending nurse, and a board-certified anesthesiologist was present at the time of injection and remained available throughout the duration of the experiment to treat any potential adverse reactions. Physiological monitoring of blood pressure, electrocardiogram (EKG), oxygen saturation, and mental status was performed just before and for one hour after the injection of scopolamine. Low doses of scopolamine (such as 500 mcg) are commonly administered in clinical settings and generally do not present major risk factors58. There were no reports of adverse events related to substance administration under drug or placebo conditions.

Peer review material

Version of Record:
Reviewer #1 (Public review): https://doi.org/10.7554/eLife.108972.3.sa1
Reviewer #2 (Public review): https://doi.org/10.7554/eLife.108972.3.sa2
Author response https://doi.org/10.7554/eLife.108972.3.sa3

Reviewed preprint v2
Reviewer #1 (Public review): https://doi.org/10.7554/eLife.108972.2.sa2

Reviewer #2 (Public review): https://doi.org/10.7554/eLife.108972.2.sa1
Author response: https://doi.org/10.7554/eLife.108972.2.sa0
eLife Assessment https://doi.org/10.7554/eLife.108972.2.sa3

Reviewed preprint v1
Reviewer #1 (Public review): https://doi.org/10.7554/eLife.108972.1.sa2
Reviewer #2 (Public review): https://doi.org/10.7554/eLife.108972.1.sa1
Author response: https://doi.org/10.7554/eLife.108972.1.sa0
eLife Assessment https://doi.org/10.7554/eLife.108972.1.sa3

Additional files

Supplementary files

Supplementary file 1. Subject demographics and electrode coverage. Table showing subject demographics and number of electrodes across all regions of interest, including the anterior hippocampus (AH), the posterior hippocampus (PC), and the entorhinal cortex (EC).

MDAR checklist

Data availability

The data supporting the findings of this study are openly available in Zenodo at https://doi.org/10.5281/zenodo.23327706. All data analyses were performed in Python (3.6), using publicly available software for the analysis of human electrophysiology data (https://github.com/pennmem/ptsa, ***DePalatis et al., 2026*** and https://github.com/pennmem/cmlreaders, ***DePalatis et al., 2025***). Other custom code is available at https://github.com/tgedankien/AR_scopolamine, copy archived at ***Gedankien, 2025***.

The following dataset was generated:

Author(s)	Year	Dataset title	Dataset URL	Database and Identifier
Gedankien T, Kriegel J, Zabeh E, McDonagh D, Lega B, Jacobs J	2026	Cholinergic blockade reveals a role for human hippocampal theta in memory encoding but not retrieval	https://doi.org/10.5281/zenodo.23327706	Zenodo, 10.5281/zenodo.23327706

References

Aghourian M, Legault-Denis C, Soucy JP, Rosa-Neto P, Gauthier S, Kostikov A, Gravel P, Bédard MA. 2017. Quantification of brain cholinergic denervation in Alzheimer's disease using PET imaging with [^{18}F]-FEOBV. *Molecular Psychiatry* **22**:1531–1538. DOI: https://doi.org/10.1038/mp.2017.183, PMID: 28894304

Atri A, Sherman S, Norman KA, Kirchhoff BA, Nicolas MM, Greicius MD, Cramer SC, Breiter HC, Hasselmo ME, Stern CE. 2004. Blockade of central cholinergic receptors impairs new learning and increases proactive interference in a word paired-associate memory task. *Behavioral Neuroscience* **118**:223–236. DOI: https://doi.org/10.1037/0735-7044.118.1.223, PMID: 14979800

Beatty WW, Bierley RA. 1986a. Scopolamine impairs encoding and retrieval of spatial working memory in rats. *Physiol. Psychol* **14**:82–86.

Beatty WW, Butters N, Janowsky DS. 1986b. Patterns of memory failure after scopolamine treatment: implications for cholinergic hypotheses of dementia. *Behavioral and Neural Biology* **45**:196–211. DOI: https://doi.org/10.1016/s0163-1047(86)90772-7, PMID: 2938571

Bowen DM, Smith CB, White P, Davison AN. 1976. Neurotransmitter-related enzymes and indices of hypoxia in senile dementia and other abiotrophies. *Brain* **99**:459–496. DOI: https://doi.org/10.1093/brain/99.3.459, PMID: 11871

de Chastelaine M, Wang TH, Minton B, Muftuler LT, Rugg MD. 2011. The effects of age, memory performance, and callosal integrity on the neural correlates of successful associative encoding. *Cerebral Cortex* **21**:2166–2176. DOI: https://doi.org/10.1093/cercor/bhq294, PMID: 21282317

DePalatis M, Duey Z, Davis A, Rudoler J, Colyer R. 2025. Cmlreaders. a6f3408. Github. https://github.com/pennmem/cmlreaders

DePalatis M, Sederberg PB, Davis A, Weidemann CT, DeHaan R. 2026. Ptsa. 2d5490a. Github. https://github.com/pennmem/ptsa

Drachman DA, Leavitt J. 1974. Human memory and the cholinergic system: a relationship to aging? arch. *Neurol* **30**:113–121. DOI: https://doi.org/10.1001/archneur.1974.00490320001001
Gallego JA, Perich MG, Miller LE, Solla SA. 2017. Neural manifolds for the control of movement. *Neuron* **94**:978–984. DOI: https://doi.org/10.1016/j.neuron.2017.05.025, PMID: 28595054
Gedankien T, Tan RJ, Qasim SE, Moore H, McDonagh D, Jacobs J, Lega B. 2023. Acetylcholine modulates the temporal dynamics of human theta oscillations during memory. *Nature Communications* **14**:5283. DOI: https://doi.org/10.1038/s41467-023-41025-y, PMID: 37648692
Gedankien T. 2025. AR_scopolamine. swh:1:rev:b47f8ab42ab15863fa504eba54e8be39e80b8f98. Software Heritage. https://archive.softwareheritage.org/swh:1:dir:8c33e148c698a9da5c8a34de812c98683f64264c;origin=https://github.com/tgedankien/AR_scopolamine;visit=swh:1:snp:e8c1f3b3edad3f067928e29ef0f2855f3c97c46e;anchor=swh:1:rev:b47f8ab42ab15863fa504eba54e8be39e80b8f98
Ghoneim MM, Mewaldt SP. 1975. Effects of diazepam and scopolamine on storage, retrieval and organizational processes in memory. *Psychopharmacologia* **44**:257–262. DOI: https://doi.org/10.1007/BF00428903, PMID: 1208760
Griffiths BJ, Mayhew SD, Mullinger KJ, Jorge J, Charest I, Wimber M, Hanslmayr S. 2019. Alpha/beta power decreases track the fidelity of stimulus-specific information. *eLife* **8**:e49562. DOI: https://doi.org/10.7554/eLife.49562, PMID: 31782730
Grober E, Buschke H, Korey SR. 1987. Genuine memory deficits in dementia. *Developmental Neuropsychology* **3**:13–36. DOI: https://doi.org/10.1080/87565648709540361
Grober E, Leipzig RM, Lipton RB, Wisniewski W, Schroeder M, Davies P, Ritter W, Buschke H. 1989. Does scopolamine directly impair memory? *Journal of Cognitive Neuroscience* **1**:327–335. DOI: https://doi.org/10.1162/jocn.1989.1.4.327, PMID: 23971984
Hamburg MD. 1967. Retrograde amnesia produced by intraperitoneal injection of physostigmine. *Science* **156**:973–974. DOI: https://doi.org/10.1126/science.156.3777.973, PMID: 6067162
Hampel H, Mesulam MM, Cuello AC, Farlow MR, Giacobini E, Grossberg GT, Khachaturian AS, Vergallo A, Cavedo E, Snyder PJ, Khachaturian ZS. 2018. The cholinergic system in the pathophysiology and treatment of Alzheimer's disease. *Brain* **141**:1917–1933. DOI: https://doi.org/10.1093/brain/awy132
Hasselmo ME, Bower JM. 1993. Acetylcholine and memory. *Trends in Neurosciences* **16**:218–222. DOI: https://doi.org/10.1016/0166-2236(93)90159-j, PMID: 7688162
Hasselmo ME, Schnell E. 1994. Laminar selectivity of the cholinergic suppression of synaptic transmission in rat hippocampal region CA1: computational modeling and brain slice physiology. *The Journal of Neuroscience* **14**:3898–3914. DOI: https://doi.org/10.1523/JNEUROSCI.14-06-03898.1994, PMID: 8207494
Hasselmo ME, Bodelón C, Wyble BP. 2002. A proposed function for hippocampal theta rhythm: separate phases of encoding and retrieval enhance reversal of prior learning. *Neural Computation* **14**:793–817. DOI: https://doi.org/10.1162/089976602317318965, PMID: 11936962
Hasselmo ME, McGaughy J. 2004. High acetylcholine levels set circuit dynamics for attention and encoding and low acetylcholine levels set dynamics for consolidation. In: Hasselmo ME (Ed). *Progress in Brain Research*. Elsevier. p. 207–231. DOI: https://doi.org/10.1016/S0079-6123(03)45015-2, PMID: 14650918
Hasselmo ME. 2005. What is the function of hippocampal theta rhythm?--Linking behavioral data to phasic properties of field potential and unit recording data. *Hippocampus* **15**:936–949. DOI: https://doi.org/10.1002/hipo.20116, PMID: 16158423
Hasselmo ME. 2006. The role of acetylcholine in learning and memory. *Current Opinion in Neurobiology* **16**:710–715. DOI: https://doi.org/10.1016/j.conb.2006.09.002
Hasselmo ME, Giocomo LM. 2006. Cholinergic modulation of cortical function. *Journal of Molecular Neuroscience* **30**:133–135. DOI: https://doi.org/10.1385/JMN:30:1:133, PMID: 17192659
Hodges JR, Salmon DP, Butters N. 1992. Semantic memory impairment in Alzheimer's disease: failure of access or degraded knowledge? *Neuropsychologia* **30**:301–314. DOI: https://doi.org/10.1016/0028-3932(92)90104-t, PMID: 1603295
Huang ZB, Wang H, Rao XR, Zhong GF, Hu WH, Sheng GQ. 2011. Different effects of scopolamine on the retrieval of spatial memory and fear memory. *Behavioural Brain Research* **221**:604–609. DOI: https://doi.org/10.1016/j.bbr.2010.05.032, PMID: 20553767
Jacobs J, Kahana MJ. 2009. Neural representations of individual stimuli in humans revealed by gamma-band electrocorticographic activity. *The Journal of Neuroscience* **29**:10203–10214. DOI: https://doi.org/10.1523/JNEUROSCI.2187-09.2009, PMID: 19692595
Jutras MJ, Fries P, Buffalo EA. 2013. Oscillatory activity in the monkey hippocampus during visual exploration and memory formation. *PNAS* **110**:13144–13149. DOI: https://doi.org/10.1073/pnas.1302351110, PMID: 23878251
Kaufman MT, Churchland MM, Ryu SI, Shenoy KV. 2014. Cortical activity in the null space: permitting preparation without movement. *Nature Neuroscience* **17**:440–448. DOI: https://doi.org/10.1038/nn.3643, PMID: 24487233
Kilimann I, Grothe M, Heinsen H, Alho EJL, Grinberg L, Amaro Jr. E, dos Santos GAB, da Silva RE, Mitchell AJ, Frisoni GB, Bokde ALW, Fellgiebel A, Filippi M, Hampel H, Klöppel S, Teipel SJ. 2014. Subregional basal forebrain atrophy in Alzheimer's disease: a multicenter study. *Journal of Alzheimer's Disease* **40**:687–700. DOI: https://doi.org/10.3233/JAD-132345
Kunec S, Hasselmo ME, Kopell N. 2005. Encoding and retrieval in the CA3 region of the hippocampus: a model of theta-phase separation. *Journal of Neurophysiology* **94**:70–82. DOI: https://doi.org/10.1152/jn.00731.2004, PMID: 15728768

Kunz L, Staresina BP, Reinacher PC, Brandt A, Guth TA, Schulze-Bonhage A, Jacobs J. 2024. Ripple-locked coactivity of stimulus-specific neurons and human associative memory. *Nature Neuroscience* **27**:587–599. DOI: https://doi.org/10.1038/s41593-023-01550-x, PMID: 38366143

Lega BC, Jacobs J, Kahana M. 2012. Human hippocampal theta oscillations and the formation of episodic memories. *Hippocampus* **22**:748–761. DOI: https://doi.org/10.1002/hipo.20937, PMID: 21538660

Manning JR, Polyn SM, Baltuch GH, Litt B, Kahana MJ. 2011. Oscillatory patterns in temporal lobe reveal context reinstatement during memory search. *PNAS* **108**:12893–12897. DOI: https://doi.org/10.1073/pnas.1015174108, PMID: 21737744

McClelland JL, McNaughton BL, O'Reilly RC. 1995. Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of connectionist models of learning and memory. *Psychological Review* **102**:419–457. DOI: https://doi.org/10.1037/0033-295X.102.3.419, PMID: 7624455

McGaughy J, Koene RA, Eichenbaum H, Hasselmo ME. 2005. Cholinergic deafferentation of the entorhinal cortex in rats impairs encoding of novel but not familiar stimuli in a delayed nonmatch-to-sample task. *The Journal of Neuroscience* **25**:10273–10281. DOI: https://doi.org/10.1523/JNEUROSCI.2386-05.2005, PMID: 16267235

Mewaldt SP, Ghoneim MM. 1979. The effects and interactions of scopolamine, physostigmine and methamphetamine on human memory. *Pharmacology, Biochemistry, and Behavior* **10**:205–210. DOI: https://doi.org/10.1016/0091-3057(79)90088-1, PMID: 450929

Miller KJ, Zanos S, Fetz EE, den Nijs M, Ojemann JG. 2009. Decoupling the cortical power spectrum reveals real-time representation of individual finger movements in humans. *The Journal of Neuroscience* **29**:3132–3137. DOI: https://doi.org/10.1523/JNEUROSCI.5506-08.2009, PMID: 19279250

Moscovitch M, Cabeza R, Winocur G, Nadel L. 2016. Episodic memory and beyond: the hippocampus and neocortex in transformation. *Annual Review of Psychology* **67**:105–134. DOI: https://doi.org/10.1146/annurev-psych-113011-143733, PMID: 26726963

Nyberg L, Habib R, McIntosh AR, Tulving E. 2000. Reactivation of encoding-related brain activity during memory retrieval. *PNAS* **97**:11120–11124. DOI: https://doi.org/10.1073/pnas.97.20.11120, PMID: 11005878

O'Keefe J, Recce ML. 1993. Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus* **3**:317–330. DOI: https://doi.org/10.1002/hipo.450030307, PMID: 8353611

Pereira F, Mitchell T, Botvinick M. 2009. Machine learning classifiers and fMRI: a tutorial overview. *NeuroImage* **45**:S199–S209. DOI: https://doi.org/10.1016/j.neuroimage.2008.11.007, PMID: 19070668

Perri R, Carlesimo GA, Serra L, Caltagirone C, Alberoni M, Appollonio I, Bonaiuto S, Bottini G, Caffarra P, Carlomagno S, Carolei A, Bastiani P, Luca M, Franceschi M, Gallucci M, Gambina G, Ghidoni E, Girotti F, Giubilei F, Lorusso S, et al. 2005. Characterization of memory profile in subjects with amnestic mild cognitive impairment. *Journal of Clinical and Experimental Neuropsychology* **27**:1033–1055. DOI: https://doi.org/10.1080/13803390490919317, PMID: 16273685

Petersen RC. 1977. Scopolamine induced learning failures in man. *Psychopharmacology* **52**:283–289. DOI: https://doi.org/10.1007/BF00426713

Polyn SM, Natu VS, Cohen JD, Norman KA. 2005. Category-specific cortical activity precedes retrieval during memory search. *Science* **310**:1963–1966. DOI: https://doi.org/10.1126/science.1117645, PMID: 16373577

Qasim SE, Fried I, Jacobs J. 2021. Phase precession in the human hippocampus and entorhinal cortex. *Cell* **184**:3242–3255. DOI: https://doi.org/10.1016/j.cell.2021.04.017, PMID: 33979655

Ranganath C, Ritchey M. 2012. Two cortical systems for memory-guided behaviour. *Nature Reviews Neuroscience* **13**:713–726. DOI: https://doi.org/10.1038/nrn3338

Renner UD, Oertel R, Kirch W. 2005. Pharmacokinetics and pharmacodynamics in clinical use of scopolamine. *Therapeutic Drug Monitoring* **27**:655–665. DOI: https://doi.org/10.1097/01.ftd.0000168293.48226.57, PMID: 16175141

Ritchey M, Montchal ME, Yonelinas AP, Ranganath C. 2015. Delay-dependent contributions of medial temporal lobe regions to episodic memory retrieval. *eLife* **4**:e05025. DOI: https://doi.org/10.7554/eLife.05025, PMID: 25584461

Roy R, Niccolini F, Pagano G, Politis M. 2016. Cholinergic imaging in dementia spectrum disorders. *European Journal of Nuclear Medicine and Molecular Imaging* **43**:1376–1386. DOI: https://doi.org/10.1007/s00259-016-3349-x, PMID: 26984612

Rugg MD, Vilberg KL. 2013. Brain networks underlying episodic memory retrieval. *Current Opinion in Neurobiology* **23**:255–260. DOI: https://doi.org/10.1016/j.conb.2012.11.005, PMID: 23206590

Rutishauser U, Ross IB, Mamelak AN, Schuman EM. 2010. Human memory strength is predicted by theta-frequency phase-locking of single neurons. *Nature* **464**:903–907. DOI: https://doi.org/10.1038/nature08860, PMID: 20336071

Sabri O, Meyer PM, Gräf S, Hesse S, Wilke S, Becker GA, Rullmann M, Patt M, Luthardt J, Wagenknecht G, Hoepping A, Smits R, Franke A, Sattler B, Tiepolt S, Fischer S, Deuther-Conrad W, Hegerl U, Barthel H, Schönknecht P, et al. 2018. Cognitive correlates of α4β2 nicotinic acetylcholine receptors in mild Alzheimer's dementia. *Brain* **141**:1840–1854. DOI: https://doi.org/10.1093/brain/awy099, PMID: 29672680

Sederberg PB, Kahana MJ, Howard MW, Donner EJ, Madsen JR. 2003. Theta and gamma oscillations during encoding predict subsequent recall. *The Journal of Neuroscience* **23**:10809–10814. DOI: https://doi.org/10.1523/JNEUROSCI.23-34-10809.2003, PMID: 14645473

Staresina BP, Michelmann S, Bonnefond M, Jensen O, Axmacher N, Fell J. 2016. Hippocampal pattern completion is linked to gamma power increases and alpha power decreases during recollection. *eLife* **5**:e17397. DOI: https://doi.org/10.7554/eLife.17397, PMID: 27508355

Summers WK, Majovski LV, Marsh GM, Tachiki K, Kling A. 1986. Oral tetrahydroaminoacridine in long-term treatment of senile dementia, Alzheimer type. *The New England Journal of Medicine* **315**:1241–1245. DOI: https://doi.org/10.1056/NEJM198611133152001, PMID: 2430180

Tallon-Baudry C, Bertrand O, Delpuech C, Pernier J. 1996. Stimulus specificity of phase-locked and non-phase-locked 40 Hz visual responses in human. *The Journal of Neuroscience* **16**:4240–4249. DOI: https://doi.org/10.1523/JNEUROSCI.16-13-04240.1996, PMID: 8753885

Tang YI, Mishkin M, Aigner TG. 1997. Effects of muscarinic blockade in perirhinal cortex during visual recognition. *PNAS* **94**:12667–12669. DOI: https://doi.org/10.1073/pnas.94.23.12667

Teles-Grilo Ruivo LM, Mellor JR. 2013. Cholinergic modulation of hippocampal network function. *Frontiers in Synaptic Neuroscience* **5**:2. DOI: https://doi.org/10.3389/fnsyn.2013.00002, PMID: 23908628

Ter Wal M, Linde-Domingo J, Lifanov J, Roux F, Kolibius LD, Gollwitzer S, Lang J, Hamer H, Rollings D, Sawlani V, Chelvarajah R, Staresina B, Hanslmayr S, Wimber M. 2021. Theta rhythmicity governs human behavior and hippocampal signals during memory-dependent tasks. *Nature Communications* **12**:7048. DOI: https://doi.org/10.1038/s41467-021-27323-3, PMID: 34857748

Tröster AI, Beatty WW, Staton RD, Rorabaugh AG. 1989. Effects of scopolamine on anterograde and remote memory in humans. *Psychobiology* **17**:12–18. DOI: https://doi.org/10.3758/BF03337812

van Diepen RM, Mazaheri A. 2018. The caveats of observing inter-trial phase-coherence in cognitive neuroscience. *Scientific Reports* **8**:2990. DOI: https://doi.org/10.1038/s41598-018-20423-z, PMID: 29445210

Weng PH, Chen JH, Chen TF, Sun Y, Wen LL, Yip PK, Chu YM, Chen YC. 2013. CHRNA7 polymorphisms and response to cholinesterase inhibitors in Alzheimer's disease. *PLOS ONE* **8**:e84059. DOI: https://doi.org/10.1371/journal.pone.0084059, PMID: 24391883

Whitehouse PJ, Price DL, Clark AW, Coyle JT, DeLong MR. 1981. Alzheimer disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Annals of Neurology* **10**:122–126. DOI: https://doi.org/10.1002/ana.410100203, PMID: 7283399

Yoo HB, Umbach G, Lega B. 2021. Neurons in the human medial temporal lobe track multiple temporal contexts during episodic memory processing. *NeuroImage* **245**:118689. DOI: https://doi.org/10.1016/j.neuroimage.2021.118689, PMID: 34742943