


Figure 7-figure supplement 1

A



B

The figure displays six 2x2 ANOVA tables arranged in a 2x3 grid. The top row shows results for the 'Fraction of MCs' and the bottom row shows results for the 'Fraction of MC-odorant pairs'. The columns represent different response types: 'Excitatory or suppressive response', 'Excitatory response', and 'Suppressive response'. The leftmost column is for the 'task' condition, and the rightmost column is for the 'passive' condition. Within each column, the top table is for the 'Ablation' group and the bottom table is for the 'Control' group. The tables show p-values for the main effects of Ablation and Control, and their interaction.

Response Type	Condition	Ablation	Control
Excitatory or suppressive response	task	p < 0.0001	p < 0.0001
	passive	p = 0.1592	p < 0.0001
Excitatory response	task	p = 0.1345	p = 0.0008
	passive	p = 0.9963	p = 0.9705
Suppressive response	task	p < 0.0001	p < 0.0001
	passive	p = 0.4718	p = 0.7066
Excitatory or suppressive response	task	p < 0.0001	p < 0.0001
	passive	p = 0.0145	p < 0.0001
Excitatory response	task	p = 0.0019	p = 0.0044
	passive	p = 0.9995	p = 0.9730
Suppressive response	task	p < 0.0001	p < 0.0001
	passive	p = 0.4015	p = 0.2262