

Neuroscience

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

31 May 2023 (this version)

Posted to bioRxiv

4 April 2023

Sent for peer review

29 March 2023

Somatotopic organization among parallel sensory pathways that promote a grooming sequence in *Drosophila*

Katharina Eichler, Stefanie Hampel , Adrián Alejandro-García, Steven A. Calle-Schuler, Alexis Santana-Cruz, Lucia Kmecova, Jonathan M. Blagburn, Eric D. Hoopfer, Andrew M. Seeds 

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico • Neuroscience Program, Carleton College, Northfield, Minnesota

 (https://en.wikipedia.org/wiki/Open_access)

 (<https://creativecommons.org/licenses/by/4.0/>)

Abstract

Mechanosensory neurons located across the body surface respond to tactile stimuli and elicit diverse behavioral responses, from relatively simple stimulus location-aimed movements to complex movement sequences. How mechanosensory neurons and their postsynaptic circuits influence such diverse behaviors remains unclear. We previously discovered that *Drosophila* perform a body location-prioritized grooming sequence when mechanosensory neurons at different locations on the head and body are simultaneously stimulated by dust (Hampel et al., 2017; Seeds et al., 2014). Here, we identify nearly all mechanosensory neurons on the *Drosophila* head that individually elicit aimed grooming of specific head locations, while collectively eliciting a whole head grooming sequence. Different tracing methods were used to reconstruct the projections of these neurons from different locations on the head to their distinct arborizations in the brain. This provides the first synaptic resolution somatotopic map of a head, and defines the parallel-projecting mechanosensory pathways that elicit head grooming.

eLife assessment

This **valuable** work provides a near-complete description of the mechanosensory bristles on the *Drosophila melanogaster* head and the anatomy and projection patterns of the bristle mechanosensory neurons that innervate them. The data presented are **solid**. The study has generated numerous invaluable resources for the community that will be of interest to neuroscientists in the field of circuits and behaviour, particularly those interested in mechanosensation and behavioural sequence generation.

Introduction

The ability to produce complex behaviors by assembling sequences of different movements is essential for purposeful behavior and survival. One prominent model that describes how

the brain produces movement sequences is called a “parallel model”. This model proposes that the premotor elements of different movements to be executed in sequence are activated (or readied) in parallel and then selected sequentially through a mechanism where movements occurring earlier in the sequence suppress later ones (Bohland et al., 2010; Bullock, 2004; Houghton and Hartley, 1995; Lashley, 1951). A hallmark feature of this model is a parallel circuit architecture that ensures all mutually exclusive actions to be performed in sequence are simultaneously readied and competing for output. Performance order is established by an activity gradient among parallel circuits where earlier actions have the highest activity and later actions have the lowest. A winner-take-all network selects the action with the highest activity and suppresses the others. The selected action is performed and then terminated to allow a new round of competition and selection of the next action. This model is supported by physiological and behavioral evidence from the movement sequences of different animals (Averbeck et al., 2002; Mushiaki et al., 2006; Seeds et al., 2014). Yet, despite some movement sequences exhibiting features consistent with the parallel model, our understanding of the underlying neural circuit mechanisms is limited.

The grooming behavior of fruit flies (*Drosophila melanogaster*) can be studied to define the circuit mechanisms that produce movement sequences. Making flies dirty by coating them in dust elicits a grooming sequence that starts with the cleaning of different locations on the head and proceeds to the abdomen, wings, and thorax (Mueller et al., 2019; Phillis et al., 1993; Seeds et al., 2014). We previously determined that the sequence is produced by a mechanism that is consistent with a parallel model (Seeds et al., 2014). The sequence begins when different aimed grooming movements that clean specific locations of the head or body become activated in parallel by dust. The resulting competition among mutually exclusive grooming movements is resolved through hierarchical suppression. For example, grooming of the eyes occurs first because eye grooming suppresses grooming of other locations on the head and body. This parallel model of hierarchical suppression provides a conceptual framework for dissecting the neural circuit architecture that produces *Drosophila* grooming. Here, we examine the input layer of the hypothesized architecture (Seeds et al., 2014), the mechanosensory neurons that detect dust at different body locations.

Different mechanosensory structures are distributed across the head and body surface that respond to mechanical stimuli and elicit grooming. The most abundant of these structures are mechanosensory bristles (aka hairs or setae). Tactile displacement of individual bristles elicits grooming movements in which the legs are precisely aimed at the stimulus location (Corfas and Dudai, 1989; Page and Matheson, 2004; Vandervorst and Ghysen, 1980). Each bristle is innervated by a single *bristle mechanosensory neuron* (BMN) that is excited by displacement of that bristle (Corfas and Dudai, 1990; Tuthill and Wilson, 2016a; Walker, 2000). Thus, bristles and their corresponding BMNs can be ascribed to specific, aimed leg grooming movements. Other mechanosensory structures, including chordotonal organs and stretch receptors, also elicit stimulus location-aimed grooming (Hampel et al., 2017; Zhang et al., 2020). Simultaneous (parallel) optogenetic activation of mechanosensory neurons across the body elicits a grooming sequence that proceeds in the same order as the “natural” dust-induced sequence (Hampel et al., 2017; Zhang et al., 2020). Thus, the sequence is elicited by parallel mechanosensory pathways that each produce a movement that grooms a specific location on the head or body.

BMNs project their axons from different locations on the head or body, through different nerves, and into the *central nervous system* (CNS). Previous studies demonstrated that BMNs show somatotopic organization in their CNS projections (Johnson and Murphey, 1985; Murphey et al., 1989b; Newland, 1991; Newland et al., 2000; Tsubouchi et al., 2017). That is, particular projection zones in the CNS correspond to specific body locations. Somatotopic organization among mechanosensory neurons and their postsynaptic circuits is consistent with the parallel model that underlies the body grooming sequence (Seeds et al., 2014). In

this model, parallel-projecting mechanosensory neurons that respond to stimuli at specific locations on the head or body could connect with somatotopically-organized parallel circuits that elicit grooming of those locations. However, this circuit architecture has not been verified, in part, because the somatotopy among the mechanosensory neurons has not been comprehensively defined.

Here, we map the somatotopic organization among BMNs that elicit grooming of the head. Flies use their front legs to groom their heads in a sequence that starts with the eyes and proceeds to other locations, such as the antennae and proboscis (Seeds et al., 2014). Two mechanosensory structures on the head (i.e. chordotonal organs and bristles) are implicated in grooming. The antennal *Johnston's organ* (JO) is a chordotonal organ containing mechanosensory neurons called *JO neurons* (JONs) that detect stimulations of the antennae and elicit aimed grooming (Hampel et al., 2015; 2020a). There are over 1,000 bristles located on the head whose stimulation we hypothesized could also elicit aimed grooming responses. In support of this hypothesis, BMNs innervating bristles on the eyes were previously shown to elicit grooming of the eyes (Hampel et al., 2017; Zhang et al., 2020). Here, we produce a map of BMN projections from nearly all bristles on the head. We use transgenic expression, dye fills, and electron microscopy (EM) reconstructions to trace the projections of the BMNs, from their bristles, through their respective nerves, and into the CNS. Using optogenetic tools, we show that activation of subsets of BMNs at specific head locations elicits aimed grooming. Thus, we produce a synaptic resolution map of the parallel-projecting mechanosensory neurons that elicit head grooming.

Results

Classification and quantification of the head bristles

A prerequisite for determining the somatotopy of head BMNs was to define the locations of their respective bristles on the head. Different populations of bristles are located on the eyes, antennae, proboscis, and other areas on the head. While the identities of most of these populations were known (Bodenstein et al., 1994), some were poorly described and their bristle numbers were not reported. Therefore, we imaged the bristles on the head and then classified and quantified each population. We developed a unified nomenclature for the different bristle populations that was based partially on published nomenclature. Most of the bristles were easily observed by imaging white light-illuminated heads (**Figure 1A-D**), and color-coded depth maps further helped to distinguish between bristles while they were being counted (**Figure 1 – figure supplement 1A-H**). Some bristles could not be counted from these images because of their small size, position on the head, or because they could not be distinguished from one another (**Figure 1E**, asterisk with bristle number range). Therefore, we used confocal microscopy images, or referred to published work to estimate or obtain the numbers of bristles in these populations (see Materials and methods).




Figure 1.

Classification and quantification of head bristles.

(A-D) Bristles on the anterior (A), posterior (B), ventral (C), and dorsal (D) male head. The bristles on the right half are marked with color-coded dots to indicate their classification. Bristle names are abbreviated (Abv.), and full names and color codes are listed in (E). (E) Quantification of bristle populations on the head (per half). Range indicates the lowest and highest number of bristles counted across individuals for each population (N=8). Bristle number average (Avg.) and standard deviation (SD) across individuals for each population are shown. Bristle counting was facilitated using color-coded depth maps (examples shown in Figure 1 – figure supplement 1). Quantification of bristles on female heads and male/female comparisons are shown in Figure 1 – figure supplement 2. See Supplementary file 1 for bristle counts for each head. *InOm and Taste bristle number ranges are based on published data while dPoOr, vPoOr, and vOcci bristles were counted using confocal microscopy (see Materials and methods). Bristles are organized into nerve groups based on the nerve each bristle’s corresponding bristle mechanosensory neuron (BMN) projects through to enter the brain (evidence shown in Figure 2). Dorsal (d) and Ventral (v).

We next produced a map of the different bristles at their stereotyped locations on the head, and determined how the numbers of bristles in each population varied across individual flies (Figure 1A-E). By counting the bristles on both male and female heads, we found no significant gender-based differences in their numbers (Figure 1 – figure supplement 2A-E, Supplementary file 1). Given that the bristles are singly innervated (Tuthill and Wilson, 2016b), we could use the bristle counts to estimate the number of BMNs for each bristle population. This provided a framework for us to define the somatotopic projections of BMNs that innervate particular bristles.

Light microscopy-based reconstruction of BMNs innervating the head bristles

BMNs project from bristles at specific head locations and then through their respective nerves to enter the brain. While the nerve projections of BMNs innervating bristles on the eyes, proboscis, and antennae were previously reported (Hampel et al., 2017; Homberg et al., 1989; Melzig et al., 1996; Singh and Nayak, 1985; Stocker, 1994), the projections of BMNs innervating other head bristles were unknown. We determined these projections using a transgenic driver line (R52A06-GAL4) that labels BMNs on the head (Hampel et al., 2017). R52A06-GAL4 was used to express membrane-targeted *green fluorescent protein* (mCD8::GFP), and the anterior and posterior head was imaged with a confocal microscope. The GFP-labeled neurons had all the characteristic morphological features of BMNs (Tuthill and Wilson, 2016b), including a dendrite innervating a bristle, a cell body, and an axon (Figure 2A,B). R52A06-GAL4 labeled almost all BMNs on the head, but did not label any associated with the *postocellar* (PoOc) or *supracervical* (Su) bristles (Figure 2 – figure supplement 1A-H). We used the software neuTube (Feng et al., 2015) to reconstruct the GFP-labeled projections of head BMNs from confocal z-stacks (Figure 2C-H). The reconstructions enabled us to classify the BMNs into ‘nerve groups’, based on the nerves they project through to enter the brain (Figure 2I,J, groups listed in Figure 1E). This revealed that BMNs innervating bristles at different locations on the head project through specific nerves, including the antennal, eye, occipital, and labial nerves. Below we introduce the BMNs in each nerve group and the bristles that they innervate.




Figure 2.

BMNs on the head project through specific nerves.

(A–D) Confocal z-stack maximum intensity projections of the anterior (A,C) and posterior (B,D) head in which the driver line R52A06-GAL4 drives expression of GFP in BMNs (green). Cuticle is magenta. (A,B) Magnified views of the boxed areas indicated in C and D. The *dendrite* (De), *axon* (Ax), *cell body* (CB), and *innervated bristle* (Br) of a BMN are indicated in each panel. (C,D) The left half of the head is shown as a maximum projection, while z-stack-reconstructed BMNs are shown for the right half. Maximum projections of the right half of the head is shown in Figure 2 – figure supplement 1A–F. (E–H) Magnified images of the reconstructions. The magnified areas are indicated by vertical lines on the right in C and D. Reconstructed BMNs are color-coded and labeled according to the nerve that they project through: AntNv (blue); OcciNv (green); EyeNv (red); LabNv (brown). Unreconstructed portion of the antennal nerve is indicated by an asterisk. Innervated bristles are indicated with black arrows. Black arrowheads in F and H indicate partially reconstructed axons of BMNs innervating the InOm, Vib, and Taste bristles. Scale bars: 25 μ m (B), 100 μ m (D). (I,J) Summary of bristles innervated by BMNs that belong to particular nerve groups on the anterior (I) and posterior (J) head. Nerve groups also listed in Figure 1E.

reconstructed BMNs are shown for the right half. Maximum projections of the right half of the head is shown in Figure 2 – figure supplement 1A–F. (E–H) Magnified images of the reconstructions. The magnified areas are indicated by vertical lines on the right in C and D. Reconstructed BMNs are color-coded and labeled according to the nerve that they project through: AntNv (blue); OcciNv (green); EyeNv (red); LabNv (brown). Unreconstructed portion of the antennal nerve is indicated by an asterisk. Innervated bristles are indicated with black arrows. Black arrowheads in F and H indicate partially reconstructed axons of BMNs innervating the InOm, Vib, and Taste bristles. Scale bars: 25 μ m (B), 100 μ m (D). (I,J) Summary of bristles innervated by BMNs that belong to particular nerve groups on the anterior (I) and posterior (J) head. Nerve groups also listed in Figure 1E.

Head BMNs project to the brain through specific nerves

BMNs innervating the 18-22 *antennal* (Ant) bristles were previously reported to project through the *antennal nerve* (AntNv) that also carries the axons of JONs and olfactory neurons (Homborg et al., 1989; Melzig et al., 1996). We identified additional BMNs projecting through the AntNv that innervate bristles located on the anterior and dorsal head (**Figure 2E,I**, blue). These include 4-7 *frontal* (Fr) bristles located medially, 3 *orbital* (Or) and 4-6 *frontoorbital* (FrOr) bristles located laterally, and 1 *ocellar* (Oc) and 3-4 *interocellar* (InOc) bristles located on the dorsal head. BMNs projecting from these bristles form a bundle below the cuticle that projects ventrally to join the AntNv. We also identified BMNs that innervate 1-3 of the small anterior *vibrissae* (Vib) on the ventral head whose axons project dorsally to join the AntNv.

BMNs innervating bristles on the dorsal half of the posterior head project through a previously undescribed nerve that we named the *occipital nerve* (OcciNv) (**Figure 2F,J**, green). This includes the 3-4 *vertical* (Vt), 2-5 *dorsal occipital* (dOcci), and 5-9 *dorsal postorbital* (dPoOr) bristles. BMNs that innervate these different bristles form the OcciNv that projects under the cuticle ventromedially towards the brain.


Each eye contains between 645 and 828 regularly spaced ommatidia, many of which have an associated *interommatidial* (InOm) bristle (Ready et al., 1976). We estimated that there are between 607 and 645 InOm bristles on each eye based on published data (see Materials and methods). BMNs that innervate the InOm bristles were previously found to form a nerve that projects to the brain from the posterior head (Hampel et al., 2017). Because this nerve was not previously named, it is referred to here as the *eye nerve* (EyeNv). We found that the EyeNv also carries the projections of BMNs innervating bristles on the posterior and ventral head (**Figure 2E-J**, red). Those on the posterior head innervate the 12-18 *ventral occipital* (vOcci) and 9-15 *ventral postorbital* (vPoOr) bristles. Those on the ventral head innervate most of the 13-18 Vib bristles.

The proboscis has bristles on the labellum, haustellum, and maxillary palps. Each half of the labellum has 31-42 *Taste* bristles whose associated BMNs project through the *labial nerve* (LabNv) (Falk et al., 1976; Jeong et al., 2016; Nayak and Singh, 1983; Shanbhag et al., 2001; Stocker, 1994). The LabNv also carries mechanosensory neurons innervating the labellar taste pegs, along with gustatory neurons innervating either the taste pegs or taste bristles (Stocker and Schorderet, 1981). We found that BMNs innervating the 5 *haustellum* (Hau) bristles also project through the LabNv (**Figure 2H,J**, brown). BMNs that innervate the 14-18 *maxillary palp* (MaPa) bristles project through the *maxillary nerve* (MaxNv) that also carries the axons of olfactory neurons (Singh and Nayak, 1985). The Lab- and MaxNvs merge as they approach the head, and in this work we refer to the merged nerve as the LabNv. The LabNv then merges with the EyeNv in the ventral head, suggesting that these nerves project into the brain at the same location (**Figure 2H**).

Head BMNs project into discrete zones in the ventral brain

BMNs in the Ant-, Eye-, and LabNvs were previously reported to project into a region of the ventral brain called the *subesophageal zone* (SEZ) (**Figure 3A,B**) (Hampel et al., 2017; Jeong et al., 2016; Kamikouchi et al., 2006; Mitchell et al., 1999; Singh and Nayak, 1985; Stocker, 1994). To determine if all head BMNs project into the SEZ, we used R52A06-GAL4 to label their projections in a dissected brain (**Figure 3C**). The AntNv was identified in the R52A06-GAL4 pattern based on its reported dorsal-arriving projection into the SEZ (Kamikouchi et al., 2006; Stocker, 1994), while the Eye- and LabNvs were identified based on their reported ventral-arriving projections (Hampel et al., 2017; Stocker, 1994). We found that the Eye- and LabNvs project into the ventral SEZ at the same location (**Figure 3C**), consistent with the observation

that they merge as they approach the brain (**Figure 2F,H**). We tentatively identified the OcciNv projecting into the SEZ from a lateral direction, revealing that all head BMN nerves project into the SEZ. R52A06-GAL4 also labels the antennal chordotonal JONs that are known to project through the AntNv into a dorsal area of the SEZ (Hampel et al., 2017; 2020a; Kamikouchi et al., 2006; Kim et al., 2020). Visualization of JONs and BMNs in the same expression pattern revealed that most of the BMNs project into more ventral regions of the SEZ than the JONs (**Figure 3C**).



Reconstructed BMNs on half of the head that are labeled by the following driver lines: InOmBMN-LexA (**D**), dBMN-spGAL4 (**E**), pBMN-spGAL4 (**F**), and TasteBMN-spGAL4 (whole proboscis shown) (**G**). Images of the heads used for each reconstruction are shown in **Figure 3 – figure supplement 1A-D**. Reconstructed neurons are color-coded and labeled as described in **Figure 2**. (**D'-G'**) SEZ projections of BMNs from both halves of the head that are labeled by InOmBMN-LexA (**D'**), dBMN-spGAL4 (**E'**), pBMN-spGAL4 (**F'**), and TasteBMN-spGAL4 (**G'**). (**H**) Table of BMNs innervating specific bristles that are labeled by each driver line, indicated by box shading (numbers of labeled BMNs innervating different bristles shown in **Figure 3 – figure supplement 1E**). Shade color indicates the nerve that each BMN projects through. (**I**) Driver line names and identifiers. (**J**) Aligned expression patterns of InOmBMN-LexA (red), dBMN-spGAL4 (green), and TasteBMN-spGAL4 (brown).

We next used different transgenic driver lines that express in specific populations of head BMNs to independently label and visualize the different nerves (**Figure 3D-I**, **Figure 3 – figure supplement 1A-D**). The EyeNv was labeled using a previously identified driver line (VT017251-LexA) that expresses in BMNs innervating the InOm bristles (**Figure 3D**) (Hampel et al., 2017). Here, we refer to this line as InOmBMN-LexA. We also used a screening approach to produce three new Split GAL4 (spGAL4) combinations that express in BMNs innervating bristles at other locations on the head (see Materials and methods). One line named dBMN-spGAL4 labels BMNs innervating some dorsally located bristles (InOc, Vt, and dPoOr) that project through the Ant- and OcciNvs (**Figure 3E**). Another line named pBMN-spGAL4 labels BMNs innervating bristles on the posterior head (Vt, dOcci, dPoOr, and vOcci) that project through the Occi- and EyeNvs (**Figure 3F**). The third line named TasteBMN-

Figure 3.

Head BMNs project into the ventral brain region called the subesophageal zone (SEZ).

(**A**) Schematic of BMNs projecting from different nerves into the SEZ. (**B**) Anterior view of the brain immunostained for Bruchpilot (magenta) to visualize the neuropile. White box indicates the SEZ. Scale bar, 100 μm. (**C**) Image of the SEZ in which R52A06-GAL4 expressed GFP in BMNs and JONs. Brains were immunostained for GFP (green) and Bruchpilot (magenta). BMN nerves and JONs are labeled. Scale bar, 25 μm. (**D-G**) Driver lines that label BMNs from different nerves.

Reconstructed BMNs on half of the

spGAL4 labels BMNs innervating Taste bristles on the labellum that project through the LabNv (**Figure 3G**). These driver lines each provided independent labeling of one or two different nerves (**Figure 3H**).

Consistent with what we observed using R52A06-GAL4, each driver line labeled BMNs that projected into the SEZ and no other areas of the brain or *ventral nerve cord* (VNC) (**Figure 3D'-G'**, **Figure 3 – figure supplement 1A'-D'**). dBMN-spGAL4 and pBMN-spGAL4 both labeled the OcciNv that was found to project into the SEZ from a lateral direction (**Figure 3E',F'**), in agreement with what we observed in the R52A06-GAL4 pattern (**Figure 3C**). A comparison of the nerves labeled by the different driver lines revealed that each nerve has morphologically distinct projections. To further visualize the spatial relationships between these projections, we computationally aligned the expression patterns of the different driver lines into the same brain space (**Figure 3J**). Indeed, BMNs from different nerves were found to project into distinct zones of the ventral SEZ. However, we also observed potential zones where overlap could occur between the projections of BMNs from different nerves (discussed more below).

Brain projections of BMNs that innervate specific head bristles

Our results suggested that different BMN “types” innervate specific populations of bristles on the head and project into distinct zones in the SEZ. However, it was unclear to what extent BMNs of the same type projected to the same zones, and if other BMN types had distinct or overlapping projections. Therefore, we next compared the projections of individual BMNs from different populations of bristles.

The head contains different sized bristles, ranging from large Vt bristles on the dorsal head, to small vOcci bristles on the posterior head. We performed dye fills to label individual BMNs that innervate the largest bristles. This was done by modifying a previously published method for filling BMNs innervating bristles on the thorax (Kays et al., 2014). In the modified method, a particular bristle was plucked from the head and a small volume of dye (DiD) pipetted into the exposed socket containing the dendrite of the associated BMN. The dye then diffused into the neuron, and its projection morphology in the brain was imaged using a confocal microscope (experiment schematic and example fills shown in **Figure 4 – figure supplement 1A-E**). This method was particularly amenable to large bristles that were relatively easy to pluck. We successfully filled individual BMNs that innervate the Oc, Or, Ant, Vib, and Vt bristles (**Figure 4C-Q**). The BMNs were named based on the bristle populations that they innervate. For example, BMNs that innervate the Ant bristles were named *bristle mechanosensory Ant neurons* (BM-Ant neurons).




Figure 4.

Projections of BMNs that innervate specific head bristles.

(A,B) Bristles on the anterior (A) and posterior (B) head whose associated BMNs were labeled using dye fill (C–Q, fill) or multicolor flipout (R–V, MCFO) techniques. (C–V) SEZ projections of individual BMNs that innervate the bristle indicated in the upper right corner (anterior view). BMNs are oriented as if they are projecting from the right side of the head. Scale bar, 50 μ m. (C–Q) BMNs labeled by dye filling. Schematic of the filling technique and whole brain examples shown in Figure 4 – figure supplement 1. Filled BMNs innervate the Oc (C), Or (D–F), Ant (G–J), Vib (K–N), and Vt (O–Q) bristles. All fill trials for the different bristles are shown in Figure 4 – figure supplement 2, Figure 4 – figure supplement 3, Figure 4 – figure supplement 4, and Figure 4 – figure supplement 5. (R–V) MCFO-labeled BMNs innervate the InOc (R), dPoOr (S), dOcci/dPoOr (T), vOcci (U), and Taste (V) bristles. BMNs were MCFO labeled using the following driver lines: dBMN-spGAL4 (R,S), pBMN-spGAL4 (T,U), and TasteBMN-spGAL4 (V). All MCFO trials for the different bristles are shown in Figure 4 – figure supplement 6, Figure 4 – figure supplement 7, and Figure 4 – figure supplement 8. The number (N) of fill or MCFO trials obtained for each BMN is indicated in the upper right corner.

The large bristles are invariant in number and location across individuals (Figure 1A–E). For example, all flies have a single Oc bristle on each half of the head that is always in the same location. We therefore performed dye fills on the same bristles from multiple different heads. This revealed that BMNs innervating the same bristle have the same general projection morphology across individual flies (Figure 4 – figure supplements 2–5). We also performed dye fills on different bristles from the same population, such as the Ant 1, Ant 2, Ant 3, and Ant 4 bristles (Figure 4 – figure supplement 3A–M). BMNs innervating the same populations were found to have similar projections. For example, BM-Ant neurons all showed similar ipsilateral and midline projecting branches (Figure 4G–J). Morphological similarity among BMNs innervating the same bristle populations was also observed for the BM-Or (Figure 4D–F), –Vib (Figure 4K–N), and –Vt (Figure 4O–Q) neurons.

While BMNs innervating the largest bristles could be labeled using dye fills, we could not label BMNs innervating small bristles using this method. Therefore, we used the *multicolor flipout* (MCFO) method (Nern et al., 2015) to stochastically label individual BMNs innervating

bristles within the expression patterns of the driver lines shown in [Figure 3E'-G'](#). This enabled us to determine the morphologies of BMNs that innervate the InOc, dOcci, dPoOr, vOcci, and Taste bristles ([Figure 4R-V](#)). Unlike the dye-filled BMNs, the MCFO-labeled BMNs could not be linked to specific bristles within a population (e.g. Ant 1 or Ant 2), but only to a specific population (e.g. Ant). In agreement with what we observed with dye-filled BMNs innervating the same populations of large bristles, the MCFO-labeled BMNs innervating the same populations of small bristles also showed similar projection morphologies ([Figure 4 – figure supplements 6–8](#)).

We next compared the projections of the dye-filled and MCFO-labeled BMNs ([Figure 4C-V](#)). This revealed that some BMNs innervating neighboring bristle populations have similar morphologies. For example, BM-InOc and -Oc neurons have similar morphology, including ipsilateral and midline-crossing projections ([Figure 4C,R](#)), while BM-dPoOr, -dOcci, and -vOcci neurons show similar ipsilateral projections. This suggested that BMNs innervating neighboring head bristle populations show similar morphology and project into overlapping zones in the SEZ.

EM-based reconstruction of the head BMN projections in a full adult brain

We next used a previously reported serial-section EM volume of a *full adult fly brain* (FAFB) to reconstruct the SEZ projections of all head BMNs and produce a comprehensive map of their organization ([Zheng et al., 2018](#)). FAFB consists of a brain that was dissected from the head capsule, making it impossible to reconstruct BMNs all the way from their bristles. Instead, the severed Ant-, Occi-, Eye-, and LabNvs were identified in FAFB at the same anatomical locations that we had observed using light microscopy ([Figure 3C,J](#), [Figure 5 – figure supplement 1A](#)). We used the FlyWire.ai platform ([Dorkenwald et al., 2022](#)) to seed all automatically-segmented neurons within the different nerve bundles as they entered the neuropil (right brain hemisphere nerves, [Figure 5 – figure supplement 1B-D](#)), and the neurons were then fully proofread and edited by human experts to identify their individual morphologies. The morphologies of the majority of the reconstructed neurons matched those of mechanosensory neurons, including BMNs (discussed below), JONs ([Hampel et al., 2020a](#); [Kamikouchi et al., 2006](#); [Kim et al., 2020](#)), and labellar taste peg mechanosensory neurons (TPMNs) ([Jeong et al., 2016](#); [Miyazaki and Ito, 2010](#); [Zhou et al., 2019](#)) ([Figure 5 – figure supplement 1E](#)). The remaining neurons included gustatory neurons ([Engert et al., 2022](#)), unidentified sensory neurons ([Figure 5 – figure supplement 2A-Y](#)), and interneurons (not shown).

We identified 705 BMNs among the EM-reconstructed neurons by comparing their SEZ projection morphologies with light microscopy imaged BMNs ([Figure 3C,J](#), [Figure 4C-V](#)). In agreement with the light microscopy data, the reconstructed BMNs project through different nerves into distinct zones in the SEZ ([Figure 5A,B](#), [FlyWire.ai link 1](#)). For example, BMNs from the Eye- and LabNv have distinct ventral and anterior projections, respectively. This shows how the BMNs are somatotopically organized, as their distinct projections correspond to different bristle locations on the head ([Figure 5B,C](#)).




Figure 5.

Electron microscopy-based reconstruction of head BMNs.

(A) All reconstructed BMNs projecting into the brain from the right side of the head (anterior, dorsal, and lateral views shown). BMN colors correspond to the nerves that they project through, including the AntNv (blue), EyeNv (red), OcciNv (green), and LabNv (brown). Scale bars, 50 μ m. (B) Zoomed anterior (left) and lateral (right) views of the BMNs in the SEZ. Labeled arrows for each incoming nerve indicate BMN projection direction. Scale bars, 10 μ m. (C) Bristles on the anterior (left) and posterior (right) head that are innervated by BMNs in the nerve groups indicated by their color. [Figure 5 – figure supplement 1](#) summarizes the EM reconstruction strategy. Sensory neurons that could not be assigned an identity are shown in [Figure 5 – figure supplement 2](#).

Matching the reconstructed head BMNs with their bristles

The reconstructed BMN projections were next matched with their specific bristle populations. The projections were first clustered based on morphological similarity using the NBLAST algorithm ([Figure 6 – figure supplement 1A,B](#), [Supplementary file 2](#), [FlyWire.ai link 2](#)) (Costa et al., 2016). These clusters were then assigned as BMN types based on their similarity to light microscopy images of BMNs known to innervate specific bristles. 10 types were matched with dye-filled or MCFO-labeled BMNs (BM-InOc, -Oc, -Ant, -Or, -Vib, -Vt, -dPoOr, -dOcci, vOcci, and -Taste neurons, [Figure 6 – figure supplement 2A-M](#)). BM-MaPa neurons were matched using published images of labeled MaxNv projections (Singh and Nayak, 1985). 4 types were matched by comparison with BMNs innervating neighboring bristles that showed similar morphology (BM-Fr, -FrOr, -vPoOr, and -Hau neurons). Among these, the BM-vPoOr neurons were so morphologically similar to the MCFO matched BM-vOcci neurons that they could not be distinguished from each other, and were therefore treated as a single group (BM-vOcci/vPoOr neurons). The collective projections of the 555 reconstructed BM-InOm neurons were matched with BMNs labeled using the InOmBMN-LexA driver line ([Figure 6 – figure supplement 2N](#)). This matching involved combining 11 different NBLAST clusters ([Figure 6 – figure supplement 1A,B](#)) and revealed morphological diversity among the BM-InOm neurons.

Additional evidence was used to support our BMN type assignments (**Figure 6 – figure supplement 3A**), including a comparison of the morphology and numbers of reconstructed BMNs on both sides of the brain (for the small OcciNv, **Figure 6 – figure supplement 4A-D**), and determining that BMNs of the same type show common postsynaptic connectivity (described below). Finally, we verified that the numbers of BMNs for each type were consistent with their corresponding bristle numbers (**Figure 6 – figure supplement 3B-F**). This consistency of the BMN/bristle numbers, and completeness of sensory neuron proofreading in each nerve suggested that nearly all BMNs were reconstructed. Thus, we produced a near complete brain projection map of 15 BMN types that innervate the different bristle populations on the head (**Figure 6A-R**, listed in **Supplementary file 2**, FlyWire.ai link 3).




Figure 6.

BMN types that innervate specific head bristles.


(A-B) Different bristle populations indicated by labeled and colored dots are innervated by BMNs shown in C-R. The anterior (A) and posterior (B) head are shown. (C-R) Reconstructed SEZ projections of BMN types that are labeled and plotted in colors indicating the bristles that they innervate. Shown are the dorsal views of all BMNs (C), BM-InOc (D), BM-Oc (E), BM-Fr (F), BM-Ant (G), BM-Or (H), BM-FrOr (I), BM-InOm (J), BM-Vib (K), BM-MaPa (L), BM-Taste (M), BM-Hau (N), BM-Vt/PoOc (O), BM-dOcci (P), BM-dPoOr (Q), and BM-vOcci/vPoOr (R) neurons. The number of reconstructed BMNs for each type is indicated.

Evidence used for assigning the different BMN types is shown in **Figure 6 – figure supplements 1**, **Figure 6 – figure supplements 2**, **Figure 6 – figure supplements 3**, and **Figure 6 – figure supplements 4**.

BMN somatotopic map

The projection map defined above revealed three features of somatotopic organization among the BMN types (**Figure 6C-R**). First, each type has a unique branch morphology that defines its projections into distinct zones in the SEZ. Second, types that innervate neighboring bristle populations have branches that project into partially overlapping zones. For example, BMNs that innervate bristles on the dorsal head all have a common ipsilateral projection (**Figure 6C-I,O**, lateral branch in each panel). In contrast, BMNs that innervate bristles at distant locations (e.g. dorsal and ventral head) show little or no projection overlap. Third, the projections of BMNs either remain in the ipsilateral brain hemisphere or cross the midline to the contralateral side, depending on the locations of their corresponding bristle populations (**Figure 7A-E**). That is, BMNs innervating populations located medially on

the anterior head have midline-crossing projections, whereas BMNs innervating lateral, eye, and posterior head populations have ipsilateral-only projections. BMNs innervating bristles on the proboscis showed mixtures of ipsilateral-only and midline-crossing projections. These somatotopic features reveal how BMNs have distinct and overlapping SEZ projections that reflect their relative locations and proximities on the head.



bristles at distant locations showed low NBLAST similarity and were not connected, consistent with these BMNs projecting into distinct zones. Interestingly, the different BMN projection zones defined by the NBLAST and connectivity data correspond roughly to the eye, ventral, dorsal, and posterior head.

Somatotopically-organized parallel BMN pathways

The map of somatotopically organized BMN projection zones provided evidence of the parallel sensory pathways predicted by the model of hierarchical suppression underlying grooming (Hampel et al., 2017; Seeds et al., 2014). In the model, sensory neurons detect dust at different head locations and elicit aimed grooming through distinct postsynaptic circuits that function in parallel. The projection zones could be where BMNs synapse with these circuits. We examined the postsynaptic connectivity of the different BMN types to test if they form parallel connections with distinct partners. Nearly all neurons postsynaptic to the BMNs were first proofread in FlyWire.ai by our group and the wider proofreading community (Dorkenwald et al., 2022). We then compared the connectivity of the BMNs with their postsynaptic partners using cosine similarity-based clustering.

We performed cosine similarity clustering of the BMNs without the BM-InOm neurons. This was because the 555 BM-InOm neurons were present in higher numbers and with fewer presynaptic sites than the 150 BMNs of other types, and clustering all BMNs together resulted in obscured clustering (presynaptic site counts in [Figure 7 – figure supplement 1A](#)). Indeed, the resulting clusters revealed parallel postsynaptic connectivity that reflected the BMN somatotopic map. Examples are shown in [Figure 7F–L](#) as plots of BMN clusters from the dendrogram in [Figure 7 – figure supplement 2](#) (cut height 4.5, FlyWire.ai link 4). The lowest level clusters included BMNs of the same type that were strikingly clustered together, and thus have the highest postsynaptic connectivity similarity ([Figure 7H,I](#), BM-Vib and -Taste neuron clusters shown). Some BMNs of the same type fell into more than one cluster, suggesting that there are BMN subtypes that connect with different postsynaptic partners ([Figure 7I,K](#), BM-Taste (yellow); [Figure 7J,L](#), BM-Ant (teal)). This intratype clustering was also observed among the morphologically diverse BM-InOm neurons when they were compared with each other ([Figure 7M](#), clustering shown in [Figure 7 – figure supplement 3](#), FlyWire.ai link 5). At higher levels of clustering, BMN types innervating neighboring bristle populations showed high similarity, falling into clusters that corresponded to the dorsal, ventral, and posterior head ([Figure 7J,K](#), dorsal and ventral head shown). Thus, the somatotopic organization of the BMN projections is preserved among their postsynaptic partners to form parallel sensory pathways. However, we also found evidence that some postsynaptic partners might not be head location specific, as one cluster included BMNs that innervate bristles on the anterior and posterior head ([Figure 7L](#)).

Activation of subsets of head BMNs elicits aimed grooming of specific locations

We next tested the extent to which the parallel-projecting BMNs elicited aimed grooming of specific head locations. The driver lines described above ([Figure 3D–I](#)) were used to express the light-gated neural activator CsChrimson (Klapoetke et al., 2014) in different subsets of BMNs ([Figure 8A](#)). Flies were placed in chambers where they could move freely and then exposed to red light to activate the CsChrimson-expressing BMNs. We manually annotated the movements elicited by optogenetic activation of BMNs from recorded video ([Figure 8 – figure supplement 1](#), Videos 1–4).




Figure 8.

Optogenetic activation of BMNs at specific head locations elicits aimed grooming.

(A) Bristles shaded black on the anterior (left) and posterior (right) head are innervated by BMNs that express CsChrimson under control of the indicated driver lines. Control-spGAL4 shows no expression. (B) Histograms of manually annotated video for each line show movements elicited with red-light induced optogenetic activation. The fraction of flies performing each movement are plotted in one second bins (N = 10 flies per line). Grooming movements are indicated by different colors, including eye (magenta), dorsal head (blue), and ventral head (orange) grooming. Other elicited movements include

backward motion (black) and head nodding (gray). Gray bars indicate a 5 second red-light stimulus. Most driver lines were tested using 30 second interstimulus intervals, while pBMN-spGAL4 elicited more reliable behavior using 10 second intervals. Movements are mutually exclusive except head nodding. Representative experimental trials shown in **Video 1**, **Video 2**, **Video 3**, **Video 4**, and **Video 5**. **Figure 8 – figure supplement 1** shows additional controls and ethograms for individual flies tested. (C) Box plots show the percent time that flies spent performing each movement during the experiment shown in B. Bottom and top of the boxes indicate the first and third quartiles, respectively; median is shown in each box; whiskers show the minimum and maximum values. Asterisks indicate *p,0.05, **p,0.001, ***p,0.0001 from Mann-Whitney U pairwise tests between each experimental line and its corresponding control after application of Bonferroni correction. **Figure 8 – source data 1** contains numerical data used for producing each box plot.

Optogenetic activation of BMN types labeled by each driver line elicited grooming by the front legs that was aimed at specific head locations (**Figure 8B,C**). For example, a line that expressed in different BMN types on the dorsal head elicited aimed dorsal head grooming (dBMN-spGAL4; BM-InOc, -Vt, and -dPoOr neurons, blue trace, **Video 1**). Two lines expressed exclusively in specific BMN types, which enabled us to test the extent to which grooming was aimed specifically at those BMNs (i.e. BM-Taste and -InOm neurons). Indeed, BM-Taste neurons on the labellum elicited labellar grooming, but also grooming of neighboring locations on the proboscis and ventral head (TasteBMN-spGAL4, yellow trace includes proboscis and ventral head grooming, **Video 2**). Activation of BM-InOm neurons (InOmBMN-LexA, **Video 3**) elicited eye grooming (red trace), but also grooming of the neighboring dorsal head (blue trace). This suggested that head BMNs elicit aimed grooming of their corresponding bristle locations, but also neighboring locations.

Activation of BMNs on the posterior head elicited some dorsal head grooming (blue trace), but mostly a forward head nodding movement (**Figure 8A-C**, pBMN-spGAL4; BM-Vt, -dOcci, -dPoOr, and -vOcci neurons, **Video 4**). Nodding was an apparent avoidance response to

posterior touches of the head, and occurred mostly while the flies either stood in place or walked around. However, nodding was also observed during dorsal head grooming. Such nodding movements during head grooming were previously shown to help the legs reach particular locations (Honegger et al., 1979). Nodding also occurred with the dorsal head grooming elicited using the dBMN-spGAL4 and InOmBMN-LexA driver lines, but these lines did not elicit nodding in the absence of grooming as we observed with pBMN-spGAL4. This suggested that BMN-activated nodding occurs in two different behavioral contexts: during dorsal head grooming and as an avoidance response. Different evidence led us to hypothesize that nodding in these contexts was elicited by distinct BMN types. First, pBMN- and dBMN-spGAL4 driver lines show overlapping expression in BM-Vt and -dPoOr neurons, and both elicit dorsal head grooming accompanied by nodding. Second, pBMN-spGAL4 is the only tested line that expressed in BM-dOcci and -vOcci neurons and also the only line that elicited nodding in the absence of grooming. When taken together, our experiments suggest that nodding-only movements are elicited by BM-dOcci and -vOcci neurons and dorsal head grooming is elicited by BM-InOc, -Vt, -dPoOr, and -InOm neurons.

In addition to grooming, BMNs on the dorsal head and eyes elicited backwards motions that appeared as if flies were avoiding something that touched the head (dBMN-spGAL4 and InOmBMN-LexA). The backward motion and grooming were mutually exclusive and sequential, as the backward motion occurred transiently at the stimulus onset and was followed by grooming. As we reported previously (Hampel et al., 2020a), the red-light stimulus also elicited backward motions with control flies (Figure 6B,C, control, black trace, Video 5). However, control flies only responded in 33% of trials, whereas BMN activation flies responded with backward motions in most trials (73% for dorsal head BMNs, 100% BM-InOm neurons). Taken together, this study reveals that the somatotopically organized head BMNs elicit both aimed grooming and avoidance responses.

Discussion

Comprehensive definition of head BMNs

A major outcome of this work was the definition of nearly all BMNs on the *Drosophila* head. Although there were previous descriptions of the BMNs from different body parts, there were no comprehensive descriptions of all BMNs for any part. Furthermore, the head BMNs were among the least well described. Here, we modified a previously reported BMN dye fill method and produced new transgenic driver lines to define the projection morphologies of the different head BMN types that innervate specific bristle populations on the head. We then identified and reconstructed these types in the FAFB EM dataset. This provides the most comprehensive definition of the BMNs for any body part of *Drosophila* (or any other insect), and an essential resource for future studies. The annotated neurons can be linked to the ongoing neural circuit reconstructions in FAFB (Dorkenwald et al., 2022), or identified in anticipated new EM reconstructions of the brains of other individuals using available and emerging tools (Galili et al., 2022).

While nearly all head BMNs were reconstructed in this work, different knowledge gaps remain. It is unclear if the PoOc and Su bristles are innervated by BMNs because they could not be observed using transgenic driver lines or dye filling methods. We proposed that one of the BM-Vt/PoOc neurons innervates the PoOc bristle, based on proximity and presumed similarity to the neighboring BM-Vt, -InOc, and -Oc neurons. For the Su bristles, one possibility is that they are innervated by some of the 25 unknown sensory neurons reconstructed in this work (Figure 5 – figure supplement 2A-Y). It also remains unclear what neurotransmitter(s) are used by the BMNs. A machine learning approach was recently

developed that can predict whether a neuron in FAFB uses any of 6 major neurotransmitters with high accuracy (Eckstein et al., 2020). Given that the neurotransmitter predictions for the BMNs were overwhelmingly cholinergic (not shown), and a previous study indicated that leg BMNs are sensitive to a nicotinic acetylcholine receptor antagonist (Tuthill and Wilson, 2016a), the parsimonious explanation is that the BMNs are cholinergic. However, other studies suggest that BMNs could use histamine as a neurotransmitter (Melzig et al., 1996; Salvaterra and Kitamoto, 2001; Yasuyama and Salvaterra, 1999). Thus, the extent to which head BMNs use acetylcholine, histamine, or other neurotransmitters remains unresolved.

Resource: nearly all head mechanosensory neurons reconstructed and annotated in FAFB

In conjunction with two previous studies, work presented here contributes to the FAFB reconstruction and annotation of neurons associated with the major head mechanosensory structures, including the bristles (BMNs), JO (JONs), and taste pegs (TPMNs). The BMNs and TPMNs were reconstructed using the FlyWire.ai platform in the present work, and the JONs were previously reconstructed using the CATMAID platform (Hampel et al., 2020a; Kim et al., 2020). The TPMNs (38 reconstructed) respond to tactile displacements of the taste pegs and are implicated in feeding behavior (Jeong et al., 2016; Sánchez-Alcañiz et al., 2017; Zhou et al., 2019). Subpopulations of ~480 JONs have been previously defined that respond to diverse mechanical forces that move the antennae (JO-A, -B, -C, -D, -E, -F, and -mz neurons), including sound, gravity, wind, and tactile displacements (Hampel et al., 2015; 2020a; Ishikawa et al., 2017; Kamikouchi et al., 2009; Mamiya and Dickinson, 2015; Matsuo et al., 2014; Patella and Wilson, 2018). The JONs are implicated in different behaviors including courtship, flight, locomotion, gravitaxis, wind-guided orientation, escape, and head grooming (Hampel et al., 2015; 2020a; Kamikouchi et al., 2009; Lehnert et al., 2013; Mamiya et al., 2011; Mamiya and Dickinson, 2015; Suver et al., 2019; Tootoonian et al., 2012; Vaughan et al., 2014; Yorozu et al., 2009). The reconstruction and annotation of head mechanosensory neurons in FAFB provides an important resource for connectomics-based studies of mechanosensory processing (**Supplementary file 2**). While the majority of mechanosensory neurons on the head are now identified in FAFB, some remain to be identified, such as multidendritic and pharyngeal mechanosensory neurons on the proboscis (Yang et al., 2021; Zhang et al., 2016).

The reconstructed JONs, BMNs, and TPMNs project into distinct regions in the SEZ (**Figure 5 – figure supplement 1A,E**), and therefore show modality-specific projections. For example, the JONs (chordotonal neurons) define a region of the SEZ called the antennal mechanosensory and motor center (AMMC) while the BMNs project more ventrally. While the BMN projections are based on head location (somatotopic), the JON projections are based on mechanical stimulus modality, such as their responses to vibrational or tonic antennal movements (tonotopic) (Hampel et al., 2020a; Kamikouchi et al., 2006; Kim et al., 2020; Patella and Wilson, 2018). However, there are potential overlapping projections between the most ventral projecting JONs (JO-F neurons) and some BMNs projecting through the AntNv (**Figure 5 – figure supplement 1**). Among the BMNs that appear to overlap with the JO-F neurons are the BM-Ant neurons that are located on the same antennal segment as the JONs (pedicel). This suggests that the JO-F neuron projections are somatotopic like the BMNs. Modality-specific mechanosensory projections are also reported in the VNC of *Drosophila* and other insects (e.g. BMNs, hair plates, campaniform sensilla, and chordotonal neurons), revealing this organization to be fundamental in insects (Merritt and Murphey, 1992; Murphey et al., 1989a; Phelps et al., 2021; Smith and Shepherd, 1998; Tsubouchi et al., 2017; Tuthill and Wilson, 2016a).

A synaptic resolution somatotopic map of the head

This work defines the somatotopic organization of the head BMNs. Somatotopy was previously reported for BMNs innervating bristles on the bodies of *Drosophila* and other insects (Johnson and Murphey, 1985; Murphey et al., 1989b; Newland, 1991; Newland et al., 2000; Tsubouchi et al., 2017). However, these studies only produced partial somatotopic maps using dye fills or transgenic driver lines. Furthermore, there were no previous descriptions of somatotopy among the head BMNs. Here, we use EM reconstructions to produce a comprehensive synaptic resolution somatotopic map of head BMNs in the same brain.

All reconstructed *Drosophila* head BMN types terminate their projections in the SEZ. This indicates that the first layers of BMN processing for the head occur in the SEZ. In contrast, head BMNs reported in other insects project into both the SEZ and thoracic ganglia, including BMNs innervating the InOm bristles of the praying mantis and cricket (Honegger, 1977; Zack and Bacon, 1981) and wind sensitive head bristles of the locust (Tyrer et al., 1979).

The projections of head BMNs innervating the same bristle populations (same types) show the highest morphological similarity, and their morphology is stereotyped across individual flies. These characteristics likely apply to most BMNs, as numerous studies have identified the stereotyped projection morphologies of BMNs innervating specific bristles on the bodies of *Drosophila* and other insects (Burg and Wu, 1986; 1989; Burg et al., 1993; Chen et al., 2006; Ghysen, 1980; Honegger, 1977; Kays et al., 2014; Murphey et al., 1989b; Zack and Bacon, 1981). Head BMNs of the same type also show the highest postsynaptic connectivity similarity. However, some BMN types fall into multiple different NBLAST and cosine similarity clusters, revealing that there are BMN subtypes with differing morphology and postsynaptic connectivity. One notable example of such intratype diversity are the BM-InOm neurons that show differential clustering (Figure 7 – figure supplement 3). This could reflect the large surface area of the eyes that spans from the dorsal to ventral head, and the differentially clustered BM-InOm neurons may innervate bristles at different locations on the eyes. Future studies will address the organizational and functional logic of such intratype diversity.

We also find that BMN types innervating neighboring bristle populations have overlapping projections into zones that correspond roughly to the dorsal, ventral, and posterior head. The overlap is likely functionally significant, as cosine similarity analysis revealed that neighboring head BMN types have common postsynaptic partners. However, overlap between neighboring BMN types is only partial, as they show differing projections and postsynaptic connectivity. The extent of overlap likely reflects the proximity between bristles and enables postsynaptic partners to respond to mechanosensory stimulations of neighboring bristles whose corresponding BMNs are likely to show correlated activity (Tuthill and Wilson, 2016b). BMN projection overlap has been observed with other parts of the body in *Drosophila*. For example, BMNs innervating bristles on the anterior and posterior leg compartments show overlapping projections in the VNC leg neuromere anterior and posterior zones, respectively (Murphey et al., 1989b). Similarly, BMNs innervating neighboring bristles on the thorax show overlap in their projections into the accessory mesothoracic neuropil (Ghysen, 1980; Kays et al., 2014).

The somatotopic map reveals that some head BMNs have projections that remain in the ipsilateral brain hemisphere, while others have midline-crossing projections to the contralateral hemisphere (Figure 7A-E). Interestingly, BMNs innervating bristles located medially on the anterior head show midline-crossing projections, whereas those innervating more lateral populations have ipsilateral-only projections. Previous studies found that BMNs innervating medial bristles on the thorax have midline-crossing projections, while those innervating more lateral bristles have ipsilateral-only projections (Ghysen, 1980; Kays et al., 2014). Similarly, BMNs that innervate bristles located on the leg segment most medial to the

body (coxa) have midline-crossing projections (Murphey et al., 1989b; Phelps et al., 2021). This is also the case for BMNs on the legs of other insects, such as the cricket (Johnson and Murphey, 1985) and hawkmoth (Kent and Levine, 1988). Why do some BMNs have ipsilateral and midline-crossing projections? One possibility is that these BMNs can excite postsynaptic circuitry in both brain hemispheres to elicit bilateral leg grooming responses, which could be appropriate for medial stimuli. In contrast, BMNs on the proboscis have mixtures of ipsilateral-only and midline-crossing projections, while those on the posterior head show ipsilateral-only projections. Thus, the organizational logic of midline-crossing BMNs described above may not be universal.

This work provides the first synaptic resolution somatotopic map of a head (or body) for any species. Previous studies identified somatotopic maps across species, such as the vertebrate somatotopic maps of head and body (Abraira and Ginty, 2013; Adibi, 2019; Brown et al., 1977). Somatotopic organization has been found to be preserved at different layers of the nervous system and is thought to be of fundamental importance, although the full functional significance of this organization is unclear (Kaas, 1997; Thivierge and Marcus, 2007). Therefore, it remains important to produce anatomical and functional somatotopic maps and define how this somatotopy interfaces with postsynaptic circuits. It has previously not been possible to obtain a comprehensive description of a somatotopic map, as most studies were limited to sparse labeling experiments and extrapolation across the different animals. Thus, the spatial relationships among mechanosensory neurons that make up particular maps could not be definitively determined. Furthermore, none of these maps enabled the definition of the postsynaptic partners. We overcame this through the first complete EM reconstruction of a somatotopic map of a head in the same brain. This enables future work that will define the postsynaptic connectome of this complete somatotopic map. Thus, the synaptic resolution map provided here has important implications for expanding our understanding of somatotopic neural circuit organization and function.

Circuits that elicit aimed grooming of specific head locations

We report here that activation of the head BMNs elicits aimed grooming. Flies groom specific head locations, including the eyes, antennae, dorsal head, ventral head, and proboscis (Dawkins and Dawkins, 1976; Hampel et al., 2015; 2017; 2020a; Seeds et al., 2014; Szebenyi, 1969; Zhang et al., 2020). With the exception of the BM-InOm neurons, little was known about the roles of the other BMNs in eliciting head grooming. The BM-InOm neurons were originally identified as necessary for grooming in response to mechanical stimulation of the eyes in the praying mantis (Zack and Bacon, 1981). Mechanical stimulation of the *Drosophila* InOm bristles (Melzig et al., 1996) and optogenetic activation of the BM-InOm neurons (Hampel et al., 2017; Zhang et al., 2020) were later reported to elicit eye grooming. Here, we used optogenetic activation to further define the movements elicited by BM-InOm neurons, and show that other BMN types elicit grooming of the dorsal and ventral head. Previous studies in *Drosophila* and other insects showed that stimulations of bristles on the legs, wings, and thorax also elicit aimed grooming (Corfas and Dudai, 1989; Li et al., 2016; Matheson, 1997; Page and Matheson, 2004; Usui-Ishihara et al., 1995; Vandervorst and Ghysen, 1980). Thus, the BMNs are important for eliciting aimed grooming of specific locations on the head and body.

While we show that the parallel-projecting head BMNs elicit grooming of specific locations (i.e. eyes, dorsal and ventral head), the full range of aimed grooming movements that can be elicited was not explored. For example, antennal grooming was previously shown to be elicited by JON activation (Hampel et al., 2015; 2020a; Zhang et al., 2020), and we hypothesize here the BM-Ant neuron activation also elicits antennal grooming. However, we did not

identify a transgenic driver line that labels BM-Ant neurons that would enable us to test this hypothesis. Previous studies of the legs, wings, and thorax used mechanical stimulation of specific bristles, rather than BMN optogenetic activation to test the ranges of grooming movements that could be elicited. This was done by delivering mechanical stimuli directly to the bristles of decapitated flies that do not move unless stimulated. In contrast, stimulating the head bristles is relatively challenging, as it requires delivering precise mechanical stimulations to specific bristles in intact and tethered flies. We have used optogenetic analysis in this study, as it was previously demonstrated that BMN optogenetic activation elicits grooming that is comparable to mechanically stimulating their corresponding bristles (Hampel et al., 2017; Zhang et al., 2020). However, our ability to test the full range of grooming movements elicited with BMN activation was limited by the driver lines produced in this study (Figure 3D-G).

How do the parallel-projecting head BMNs interface with postsynaptic neural circuits to elicit aimed grooming of specific head locations? Different evidence supports the hypothesis that the BMNs connect with parallel circuits that each elicit a different aimed grooming movement (Seeds et al., 2014). First, cosine similarity analysis revealed parallel connectivity at the first postsynaptic layer. Second, previous studies showed that optogenetic activation of different sensory and interneuron types elicits grooming of specific head locations, suggesting that they are components of putative parallel circuits (Cande et al., 2018; Guo et al., 2022; Hampel et al., 2015; Seeds et al., 2014; Zhang et al., 2020). Third, we identified different neuron types whose activation elicit grooming of the antennae and showed that they are connected to form a neural circuit (Hampel et al., 2020b; 2015). The inputs to this circuit are JONs that detect tactile stimulations of the antennae and project to the SEZ where they excite two interneuron types (aBN1 and aBN2) and a descending neuron type (aDN) to elicit grooming. The aDNs project to a zone in the VNC where circuitry for generating antennal grooming leg movement patterns is thought to reside (Berkowitz and Laurent, 1996). While this circuit is postsynaptic to the JONs (Hampel et al., 2020b; 2015), preliminary connectomic analysis reveals that it is also postsynaptic to BMNs (not shown). Future studies will define the BMN connectivity with the antennal grooming circuit, and other neurons whose activation elicit aimed head grooming.

We find that activation of specific BMN types elicits both aimed grooming of their corresponding bristle locations and neighboring locations. This suggests overlap in the locations that are groomed with the activation of different BMN types. Such overlap provides a means of cleaning the area surrounding the stimulus location. Interestingly, our NBLAST and cosine similarity analysis indicates that neighboring BMNs project into overlapping zones in the SEZ and show common postsynaptic connectivity. Thus, we hypothesize that neighboring BMNs connect with common neural circuits (e.g. antennal grooming circuit) to elicit overlapping aimed grooming of common head locations.

BMN involvement in multiple distinct behaviors

In addition to grooming, this work identifies avoidance-like behaviors that are elicited by the head BMNs and their corresponding bristles. Previous studies implicated the InOm bristles in an avoidance response (Melzig et al., 1996), although this response was not described in detail. Here, we demonstrate that activation of the BM-InOm neurons elicits an avoidance-like response in the form of backward motions. This response was also elicited by activating BMN types on the dorsal head. Another putative avoidance-like behavior, head nodding, was found to be elicited by posterior head BMNs. Avoidance responses to bristle stimulation have been previously reported in *Drosophila* and other insects, such as limb withdrawal and postural changes (Melzig et al., 1996; Newland and Burrows, 1997; Pflüger, 1980; Vandervorst and Ghysen, 1980). Thus, BMNs across the head and body elicit grooming and avoidance responses.

Parallel circuit architecture underlying the grooming sequence

This study examines the mechanosensory layer of the parallel model of hierarchical suppression that produces the head to body grooming sequence (Hampel et al., 2017; Mueller et al., 2019; Seeds et al., 2014). This layer consists of mechanosensory neurons at specific locations on the head and body that elicit aimed grooming of those locations (Hampel et al., 2020a; 2017; 2015; Zhang et al., 2020). The aimed movements are performed in a prioritized sequence when mechanosensory neurons detect dust at different locations and become simultaneously activated (i.e. head and body completely dirty). In support of this, simultaneous optogenetic activation of mechanosensory neurons across the head and body elicits a grooming sequence that resembles the dust-induced sequence (Hampel et al., 2017; Zhang et al., 2020). Among the different mechanosensory neurons, the BMNs are particularly important, as their activation alone is sufficient to elicit a grooming sequence (Zhang et al., 2020). Thus, activation of individual BMN types elicits aimed grooming, while their simultaneous activation elicits a sequence. Here we define the parallel architecture of BMN types that elicit the head grooming sequence that starts with the eyes and proceeds to the antennae and ventral head.

The parallel-projecting head BMNs are hypothesized to connect with postsynaptic circuits that perform different functions to produce the sequence (Seeds et al., 2014). We described above how parallel circuits elicit grooming of specific locations. Simultaneous activation of this parallel architecture by dust causes competition among all movements to be performed in the sequence. This competition is resolved through a hierarchical suppression mechanism whereby earlier movements suppress later ones. Performance order is established by an activity gradient among the parallel circuits where earlier movements have the highest activity and later ones have the lowest. This gradient was proposed to be produced by controlling sensory gain among the BMNs, or through putative lateral inhibitory connections between the parallel circuits. A winner-take-all network selects the movement with the highest activity and suppresses the others. Our work here provides the foundation for studies that will examine how the BMN postsynaptic circuitry is organized to drive these different functions and produce the grooming sequence.

The BMNs are hypothesized to have roles in both eliciting and terminating the different movements in the grooming sequence through dust detection (Hampel et al., 2017; Seeds et al., 2014; Zhang et al., 2020). That is, dust on particular body parts would be detected by BMNs that are activated with displacement of their corresponding bristles and elicit aimed grooming. While a completely dirty body part would cause strong BMN activation, the level of activation would decrease as a consequence of the decreased dust levels that occur with grooming. This reduced activity would terminate the selected movement in the sequence, allowing a new round of competition among the remaining movements and selection of the next movement through hierarchical suppression. However, it has not been directly demonstrated that the BMNs elicit or terminate the sequence through dust sensing. For example, blocking BM-InOm neurons does not reduce dust-induced grooming of the head (Zhang et al., 2020). However, this may be due to compensation from other mechanosensory neurons, such as the JONs. Thus, the presumed role of the BMNs in detecting dust remains to be directly demonstrated.

Materials and methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (<i>D. melanogaster</i>)	R52A06-GAL4	Jenett <i>et al.</i> , 2012	RRID:BDSC_38810	
Genetic reagent (<i>D. melanogaster</i>)	VT017251-LexA	Hampel <i>et al.</i> , 2017		aka InOmBMN-LexA
Genetic reagent (<i>D. melanogaster</i>)	VT019023-AD	Tirian <i>et al.</i> , 2017	RRID:BDSC_71430	
Genetic reagent (<i>D. melanogaster</i>)	VT050279-DBD	Tirian <i>et al.</i> , 2017	RRID:BDSC_72433	
Genetic reagent (<i>D. melanogaster</i>)	R28D07-AD	Dionne <i>et al.</i> , 2017	RRID:BDSC_70168	
Genetic reagent (<i>D. melanogaster</i>)	VT023783-AD	Tirian <i>et al.</i> , 2017	RRID:BDSC_73261	
Genetic reagent (<i>D. melanogaster</i>)	R11D02-DBD	Dionne <i>et al.</i> , 2017	RRID:BDSC_68554	
Genetic reagent (<i>D. melanogaster</i>)	dBMN-spGAL4	This paper		Stock contains VT019023-AD and VT050279-DBD
Genetic reagent (<i>D. melanogaster</i>)	pBMN-spGAL4	This paper		Stock contains R28D07-AD and VT050279-DBD
Genetic reagent (<i>D. melanogaster</i>)	TasteBMN-spGAL4	This paper		Stock contains VT023783-AD and R11D02-DBD
Genetic reagent (<i>D. melanogaster</i>)	20XUAS-IVS-mCD8::GFP	Pfeiffer <i>et al.</i> , 2010	RRID:BDSC_32194	
Genetic reagent (<i>D. melanogaster</i>)	13XLexAop2-IVS-myr::GFP		RRID:BDSC_32209	
Genetic reagent (<i>D. melanogaster</i>)	C155-GAL4, UAS-nSyb.eGFP	Kendal Broadie	RRID:BDSC_6920	
Genetic reagent (<i>D. melanogaster</i>)	BPADZp; BPZpGDBD	Hampel <i>et al.</i> , 2015	RRID:BDSC_79603	spGAL4 control
Genetic reagent (<i>D. melanogaster</i>)	BDPLexA	Pfeiffer <i>et al.</i> , 2010	RRID:BDSC_77691	
Genetic reagent (<i>D. melanogaster</i>)	20XUAS-IVS-CsChrimson-mVenus	Klapoetke <i>et al.</i> , 2014	RRID:BDSC_55134	
Genetic reagent (<i>D. melanogaster</i>)	MCFO-5	Nern <i>et al.</i> , 2015	RRID:BDSC_64089	
Genetic reagent (<i>D. melanogaster</i>)	MCFO-3	Nern <i>et al.</i> , 2015	RRID:BDSC_64087	

Genetic reagent (<i>D. melanogaster</i>)	13XLexAop2- IVS-CsChrim son-mVenus		RRID:BDSC_55137	
Antibody	anti-GFP (Rabbit polyclonal)	Thermo Fisher Scientific	Cat# A-11122, RRID:AB_221569	IF(1:500)
Antibody	anti-Brp (Mouse monoclonal)	DSHB	Cat# nc82, RRID:AB_2314866	IF(1:50)
Antibody	anti-FLAG (Rat monoclonal)	Novus Biologicals	Cat# NBP1-06712, RRID:AB_1625981	IF(1:300)
Antibody	anti-HA (Rabbit monoclonal)	Cell Signaling Technology	Cat# 3724, RRID:AB_1549585	IF(1:500)
Antibody	anti-V5 (Mouse monoclonal)	BIO-RAD	Cat# MCA1360, RRID:AB_322378	IF(1:300)
Antibody	anti-Rabbit AF488 (Goat polyclonal)	Thermo Fisher Scientific	Cat# A-11034, RRID:AB_2576217	IF(1:500)
Antibody	anti-Mouse AF568 (Goat polyclonal)	Thermo Fisher Scientific	Cat# A-11031, RRID:AB_144696	IF(1:500)
Antibody	anti-Rat AF633 (Goat polyclonal)	Thermo Fisher Scientific	Cat# A-21094, RRID:AB_2535749	IF(1:500)
Chemical compound, drug	Paraformalde hyde 20%	Electron Microscopy Sciences	Cat# 15713	
Chemical compound, drug	DiD solid	Thermo Fisher Scientific	Cat# 07757	
Chemical compound, drug	all- <i>trans</i> -Reti nal	Toronto Research Chemicals	Cat# R240000	
Software, algorithm	neuTube	Feng <i>et al.</i> , 2015		https://www.neutracng.com/
Software, algorithm	Vcode	Hagedorn <i>et al.</i> , 2008		http://social.cs.uiuc.edu/p rojects/vcode.html
Software, algorithm	Fiji	Schindelin <i>et al.</i> , 2012		http://fiji.sc/
Software, algorithm	R			https://www.r-project.org/
Software, algorithm	CMTK	Jefferis <i>et al.</i> 2007		https://www.nitrc.org/proje cts/cmtk/
Software, algorithm	FluoRender	Wan <i>et al.</i> , 2012		http://www.sci.utah.edu/s oftware/fluorender.html
Software, algorithm	Blender version 2.79			https://www.blender.org/d ownload/releases/2-79/
Software, algorithm	MATLAB	MathWorks Inc., Natick, MA		
Software, algorithm	natverse	Bates <i>et al.</i> 2020		http://natverse.org/
Software, algorithm	Cytoscape	Shannon <i>et al.</i> 2003		https://cytoscape.org/

Rearing conditions and fly stocks

GAL4, LexA, and *Split GAL4* (spGAL4) lines were generated by the labs of Gerald Rubin and Barry Dickson, and most lines are available from the Bloomington *Drosophila* Stock Center (Dionne et al., 2017; Jenett et al., 2012; Pfeiffer et al., 2008; Tirián and Dickson, 2017). Canton S flies were obtained from Martin Heisenberg's lab in Würzburg, Germany. Other stocks used in this study are listed in the key resources table.

GAL4, spGAL4, and LexA lines were crossed to either UAS or LexAop driver lines as described below. Flies were reared on Fisherbrand Jazz-Mix food *Drosophila* food (Fisher Scientific, Fair Lawn, NJ) containing corn meal, brown sugar, yeast, agar, benzoic acid, methyl paraben, and propionic acid. The flies were kept in an incubator at 21°C and 55-65% relative humidity. Flies that were not used for optogenetic experiments were kept on a 16/8 hour light/dark cycle. Flies used for optogenetic experiments were reared on food containing 0.4 mM all-*trans*-retinal (Toronto Research Chemicals, Toronto, Canada) in vials that were wrapped in aluminum foil and kept in a box to keep them in the dark. Unless stated otherwise, the flies used for experiments were 5 to 8 day-old males.

Imaging the head bristles

1-2 mm was cut off the tip of an Eppendorf 200 µL pipette tip, then an approximately 6 mm length was cut off and the remainder discarded. A freeze-killed (> 1 hour) male or female Canton S fly was then gently pushed in with a piece of wire, until the head protruded from the tip. The tip was then mounted in a small piece of soft wax. An observation chamber was constructed on a microscope slide, by cutting a 5 mm square hole in 3 layers of Highland electrical insulation tape (3M, Saint Paul, Minnesota) and covering the bottom with a translucent white plastic square (cut from a Farmland Traditions dog treat bag). The fly, held in the tube, was mounted over the chamber using the wax, first dorsal side up (imaged), then ventral side up (imaged). The head was carefully cut off using sharpened iridectomy scissors, falling into the chamber where it was arranged anterior side up, held in place with a piece of coverslip, imaged, flipped posterior side up, and imaged again. Imaging was done with a Zeiss Axio Examiner D1 microscope equipped with a 10x Achroplan objective (0.25 NA) (Karl Zeiss, Oberkochen, Germany). The objective was surrounded with a cylinder of the same translucent white plastic in order to diffuse the light source and to avoid air movements that could move the antennae. The cylinder was illuminated from both sides at 2-3 cm distance by a Dolan-Jenner Fiber-lite (Dolan-Jenner Industries, Boxborough, Massachusetts). A small amount of additional back-lighting was provided by the microscope light source (20% power) with a blue filter. Images were captured with a Zeiss AxioCam 512 at 60 ms exposure. The focal plane was advanced in small increments manually, resulting in approximately 50 images per head.

The image Sequence function of Fiji software (<http://fiji.sc/>) was used to combine all the images into a stack. The pixel size was adjusted to 0.3125 micron and calibrated with an image of a slide micrometer. The stack was down-sized to 2048 pixels minimum dimension (usually the height), and an Unsharp Mask filter applied (3 pixel radius, 0.6 mask weight), then the sides of the stack were cropped to remove unnecessary space.

The Fiji *extended depth of field* (EDF) plugin (Alex Prudencio, EPFL, École polytechnique fédérale de Lausanne) was used to superimpose in-focus areas from the stacks. For acceptable processing times, the stack was down-sized to 1024 pixels minimum dimension. The EDF process requires square images, so to avoid excessive cropping the canvas size was increased to 2048 pixels square, resulting in a black surround. The best results were obtained by averaging (with Image Calculator) (1) the result of EDF Easy Mode Fast setting,

Gaussian-blurred by 1 pixel radius, with (2) the result of EDF Easy Mode High setting. The resulting image was cropped to 1024 minimum dimension, sharpened with Unsharp Mask, radius 1 pixel, and adjusted for optimal contrast.

With anterior views of the head, the EDF algorithm has difficulties separating the arista from the underlying eye facets. Thus, for presentation images of the front view of the head, the EDF process was carried out separately on an anterior stack with the arista present, and a more posterior stack with the antennae absent. These results were imported as layers into GIMP (GNU Image Manipulation Program) and manually combined by masking.

To facilitate bristle identification, a color-coded depth map was constructed from the downsized 1024-height stack, sharpened with Unsharp-mask (1 pixel). The color channels were split, and G and B channels discarded. The R channel was Inverted (Edit menu), the contrast was adjusted, and a Gamma correction of 1.34 was applied to the stack. A lookup table (LUT) had been previously created ranging from light blue, through white and yellow to dark red (“Stellar”). The Image: Hyperstacks: Temporal-Color Code function (LUT Stellar) was applied to the stack, giving a depth-coded image. Finally, an Enhanced Local Contrast (CLAHE, blocksize 63) was applied, followed by a gamma adjustment (1.5-1.8).

Bristle nomenclature

Published names for the different bristles were used when possible. However, bristle abbreviations are from the present work unless otherwise indicated. Some bristle names were from (Bodenstein et al., 1994), including the *frontal* (Fr), *frontoorbital* (FrOr), *orbital* (Or), *ocellar* (Oc), *interocellar* (InOc), *vertical* (Vt), *postorbital* (PoOr), and *vibrissae* (Vib). We deviated from this nomenclature in the following cases. First, although the PoOr bristles form a continuous row along the back margin of each eye (Figure 1B), we subdivided them into *dorsal* (dPoOr) and *ventral* (vPoOr) populations based on whether their associated BMNs project through the OcciNv or EyeNv, respectively (Figure 2F,J). Second, we did not use the bristle name *postvertical*, but instead used *postocellar* (PoOc) that was previously proposed to better describe the location of these bristles as posterior to the other ocellar bristles (Steyskal, 1976). Third, instead of occipital, we used the name *supracervical* (Su) for the bristles located immediately above the cervical connective on the back of the head (Steyskal, 1976). This is because we named two populations of small bristles on the back of the head the *dorsal* and *ventral occipital* (dOcci and vOcci) bristles. This name was previously proposed for these bristles in other species of flies (Steyskal, 1976). The abbreviation (Occi) was taken from the blowfly literature (Thei, 1979). Given that the Occi bristles are found as two distinct populations in *Drosophila melanogaster*, we refer to them in this work as the *dorsal* (dOcci) and *ventral* (vOcci) bristles (Figure 1B).

Our abbreviation for the *vertical* (Vt) bristles describes all vertical bristles. The largest two have been referred to as *Vt inner* (Vti) and *exterior* (Vte) bristles in different fly species (Steyskal, 1976), and were named in the present work Vt 1 and Vt 2, respectively (Figure 4 – figure supplement 5A). The other previously described Vt bristle (named Vt 3) is posterior to Vt 2. Medial to the Vt 2 bristle is a newly categorized fourth Vt bristle that we could not find a previous description of in *Drosophila* (labeled Vt 4 in Figure 4 – figure supplement 5A). This bristle could be the paraverticlar bristle that was previously described (Steyskal, 1976).

The bristles on the eyes were referred to as *interommatidial* (InOm) bristles (Honegger et al., 1979; Zack and Bacon, 1981). For the bristles on the outer labellum, we used the common name, *taste* (Taste) bristles (Stocker, 1994). We refer to the other bristles on the proboscis as *maxillary palp* (MaPa) and *haustellum* (Hau) bristles. Note: while most of the bristles on the head are termed trichoid sensilla that are innervated by a single BMN, the Taste bristles are

mostly basiconic sensilla that are each innervated by a BMN and multiple gustatory neurons. The bristles on the first and second segments of the antennae are referred to as *antennal* (Ant) bristles.

Bristle quantification

We counted the bristle populations that are defined above and shown in [Figure 1A-D](#). Most bristles were counted using color-coded depth maps (described above) that aided the identification of the bristles within each population ([Figure 1 – figure supplement 1A-H](#)). The Ant, Fr, FrOr, Or, Oc, InOc, Vt, dOcci, PoOr, Vib, MaPa, Hau, PoOc, and Su bristles were counted using this method on eight male and four female heads ([Figure 1E](#), [Figure 1 – figure supplement 2A-E](#), [Supplementary file 1](#) (Table 1)). Bristles on each half of the head were counted using the Fiji Cell Counter plugin (Kurt De Vos, University of Sheffield) and then averaged. These averages were used to calculate average and standard deviation for each bristle population from all counted male and female heads. Two tailed t-tests were performed to compare male and female bristle numbers for each population. Although in this manuscript we make a distinction between the dPoOr and vPoOr bristles ([Figure 1B](#)), we counted all PoOr bristles together when comparing their numbers between males and females ([Figure 1 – figure supplement 2E](#)). We used other approaches to obtain or estimate the numbers of dPoOr, vPoOr, InOm, vOcci, and Taste bristles (described below).

BMNs that innervate the PoOr bristles project through two different nerves, the Occi- and EyeNvs ([Figure 2F](#)). Specifically, BMNs innervating *dorsal PoOr* (dPoOr) bristles project through the OcciNv, while those innervating *ventral PoOr* (vPoOr) bristles project through the EyeNv. We determined the average number of dPoOr or vPoOr bristles based on whether they were innervated by an Occi- or EyeNv-projecting BMN. The BMNs were labeled using R52A06-GAL4 to express membrane-targeted *green fluorescent protein* (mCD8::GFP) and imaged using a confocal microscope (imaging method described below). We then counted the dPoOr and vPoOr bristles from confocal images (example shown in [Figure 2 – figure supplement 1D](#)). Multiple different heads were counted to determine the average number of dPoOr and vPoOr bristles (N = 10, [Supplementary file 1](#) (Table 2)).

The InOm bristles were too small and numerous to be counted, and it was only necessary to estimate their numbers in this work. The eyes contain the majority of bristles on the head and the numbers of these bristles can vary. Data from a previous study indicated that each eye contains between 745 and 828 regularly spaced ommatidia (776 average), and most ommatidia have an associated bristle ([Ready et al., 1976](#)). However, some ommatidia around the eye edges are not associated with bristles. Based on one example of an eye from a Canton S fly ([Ready et al., 1976](#)), we calculated that 78% of the ommatidia had an associated bristle for that eye. This percentage was used to estimate that there are between 607 and 645 bristles on each eye from the above counted bristle ranges. This estimate was sufficient for identifying the EM reconstructed BMNs that innervate the InOm bristles (described below) based on their overwhelming number relative to other head BMNs.

In this work we identified the vOcci bristles that were not previously described. These bristles were too small to be reliably observed at the level of resolution of the images shown in [Figure 1A-D](#). To help visualize and count these bristles, we used a transgenic driver line pBMN-spGAL4 (R28D07-AD $\cap$ VT050279-DBD) that labels BMNs innervating these bristles (shown in [Figure 3F,F'](#)). pBMN-spGAL4 was used to express mCD8::GFP, and the ventral posterior head was imaged with a confocal microscope (see below for imaging method). The bristles could be counted by using the labeled BMN dendrites to highlight their locations (shown in [Figure 3 – figure supplement 1C](#)). pBMN-spGAL4 labeled almost all of the visible vOcci bristles, but in some heads, we could see bristles that did not have an labeled BMN (not shown). Therefore, it is possible that the counts of these bristles using the GFP-labeled

BMNs are lower than the actual number. We determined the average number of Occi bristles by counting different heads (N = 13, **Supplementary file 1** (Table 3)).

Most of the heads that we imaged had their proboscises oriented such that we could not observe and count all of the Taste bristles. However, Taste bristles have been counted in previous studies (Falk et al., 1976; Jeong et al., 2016; Nayak and Singh, 1983; Shanbhag et al., 2001). See **Supplementary file 1** (Table 4) for Taste bristle counts from different publications (published counts for Ant and MaPa bristles are also shown). One of these studies also determined that there were no differences in the numbers of Taste bristles between males and females (Shanbhag et al., 2001). We took the highest and lowest numbers from these different references for the range that is shown in **Figure 1E**.

Head immunostaining and nerve reconstructions

R52A06-GAL4 (RRID:BDSC_38810), dBMN-spGAL4, pBMN-spGAL4, and TasteBMN-spGAL4 were crossed to *20XUAS-IVS-mCD8::GFP* (RRID:BDSC_32194) while VT017251-LexA (InOmBMN-LexA) was crossed to *13XLexAop2-IVS-myr::GFP* (RRID:BDSC_32209). Anesthetized male progeny were decapitated using a standard razor blade and heads were placed in phosphate-buffered saline (PBS). To facilitate antibody penetration for staining, we used #5 Dumoxel forceps (Fine Science Tools, Foster City, California) to tear small holes in the cuticle and pull off the antennae or proboscis. Heads were fixed in PBS with 2% paraformaldehyde for 1 hour at room temperature, and then washed with PAT (PBS, 1% Bovine Serum Albumin, 0.5 % TritonX) six times within 2 hours. Heads were blocked overnight at 4 °C in PAT with 3% normal goat serum (PAT-NGS), then incubated for 3 days (room temperature during the day and 4 °C at night) in PAT-NGS containing rabbit anti-GFP (Thermo Fisher Scientific, Waltham, MA, RRID:AB_221569). Heads were washed with PAT for 5 hours and then incubated for 3 days in PAT-NGS with goat anti-rabbit AlexaFluor-488 (Thermo Fisher Scientific, Waltham, MA, RRID:AB_2576217). Heads were washed for 2 days with several exchanges of PAT, and then in PBS for 2 hours at room temperature. A standard slide was used for mounting with a small “well” created by stacking five Avery reinforcement labels (Avery products corporation, Brea, California). A drop of Vectashield (Vector Laboratories, Inc. Burlingame, California) was added to the well and the heads were positioned either anteriorly or dorsally. The well was then covered with a circular cover slip (Electron Microscopy Sciences, Hatfield, Pennsylvania, 1.5 Micro Coverglass 12 mm diameter, Cat# 72230-01).

Heads were imaged using a Zeiss LSM800 confocal microscope (Carl Zeiss, Oberkochen, Germany) equipped with a 20x objective (Plan-Apochromat 20x/0.8). Fiji software (<http://fiji.sc/>) was used for examining the morphology of the imaged BMNs and for image processing steps, including adjustment of brightness and contrast, stitching, and image inversion. Reconstructions of the labeled head BMNs and their respective nerves from the confocal z-stacks (**Figure 2** and **Figure 3**) were performed using the software neuTube (Feng et al., 2015). Image stacks of the heads are displayed as maximum intensity projections (**Figure 2** and **Figure 3**).

CNS immunostaining and analysis

The different GAL4, spGAL4, and LexA driver lines described above were crossed to either *20XUAS-IVS-mCD8::GFP* or *13XLexAop2-IVS-myr::GFP*. Brains and ventral nerve cords were dissected and stained as previously described (Hampel et al., 2015; 2011). The following primary and secondary antibodies were used for staining GFP and the neuropil: rabbit anti-GFP, mouse anti-nc82 (Developmental Studies Hybridoma Bank, University of Iowa, RRID:AB_2314866) to stain Bruchpilot, goat anti-rabbit AlexaFluor-488, and goat anti-mouse

AlexaFluor-568 (Thermo Fisher Scientific, RRID:AB_144696). The stained CNSs were imaged, and confocal stacks processed as described above for heads.

To display the GFP expression patterns of different spGAL4 and LexA lines together as shown in **Figure 3J** we used the Computational Morphometry Toolkit (CMTK) (<https://www.nitrc.org/projects/cmtk/>) (Jefferis et al., 2007) to computationally register individual confocal stacks of each line to the JFRC-2010 standard brain (www.virtualflybrain.org (<http://www.virtualflybrain.org/>)). The PIC file of each registered stack was loaded into Fiji and merged to display each in a different color channel.

For multicolor flipout (MCFO) experiments, dBMN-spGAL4, pBMN-spGAL4, and TasteBMN-spGAL4 were crossed to the MCFO-5 stock (RRID:BDSC_64089) (Nern et al., 2015). 9-12 day-old fly brains were dissected for pBMN-spGAL4 and dBMN-spGAL4, while 4-6 day-old brains were dissected for Taste-spGAL4. Brains were stained using rat anti-FLAG (Novus Biologicals, LLC, Littleton, Colorado, RRID:AB_1625981), rabbit anti-HA (Cell Signaling Technology, Danvers, Massachusetts, RRID:AB_1549585), mouse anti-V5 (Bio-Rad, Hercules, California, RRID:AB_322378), goat anti-rabbit AlexaFluor-488, goat anti-mouse AlexaFluor-568, goat anti-rat AlexaFluor-633 (Thermo Fisher Scientific, RRID:AB_2535749). Stained brains were imaged, and confocal stacks processed as described above. Individually labeled neurons from each line are shown as maximum projections in **Figure 4** and **Figure 4 – figure supplements 6–8**. Given that very few flipout events occurred using the MCFO-5 stock in combination with the dBMN-spGAL4 and pBMN-spGAL4, we used a different MCFO stock (MCFO-3, RRID:BDSC_64087) to obtain a higher number of individually labeled neurons. Using dBMN-spGAL4 crossed with MCFO-3, we aged males for 9-12 days and obtained approximately 1-2 individually labeled neurons in 7 dissected brains. Crossing pBMN-spGAL4 with MCFO-3, we obtained about 1-4 labeled neurons in each dissected brain when we aged males 4-6 days. The flipout events in the latter cross occurred in much higher frequency given that most brains labeled more than a single neuron with the MCFO-3 stock.

Identification of driver lines that express in different subsets of head BMNs

We used the spGAL4 system to produce driver lines that expressed in different subsets of head BMNs (Luan et al., 2006; Pfeiffer et al., 2010). spGAL4 allows for independent expression of the GAL4 DNA binding domain (DBD) and activation domain (AD). When DBD and AD are expressed in the overlapping neurons, these domains reconstitute into a transcriptionally active protein. To label specific subpopulations of head BMNs, we visually screened through an image collection of the CNS expression patterns of enhancer-driven lines and identified candidate lines that were predicted to express in different subsets of head BMNs (Dionne et al., 2017; Jenett et al., 2012; Tirián and Dickson, 2017). We selected two candidate lines to express the DBD in subsets of head BMNs: VT050279 (VT050279-DBD, RRID:BDSC_72433) and R11D02 (R11D02-DBD, RRID:BDSC_68554). VT050279-DBD or R11D02-DBD flies carrying the *20XUAS-IVS-CsChrimson-mVenus* transgene (RRID:BDSC_55134) (Klapoetke et al., 2014) were crossed to 55 different candidate-ADs. The progeny were placed in behavioral chambers and exposed to red light for optogenetic activation (described below). We tested three flies for each DBD/AD combination to identify those that expressed in neurons whose activation could elicit grooming. Grooming “hits” were stained using a GFP antibody to detect CsChrimson-mVenus expression in the CNS and anti-NC82 to mark the neuropil as described above. We identified three different combinations that expressed in restricted subsets of BMNs, that included the ADs R28D07-AD (RRID:BDSC_70168), VT019023-AD (RRID:BDSC_71430), and VT023783-AD (RRID:BDSC_73261). We generated stable lines containing both the AD and DBD, including dBMN-spGAL4 (VT019023-AD $\cap$ VT050279-DBD),

pBMN-spGAL4 (R28D07-AD $\cap$ VT050279-DBD), and TasteBMN-spGAL4 (VT023783-AD $\cap$ R11D02-DBD).

Behavioral analysis procedures

For behavioral experiments, dBMN-spGAL4, pBMN-spGAL4, TasteBMN-spGAL4, and BPADZp; BPZpGDBD (spGAL4 control, RRID:BDSC_79603) were crossed to *20XUAS-CsChrimson-mVenus*. InOmBMN-LexA and BPADZp and BDPLexA (LexA control, RRID:BDSC_77691) were crossed to *13XLexAop2-IVS-CsChrimson-mVenus* (RRID:BDSC_55137). The controls used with the spGAL4 and LexA lines contain the vector backbone that was used to produce each line (including the coding regions for each spGAL4 half or LexA), but lack any enhancer to drive spGAL4 or LexA expression (Hampel et al., 2015; Pfeiffer et al., 2010; 2008).

We used a previously reported behavioral optogenetic rig, camera setup, and methods for the recording of freely moving flies (Hampel et al., 2017; 2015; Seeds et al., 2014). The stimulus parameters used were 656 nm red light at 27 mW/cm² intensity delivered at 5 Hz for 5 s (0.1 s on/off) with 10 or 30 second interstimulus intervals (total of 3 stimulations). While most of the driver lines were recorded using 30 second interstimulus intervals, pBMN-spGAL4 was recorded using 10 second intervals. This was because the elicited head nodding behavior occurred more robustly when stimulated every 10 seconds rather than 30 seconds.

Manual scoring of behavior from prerecorded video was performed using VCode software (Hagedorn et al., 2008) and analyzed in MATLAB (MathWorks Incorporated, Natick, MA). Some grooming and avoidance-like movements were annotated as previously described, including antennal, eye, ventral head, proboscis, and backward motion (Hampel et al., 2020a; 2017; 2015; Seeds et al., 2014). In this work, ventral head and proboscis grooming were combined and referred to as ventral head grooming. Head nodding was annotated when the fly tilted its head downward by any amount until it returned its head back in its original position. Dorsal head grooming was scored when the fly used one or both legs to touch its dorsal head, which included the dorsal part of the eye. During dorsal head grooming, flies sometimes nodded, rotated, or kept their heads in their original position to groom the outer dorsal areas of the head (dorsal eye and posterior dorsal eye).

Behavioral data was analyzed using nonparametric statistical tests as previously reported (Hampel et al., 2020a; 2017; 2015). The percent time flies spent performing each behavior was calculated. To compare the behavior performed by each experimental genotype with its corresponding genetic control, we performed pairwise comparisons for each behavior using a Mann-Whitney U test and applied Bonferroni correction. Note that we tested both male and female flies for optogenetic activation of behavior. Although only males are presented in this manuscript, optogenetic activation was found to elicit similar behaviors in both males and females (not shown).

Dye filling of BMNs that innervate large bristles

The dye filling protocol used in this study was adapted from one that was previously published (Kays et al., 2014). C155-GAL4, *UAS-nSyb.eGFP* flies (RRID:BDSC_6920) were used for the dye fill experiments to label the neuropil. Flies were decapitated with a standard razor blade and their heads glued to a microscope cover glass (Fisher Scientific, Pittsburgh, Pennsylvania) using TOA 400 UV cured glue (Kemxert, York, Pennsylvania). Heads were submerged in 3.7% w/v paraformaldehyde in 0.2 M carbonate-bicarbonate buffer at pH 9.5 overnight at 4°C. Heads were washed 24 hours later by dipping in 0.2 M carbonate-bicarbonate buffer at pH 9.5 for 30 seconds, and subsequently in ddH₂O for 30 seconds. Heads were gently blotted dry to prevent dye from spilling over the cuticle, and the selected bristles on the head were plucked with #5 Dumoxel forceps. Bristles were selected from

either the left or right side of the head, depending on which bristle was in the most optimal orientation for plucking and filling.

Micropipettes for dye filling were prepared from Borosilicate Thin Wall capillaries (Warner Instruments, Holliston, Massachusetts, G100T-4). Capillaries were filled with 5–10 μL of dye solution of 10 $\mu\text{g}/\mu\text{L}$ DiD (Thermo Fisher Scientific) in 100% ethanol and the tip was approached to the bristle socket with a micromanipulator. The tip of the capillary was made to contact the edge of the bristle socket such that the dye diffused into the socket until a stable bubble of solution formed. Heads were then dried for 5 minutes and then submerged in a 0.2 M carbonate-bicarbonate buffer at pH 9.5, in the dark, and at room temperature for 48 hours. The brains were then dissected and imaged immediately without fixation. Dissected brains were placed on a microscope slide with two Avery circular reinforcement labels and a circular cover slip. The brains were imaged immediately using a Zeiss LSM 800 confocal microscope. The native fluorescence of nSyb.eGFP was preserved enough at the conclusion of the experiment to image the brain neuropil ([Figure 4 – figure supplement 1B–E](#)).

We attempted to fill BMNs innervating the following head bristles: PoOc, Oc, Or, Ant, Vib, and Vt. Many attempts to fill particular bristles resulted in unfilled or partially filled BMNs. Anecdotally, there also seemed to be a difference in how well the filling method worked for the different bristle populations. For example, multiple attempts to fill the Vt bristles only resulted in one successful fill of a BMN from Vt 1 and Vt 3 bristles. Additionally, we were unable to fill a BMN with multiple attempts of the PoOc bristle. Successful fill trials and the locations of specific bristles that were filled are shown in [Figure 4C–Q](#) and [Figure 1 – figure supplements 2–5](#).

Reconstruction and analysis of bristle mechanosensory neurons from an EM volume

BMNs were reconstructed in a complete EM volume of the adult female brain (FAFB) dataset (Zheng et al., 2018) using the FlyWire.ai platform (Dorkenwald et al., 2022). We first identified the locations of the Ant-, Occi-, and merged Eye/LabNvs in the EM volume based on their identified locations from light microscopy data. We then chose a cross section of each nerve, close to where they enter brain neuropil, and where segmentation was available for all neurons in the nerve ([Figure 5 – figure supplement 1A–D](#)). We seeded every profile in the Occi- and merged Eye/LabNvs ([Figure 5 – figure supplement 1C,D](#)). The Eye/LabNvs had a bundle of soma tracts from an SEZ interneuron hemilineage crossing the seed plane that was excluded from the seeding process based on the morphology of their initial segmentation ([Figure 5 – figure supplement 1C](#)). Previous studies reconstructed major portions of JONs in FAFB AntNv using the CATMAID platform (Hampel et al., 2020a; Kim et al., 2020). Because the FlyWire FAFB brain was locally realigned, we transformed those JONs into the FlyWire space using natverse version 0.2.4 (Bates et al., 2020). After overlaying these JONs onto the seed plane, they were excluded during the seeding effort to identify BMNs in the AntNv. This left a small ventral-medial area of the nerve with previously undocumented neurons that were seeded and reconstructed ([Figure 5 – figure supplement 1B](#)). We focused our reconstructions in the right hemisphere nerves. However, we also examined the OcciNv in the left hemisphere, given that it only contained a small number of neurons ([Figure 6 – figure supplement 4A–D](#)).

The segmentations of all seeded neurons were then fully proofread by a human annotator. This process involves splitting falsely merged parts and merging falsely missing parts of a neuron using the tools available in the FlyWire neuroglancer instance (flywire.ai). FlyWire neuroglancer was also used to examine the morphologies of neurons for classification purposes.

The neurons classified as sensory origin (no soma in the brain, only axonal projections and entering through a nerve) were skeletonized using *natverse* (skeleton in *fabfseg* package version 0.10.0). Skeletons were pruned to synapse rich areas to exclude the smooth axon in the nerve bundle. We created a three dimensional mesh of the synapses (for synapses see below) and pruned the skeletons to arbors within the mesh volume. We then compared their morphology using the NBLAST algorithm (Costa et al., 2016) and clustered the similarity scores with Ward (Figure 6 – figure supplement 1A,B). The skeletons of CATMAID reconstructions of JONs from (Hampel et al., 2020a; Kim et al., 2020) and skeletonizations of sensory neurons from this publication were transformed into JRC2018F standard brain space for plotting using *natverse*. The transformed skeletons, meshes from the FlyWire proofread segmentation and brain neuropil meshes were plotted with *natverse* in RStudio 2022.02.3. We used the standard brain transformations to analyze if BMNs crossed the midline in the brain. Defining the midline accurately is possible due to the symmetric nature of this brain space. If any skeleton node coordinates of a given BMN were located in the contralateral hemisphere (x of node $>$ x of midline) we classed the BMN as midline-crossing.

FlyWire provides access to a synapse table imported from (Buhmann et al., 2021). We queried the table for pre- and postsynaptic sites belonging to the reconstructed sensory neurons for connectome analysis (Figure 7 – figure supplements 1–3). We used the cleft scores of the cleft prediction from (Heinrich et al., 2018) to filter synapses with scores below 50, which reliably excludes falsely predicted synapses.

We then analyzed the connectivity of BMNs with each other, and with other postsynaptic partners. First, we analyzed BMN type connectivity by adding up synaptic weights of BMN-to-BMN edges by type and normalizing by the number of possible edges between the groups (Figure 7 – figure supplement 1B). Graphs were plotted using Cytoscape version 3.9.1 and the RCy3 (version 2.17.1) and igraph (version 1.3.0) packages for Cytoscape control from R (Shannon et al., 2003). Further, we compared BMN types regarding their pre- and postsynaptic connection counts (Figure 7 – figure supplement 1A). We counted all entries in the FlyWire synapse table at which a given BMN was either the pre- and postsynaptic partner (not number of presynaptic sites, cleft scores $\geq$ 50). This revealed that BM-InOm neurons had fewer synaptic connections than other BMN types, which was not surprising given their small, non-complex axonic arbors in the brain. This posed a challenge when comparing BM-InOm neuron synaptic connectivity to other BMN types, as described below.

We calculated cosine similarity of BMNs to cluster them based on their connectivity similarity. Cosine similarity emphasizes the similarity of BMNs which have similar sets of downstream targets. We took advantage of the good proofreading state of the FlyWire dataset to perform postsynaptic connectivity analysis (Dorkenwald et al., 2022). Note that this strategy is agnostic about partner types. A comprehensive typing of the postsynaptic partners will be included in a follow up study as it exceeds the focus of this study. For this analysis, we excluded synapses between BMNs and only considered postsynaptic connectivity to neuron partners of other types (non-BMNs). As mentioned, BM-InOm neurons have few connections to postsynaptic partners (pre counts in Figure 7 – figure supplement 1A), so we choose a low threshold of 3 synapses and excluded edges below that. When clustering the cosine similarity (ComplexHeatmap version 2.11.2 package in R) for all BMNs, we found that subsets of the BM-InOm neurons clustered together with other BMN types (data not shown), but the clustering failed to capture meaningful groups. Choosing a higher threshold resulted in most of the BM-InOm neuron connectivity being excluded, giving low cosine similarity scores even when compared to each other. We thus choose a higher threshold of 7 synapses to analyze the cosine similarity of the BMN types excluding BM-InOm neurons (Figure 7 – figure supplement 2) and the low synapse threshold of 3 for analysis of the BM-InOm neurons separately (Figure 7 – figure supplement 3). This analysis better captured the somatotopic mapping of the BMN types in the brain. From our

preliminary analysis we expect BM-InOm neurons to share postsynaptic partners with other BMN types, but further analysis is required to investigate patterns of connectivity of all BMNs to postsynaptic targets.

Linking reconstructed BMNs to specific populations of head bristles

Manual categorization of the EM reconstructed sensory neurons revealed that they consist of BMNs, JONs, *taste peg mechanosensory neurons* (TPMNs), and *gustatory neurons* (GRNs). JONs and GRNs were previously identified and described in the EM volume using the CATMAID platform (Engert et al., 2022; Hampel et al., 2020a; Kim et al., 2020). The TPMNs were identified based on their morphological similarity to previous light microscopy descriptions (Jeong et al., 2016; Miyazaki and Ito, 2010; Zhou et al., 2019). 25 sensory neurons were reconstructed that could not be identified based on dye fill, MCFO, or published neurons. These neurons are referred to in this work as unknown and shown in [Figure 5 – figure supplement 2A-Y](#). We also identified interneurons that were not classified because sensory neurons were the focus of this study. The BMNs, JONs, and TPMNs are shown in [Figure 5 – figure supplement 1A,E](#). The GRNs and interneurons were excluded from further analysis in this study.

The 705 BMNs and 25 unknown sensory neurons were clustered based on NBLAST similarity scores as described above ([Figure 6 – figure supplement 1A,B](#)). We used these clusters in conjunction with manual anatomical inspection of light microscopy images to assign the BMNs to specific bristle populations. In particular, we compared the morphologies of the EM-reconstructed BMNs with dye-filled, stochastically-labeled (MCFO), and driver line-labeled BMNs ([Figure 6 – figure supplement 2A-N](#)). In some cases we did not have light microscopy images to enable direct matching of particular BMNs to their corresponding bristles. These BMNs were matched based on the morphology of BMNs innervating neighboring bristle populations (described for specific BMNs below). We also verified that the numbers of BMNs for each type were consistent with the numbers of their bristles ([Figure 6 – figure supplement 3B-F](#)). In the specific cases described below, we used additional evidence to match the different BMNs to their bristles, including comparing the reconstructed BMNs on both sides of the brain (OcciNv only, [Figure 6 – figure supplement 4A-D](#)) and based on common connectivity of the BMNs with their postsynaptic partners ([Figure 7F-L](#) and [Figure 7 – figure supplement 1](#)). A full list of the BMNs can be found in [Supplementary file 2](#).

The BMNs have been referred to in some previous work as external sensilla (es neurons). We adopted BMN here because it was similar to the widely adopted nomenclature for different sensory modalities, such as GRNs, ORNs, JONs. Further, it more explicitly links the neurons to the bristles.

3 BMNs that innervate the InOc bristles (BM-InOc neurons, [Figure 6D](#)) were identified based on comparison with MCFO data ([Figure 6 – figure supplement 2C](#)). 1 BMN that innervates the Oc bristle (BM-Oc neuron, [Figure 6E](#)) was identified based on comparison with dye fill data ([Figure 6 – figure supplement 2D](#)). 6 BMNs are proposed to innervate the Fr bristles (BM-Fr neurons, [Figure 6F](#)). The Fr bristles are located near the midline, immediately above the Ant bristles and below the InOc and Oc bristles. BMNs that innervate the Ant, InOc, and Oc bristles show very similar morphology with the BM-Fr neurons, including the ipsilateral and midline projections. 20 BMNs that innervate the Ant bristles (BM-Ant neurons, [Figure 6G](#)) were identified based on comparison with dye fill data ([Figure 6 – figure supplement 2E](#)). 3 BMNs that innervate the Or bristles (BM-Or neurons, [Figure 6H](#)) were identified based on comparison with dye fill data ([Figure 6 – figure supplement 2F](#)). 6 BMNs are proposed to innervate the FrOr bristles (BM-FrOr neurons, [Figure 6I](#)). The FrOr bristles are located

immediately ventral to the Or bristles. The BM-Or neurons show very similar morphology with the proposed BM-FrOr neurons, including an ipsilateral projection.

555 BMNs that innervate the InOm bristles (BM-InOm neurons, [Figure 6J](#)) were identified using previous descriptions of the projections of these neurons ([Hampel et al., 2017](#); [Zhang et al., 2020](#)). The similarity of the collective projections of the BM-InOm neurons with those labeled by the InOmBMN-LexA driver line are shown in [Figure 6 – figure supplement 2N](#). Clustering based on NBLAST similarity scores revealed that there were morphologically distinct groups of BM-InOm neurons. However, in contrast to other BMN types, the BM-InOm neurons were small with few relatively simple branches, and the clusters were likely due to relatively minor differences in these branches. For example, at the selected cut height (H=5) resulted in 11 different clusters ([Figure 6 – figure supplement 1A,B](#)). Additionally, while the BM-InOm neurons showed some differential postsynaptic connectivity based on cosine clustering results, these differences were relatively low ([Figure 7 – figure supplement 3A,B](#)). Therefore, the BM-InOm neurons are treated as a single group that innervates eye bristles in this work, while a future study will further examine the heterogeneity of the BM-InOm neurons.

18 EyeNv-projecting BMNs that innervate the Vib bristles (BM-Vib neurons, [Figure 6K](#)) were identified based on comparison with dye fill data ([Figure 6 – figure supplement 2G](#)). This includes 3 BMNs that project through the AntNv (BM-Vib (AntNv) neurons) that are morphologically similar to the BM-Vib (EyeNv) neurons. Based on 52A06-GAL4 labeling of BMNs on the head, BM-Vib (AntNv) neurons innervate smaller anterior Vib bristles that are lateral to Vib 1 and 2 bristles. In total, 15 BMNs projecting through the EyeNv (BM-Vib (EyeNv)) and 3 projecting through the AntNv (BM-Vib (AntNv)) were identified in the EM dataset.

15 BMNs are proposed to innervate the MaPa bristles (BM-MaPa neurons, [Figure 6L](#)), as their morphology matches a previous description ([Singh and Nayak, 1985](#)). It was difficult to distinguish between BM-MaPa and BM-Vib neurons based on morphology or nerve projection. However, all BM-MaPa neurons showed high cosine similarity in their postsynaptic connectivity to non-BMN neurons, and clustered together ([Figure 7 – figure supplement 2](#)).

BMNs that innervate the Taste bristles (BM-Taste neurons, [Figure 6M](#)) were identified based on their similarity to the MCFO images ([Figure 6 – figure supplement 2H](#)). We did not have dye fill or MCFO images of the BMNs that innervate the Hau bristles (BM-Hau neurons). These BMNs were presumed to show similar morphology to the BM-Taste neurons, given their close proximity. Indeed, visual inspection revealed 5 BMNs that had similar, yet slightly different morphology ([Figure 6N](#)). The axons appeared larger in diameter and showed a dorsal midline-crossing branch that was not found in BM-Taste neurons. Further, the 5 BM-Hau neurons showed high cosine similarity in their postsynaptic connectivity with non-BMN neurons ([Figure 7 – figure supplement 2](#)).

5 BMNs that project through the OcciNv that could innervate the Vt bristles (BM-Vt neurons, [Figure 6O](#)) were identified based on comparison with dye fill data ([Figure 6 – figure supplement 2I-K](#)). Because there are only 4 Vt bristles, the additional BMN in this group is proposed to innervate the PoOc bristle (BM-PoOc neuron). However, because we had no dye fills for the BM-PoOc neuron and limited dye fill examples of BM-Vt neurons, we included all 5 BMNs in the category BM-Vt/PoOc neurons. PoOc is hypothesized to project through the OcciNv and show similar morphology to the neighboring BM-Oc and -InOc neurons, and at least one BM-Vt/PoOc neuron has the expected morphology. Although the BM-InOc neuron is presumed to exist, we could not find evidence of a neuron innervating the PoOc bristle using any of the BMN driver lines reported in this study ([Figure 2 – figure supplement 1G](#)).

7 BMNs projecting through the OcciNv showed very similar morphology and were identified as innervating the dPoOr and dOcci bristles and named BM-dPoOr and BM-dOcci neurons, respectively. The morphology of the BM-dPoOr neurons was determined from MCFO experiments using dBMN-spGAL4 (**Figure 6 – figure supplement 2L**). MCFO experiments using pBMN-spGAL4 did not enable us to distinguish between BM-dPoOr and BM-dOcci neurons, as we could not tell which BMN came from which bristle population (**Figure 4S,T, Figure 4 – figure supplement 7B-D**). One difficulty in matching the reconstructed BM-PoOr and BM-dOcci neurons was that their numbers in the right brain hemisphere did not match what we expected from the numbers of their corresponding bristles. For example, there were only 4 tentatively assigned BM-dPoOr neurons, while we expected between 5 and 9 (**Figure 6 – figure supplement 3E**). Because the OcciNv contains only a small number of BMNs, we also reconstructed BMNs in the left brain hemisphere OcciNv (**Figure 6 – figure supplement 4A-D**). This revealed 4 additional BMNs on the left than on the right for the BM-dPoOr neurons, which matched what we expected (**Figure 6 – figure supplement 4D**). The numbers of the other BMN types that project through the OcciNv (i.e. BM-Vt/PoOc and BM-dOcci neurons) were the same in both hemispheres and matched with the expected number of bristles.

26 BMNs projecting through the EyeNv show very similar morphology and were identified as innervating the vPoOr and vOcci bristles. The vOcci-innervating BMNs (BM-vOcci neurons) were matched with MCFO images (**Figure 6 – figure supplement 2M**). However, we did not obtain dye fill or MCFO images of BMNs innervating the vPoOr bristles (BM-vPoOr neurons). These neurons were expected to show similar morphology with the BM-vOcci neurons based on their close proximity. Because we could not distinguish between these BMNs, they were grouped together and named (BM-vOcci/vPoOr neurons, **Figure 6R**).

Acknowledgements

We thank Brian Chen's lab at McGill University Health Centre for essential feedback on the development of the dye labeling protocol from this study. We thank the Princeton FlyWire team and members of the Murthy and Seung labs for development and maintenance of FlyWire (supported by BRAIN Initiative grant MH117815 to Murthy and Seung), and the Princeton EM proofreading team for contributing neuronal edits. We thank FlyWire users who contributed >10% proofreading of individual neurons or identification of neurons: Mala Murthy group (>10% of 31 neurons) Claire McKellar, Austin T Burke, Sarah Morejohn, Kyle Patrick Willie, Merlin Moore, Doug Bland, Ben Silverman, Jay Gager, Ryan Willie), Carlos Ribeiro group (>10% of 35 neurons) Ibrahim Tastekin, Raquel Barajas), the *Drosophila* Connectomics Group, Cambridge University (leads: M. Costa and G. Jefferis) for contributing to the tracing of FlyWire neurons (>10% of 20 neurons) Yijie Yin, Paul Brooks, Laia Serratosa Capdevila, Siqi Fang, A Javier, Paul Brooks), Rachel Wilson group (>10% of 8 neurons) Peter Gibb), Julie Simpson group (>10% of 2 neurons) Li Guo), Wolf Hütteroth group (>10% of 4 neurons), and Jinseop Kim group (>10% of 753 neurons) Hyunsoo Yim, Senongbong Yu, Chan Hyuk Kang, Taehyun Choe, Yeonju Nam, Tae Kim, Jinmook Kim).

Research reported in this publication was supported by the National Institute Of Neurological Disorders And Stroke of the National Institutes of Health under Award Number RF1NS121911. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. This work was also funded by the Whitehall Foundation (2017-12-69), Puerto Rico Science, Technology & Research Trust (2020-00195), NIMHD RCMI (MD007600), NIGMS-NIH RISE (R25GM061838), NIGMS-NIH RISE (5R25GM061151), NIH NIGMS (GM103642), and NSF (HRD-1736019).




Figure 1 - figure supplement 1.

Color-coded depth maps of the head.

(A-H) Example depth maps that were constructed from image z-stacks (see Materials and methods). Shown are the anterior (A,E), posterior (B,F), ventral (C,G), and dorsal (D,H) views of two different male heads. (A-D) Head shown in Figure 1 (Male 4). (E-H) Example of a different male head (Male 1). Colors indicate closer features in light blue, and increasingly more distant features in white, yellow, and dark red. Bristle names are indicated using abbreviations, whose full names are listed in Figure 1E.




Figure 1 - figure supplement 2.

Comparison of bristle numbers on male and female heads.

(A-D) Bristles on an anterior (A), posterior (B), ventral (C), and dorsal (D) female head (Female 1). (E) Table shows quantification of bristles on male and female heads and male/female

comparisons (male N=8, female N=4). Bristle numbers for each population are for one half of the head. Range, Avg., and SD are shown as described in [Figure 1](#). A student's t-test (2-tailed) was performed to compare the male/female bristle populations. InOc, PoOr (d+v), and Su bristles show a t-test $p \leq 0.05$, however the Bonferroni-adjusted α -value is 0.004 (14 comparisons). See **Supplementary file 1** for bristle counts for each head. Note that the PoOr (d+v) bristles were counted as a single group for comparing bristle numbers between males and females. Not determined (ND).





Figure 2 – figure supplement 1.

R52A06-GAL4 expression in head BMNs.

(A,B) Maximum intensity projections of the left and right halves of the anterior (A) and posterior (B) heads that are shown in [Figure 2C,D](#). BMNs are green and the cuticle is magenta. The BMNs on the right side of the midline (dotted line) for either the anterior or posterior heads were reconstructed and shown in [Figure 2C,D](#) (right). Anterior and posterior images are from two different heads. (C-F) Magnified images of A and B. Magnified areas are indicated by the

vertical lines. The different nerves are labeled with colored arrows. Location of unreconstructed portion of AntNv is indicated by an asterisk. Innervated bristles are indicated with white arrows. White arrowheads in D and F indicate partially reconstructed axons of BMNs innervating the InOm, Vib, and Taste bristles. (G) Posterior region of the head that includes the PoOc, Su, and dOccl bristles. Note that there are no GFP-labeled BMNs innervating the PoOc and Su bristles. (H) Brain and ventral nerve cord of R52A06-GAL4 expressing GFP and immunostained for GFP (green) and Bruchpilot (magenta) to label the neuropile. Scale bars all indicate 100 μ m.



Driver line expression in head BMNs.

(A-D') Maximum intensity projections of heads **(A-D)** and CNSs **(A'-D')** expressing GFP in BMNs that innervate different bristles. **A-D** were produced from the same confocal z-stacks that were used for the BMN reconstructions shown in [Figure 3D-G](#). Magnified views of the SEZs in **A'-D'** are shown in [Figure 3D'-G'](#). The expression patterns of the following driver lines are shown: InOmBMN-LexA **(A,A')**, dBMN-spGAL4 **(B,B')**, pBMN-spGAL4 **(C,C')**, and TasteBMN-spGAL4 **(D,D')**. Scale bars, 100 μ m **(A-C)**, 50 μ m **(D)**, 100 μ m **(A'-D')**. **(E)** Table showing the average number of GFP positive BMNs (and SD) that

innervate each bristle population from the indicated spGAL4 lines.




Figure 4 – figure supplement 1.

Overview of the dye filling technique and whole brain examples.

(A) Schematic of the dye filling technique used for labeling BMNs from specific bristles. BMNs were labeled with the anterograde dye, DiD. Each bristle was plucked from a head and DiD was pipetted onto the exposed socket to label the BMN innervating that bristle. **(B-E)** Whole brain examples of four fills shown in [Figure 4](#). Examples are from Ant 3 **(B)**, Vib 1 **(C)**, Or 1 **(D)**, and Vt 2 **(E)** bristle socket fills. Each filled BMN is magenta and the brain neuropile is labeled with pan neuronally expressed nSyb.GFP in green. Scale bar, 100 μ m.




Figure 4 – figure supplement 2.

Different fill trials for Oc and Or bristles.

(A) Oc and Or bristles whose associated BMNs were labeled by dye filling are indicated with labeled dots (dorsal view). The boxed area in the top image is shown magnified in the bottom image. (B-L) Anterior view of the SEZ projections of individual BMNs that innervate the Oc (B, C), Or 1 (D, E), Or 2 (G, H), and Or 3 (J, K) bristles. Two or three different flies were tested for each bristle (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. BMNs are oriented as if they are projecting from the right side of the head. Scale bar, 50 μm.




Figure 4 – figure supplement 3.

Different fill trials for Ant bristles.

(A) Ant bristles whose associated BMNs were labeled by dye filling are indicated with labeled dots (anterior view). The boxed area in the top image is shown magnified in the bottom image. (B-M) Anterior view of the SEZ projections of individual BMNs that innervate the Ant 1 (B, C, D), Ant 2 (E, F, G), Ant 3 (H, I, J), and Ant 4 (K, L, M) bristles. Three different flies were tested for each bristle (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. Scale bar, 50 μm.




Figure 4 - figure supplement 4.

Different fill trials for Vib bristles.

(A) Vib bristles whose associated BMNs were labeled by dye filling are indicated with labeled dots (anterior view). The boxed area in the top image is shown magnified in the bottom image. (B-M) Anterior view of the SEZ projections of individual BMNs that innervate the Vib 1 (B-D), Vib 2 (E-G), Vib 3 (H-J), and Vib 4 (K-M) bristles. Three different flies were tested for each bristle (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. Scale bar, 50 μm.




Figure 4 - figure supplement 5.

Different fill trials for Vt bristles.

(A) Vt bristles whose associated BMNs were labeled by dye filling are indicated with labeled dots (dorsal view). The boxed area in the top image is shown magnified in the bottom image. (B-E) Anterior view of the SEZ projections of individual BMNs that innervate the Vt 1 (B), Vt 2 (C,D), and Vt 3 (E) bristles. One or two different flies were tested for each bristle (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. Scale bar, 50 μm. Note: Vt 4 was not filled.




Figure 4 – figure supplement 6.

MCFO trials for BMNs innervating the InOc, dPoOr, and Vt bristles.

(A) Bristles whose associated BMNs were labeled by MCFO are indicated with labeled dots (dorsal view). The boxed area in the top image is shown magnified in the bottom image. (B-I) Anterior view of the SEZ projections of individual BMNs labeled by MCFO using dBMN-spGAL4. BMNs shown innervate the InOc (B,C), dPoOr (D-F), and Vt (G-I) bristles. At least two different flies were tested for each BMN (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. Scale bar, 50 μ m.




Figure 4 – figure supplement 7.

MCFO labeled trials for BMNs innervating the dOcci/dPoOr and vOcci bristles.

(A) Bristles whose associated BMNs were labeled by MCFO are indicated with labeled dots (posterior view). (B-J) Anterior view of the SEZ projections of individual BMNs labeled by MCFO using pBMN-spGAL4. BMNs shown innervate the dPoOr/dOcci (B-D) and vOcci (E-J) bristles. Three or six different flies were tested for each BMN (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. Scale bar, 50 μ m.




Figure 4 - figure supplement 8.

MCFO labeled trials for Taste bristles.

(A) Bristles whose associated BMNs were labeled by MCFO are indicated with labeled dots (posterior view). The boxed area in the top image is shown magnified in the bottom image. (B-K) Anterior view of the SEZ projections of individual BMNs labeled by MCFO using TasteBMN-spGAL4. BMNs shown innervate the Taste bristles. Ten different flies were tested (fly number indicated in upper right corner). Asterisk indicates the BMN example that is shown in Figure 4. Scale bar, 50 µm.




Figure 5 - figure supplement 1.

Reconstruction of mechanosensory neurons in different head nerves.

(A) Locations in each nerve where different segmented neurons in the EM volume were seeded for proof-reading and editing (black lines). (B-D) EM sections through each nerve at the locations shown in A with seeded neurons indicated with dots. Shown are AntNv (B), EyeNv/LabNv (C), and OcciNv (D) sections. Dot colors indicate the seeded neuron type, including BMNs projecting through the AntNv (blue), EyeNv (red), LabNv (brown), OcciNv (green), JONs (yellow), TPMNs (black), or other neurons (white). All neuron segments

were seeded for each nerve, with the following exceptions: 1) The Eye/LabNv had a bundle of soma tracts from an interneuron hemilineage crossing the seed plane that was excluded based on the morphology of their initial segmentation, and 2) Previously reconstructed JONs were excluded when seeding the AntNv (B, upper left), leaving a ventral-medial area of the nerve with previously undocumented neurons (bottom right). Scale bars, 2 µm. (E) Anterior (left) and lateral

(right) views of all reconstructed head mechanosensory neurons, including BMNs, JONs, and TPMNs (colors same as A-D). Arrows for each incoming nerve indicate projection direction.




Figure 5 – figure supplement 2.

Reconstructed sensory neurons that could not be assigned an identity (unknown sensory neurons).

(A-Y) SEZ projections (dorsal views) of unknown sensory neurons 001 (A), 002 (B), 003 (C), 004 (D), 005 (E), 006 (F), 007 (G), 008 (H), 009 (I), 010 (J), 011 (K), 012 (L), 013 (M), 014 (N), 015 (O), 016 (P), 017 (Q), 018 (R), 019 (S), 020 (T), 021 (U), 022 (V), 023 (W), 024 (X), and 025 (Y). Unknown sensory neurons project through the AntNv (A-H) or the EyeNv/LabNv (I-Y). Unknown neurons in S-U send their axons into the neck connective, and possibly descend to the ventral nerve cord.




Figure 6 – figure supplement 1.

NBLAST clustering of BMNs.

(A) Dendrogram of Ward clustered NBLAST similarity scores of all reconstructed sensory neurons. Identified nerve projection groups and BMN type assignments are color coded in the bars to the right (color IDs shown in the bottom right corner). The dendrogram was cut at $H=5$, resulting in 17 clusters indicated by numbers on the dendrogram. **Supplementary file 2** shows the NBLAST clusters when the dendrogram was cut at $H=1$, at which most of the BMN types are clustered individually. (B) Morphology of the BMNs in each NBLAST cluster shown in A, indicated by the same number (color-coded by type).




Figure 6 – figure supplement 2.

Matching NBLAST-clustered and individually-labeled BMNs.

(A-B) Bristles on the anterior (A) and posterior (B) head whose associated BMNs were labeled by dye filling, (fill), multicolor flipout (MCFO), or driver line expression (driver), and shown in C-N (top panel). (C-N) Top panels show representative images of the SEZ projections of labeled BMNs that innervate the bristle indicated in the bottom right corner (anterior view). All fill and MCFO trials for the different bristles are shown in Figure 4 – figure supplement 2-8.

Bottom panels show representative examples of the EM-reconstructed BMN types (types indicated in the bottom right corner). BMN types shown are BM-InOc (C), BM-Oc (D), BM-Ant (E), BM-Or (F), BM-Vib (G), BM-Taste (H), BM-Vt/PoOc (I), BM-Vt/PoOc (J), BM-Vt/PoOc (K), BM-dPoOr (L), BM-vOcci/vPoOr (M), and BM-InOm (N). Scale bar, 50 μm.




Figure 6 – figure supplement 3.

Evidence used to match the EM reconstructed BMNs with their bristles.

(A) NBLAST clusters were assigned as types that innervate particular bristle populations based on: 1) comparison of their morphology with images of dye-filled, MCFO-labeled, or published BMNs, 2) nerve projections, 3) proximity to BMNs with known morphology, and 4) common connectivity with postsynaptic partners. Postsynaptic connectivity clustering is shown in Figure 7 – figure supplement 2. The BMN types were validated because their numbers

matched the numbers of their corresponding bristles. The mismatch between the reconstructed BM-InOm neurons and the number of InOm bristles is likely due to the latter being an estimate (shown in **D**). Black shaded boxes indicate which evidence was used to match each BMN type. Detailed descriptions of the evidence that was used for assigning each BMN type can be found in Materials and methods. (**B-F**) Head bristle counts for each population and the number of reconstructed BMNs for each type. Shown are BMNs projecting through the AntNv (**B**), LabNv (**C**), EyeNv (**D**), and OcciNv (**E**), and BMNs innervating the Vib bristles (**F**). (**G**) R52A06-GAL4 expression in BM-Vib neurons on the head (green). Maximum intensity projections of the ventral head. Cuticle is magenta. BM-Vib neurons (green) that project to the brain through the AntNv or EyeNv are labeled with arrows.




Figure 6 – figure supplement 4.

EM reconstruction of OcciNv BMNs from both brain hemispheres.

(**A**) All BMNs entering the brain through the right (R) and left (L) OcciNv, color-coded by the BMN type. (**B,C**) BMN numbers are consistent across hemispheres for the BM-Vt/PoOc (**B**) and BM-dOcci (**C**) neurons (indicated below each hemisphere). (**D**) Reconstruction of the full occipital nerve on both hemispheres revealed twice as many BM-dPoOr neurons on the left than on the right.




Figure 7 – figure supplement 1.

BMN type synaptic counts and BMN/BMN connectivity.

(**A**) Number pre- and postsynaptic connections for each BMN type. Black lines indicate mean synapse number. (**B**) Synaptic connectivity among BMN types. The average synaptic count for each edge is shown on arrows indicating directionality of the connection (arrow width corresponds to the count). Total count of synapses between two given types was divided by the number of possible edges between the two types.




Figure 7 – figure supplement 2.

Cosine similarity clustering of BMN to non-BMN postsynaptic connectivity.

Heatmap of the cosine similarity among BMNs with the row and column order determined by clustering of the similarity scores. Postsynaptic partners excluded connections to other BMNs and connections with fewer than 6 synapses. Clustering of the rows and columns is the same as the dendrograms next to bars that indicate nerve group and BMN type (color-code on bottom right). Dendrogram cut height 4.5 resulted in 5 clusters (indicated by numbers on the top dendrogram). Morphologies of the neurons in each cluster are shown in Figure 7H-L.




Figure 7 – figure supplement 3.

Cosine similarity clustering of BM-InOm neurons in their connectivity with non-BMN postsynaptic partners.

(A) Heatmap of BM-InOm neuron cosine similarity with the row and column order determined by clustering of the similarity scores. Postsynaptic partners excluded connections to other BMNs and connections with fewer than 2 synapses. Clustering of the rows and columns is the same with dendrograms shown on the left and top. Dendrogram cut height 5.6 resulted in 7 clusters (indicated by numbers on the top dendrogram). **(B)** Dorsal view of the neuron morphologies in each cluster (cluster number indicated at the top left), revealed tiling of the BM-InOm innervation area in the brain. Clusters 1-2 mainly innervate an area located medial-dorsally, while clusters 6-7 innervate a lateral-ventral area. Clusters 3-5 send branches to both areas.




Figure 8 – figure supplement 1.

Ethograms of movements performed with activation of different BMNs.

Ethograms of manually scored videos showing the movements elicited with red-light induced activation. Ethograms of individual flies are stacked. Grooming movements (top plots) are indicated by different colors, including eye (magenta), dorsal head (blue), and ventral head (orange) grooming. Other elicited movements (bottom plots) include backward motion (black) and head nodding (gray). Gray bars indicate a 5 second red-light stimulus. Most driver lines were tested using 30 second interstimulus intervals, while pBMN-spGAL4 elicited more reliable behavior using 10 second intervals. Movements are mutually exclusive except head nodding. Gray bars indicate a 5 second red-light stimulus.

Supplementary file 1. Quantification of bristle populations on male and female heads.

Table 1. Bristles counted using white light-illuminated male and female heads. Results of a two-tailed t-test comparing male and female bristle numbers are also shown. Table 2 shows counts of PoOr bristles innervated by BMNs that project through the OcciNv (dPoOr bristles) or the EyeNv (vPoOr bristles). BMNs were labeled by R52A06-GAL4. Table 3 shows vOcci bristles counted based on having an associated BMN labeled in the pBMN-spGAL4 (R28D07-AD $\cap$ VT050279-DBD) pattern (males). Table 4 shows a summary of published bristle numbers.

Supplementary file 2. Table of EM-reconstructed BMNs, JONs, and TPMNs. Table includes FlyWire.ai neuron XYZ coordinates and IDs, nerve groups, names, types, NBLAST clusters, and cosine similarity clusters.

Figure 8 – source data 1. Numerical data used for producing each box plot. Rows correspond to individual flies. Columns indicate the percent time each fly spent performing different movements, including eye, dorsal, and ventral head grooming, nodding, and backward motions.

Video 1. Optogenetic activation of dorsal head BMNs elicits aimed dorsal head grooming. CsChrimson was expressed in BMNs targeted by the dBMN-spGAL4 driver line. Infrared light in the bottom right corner indicates when the red light was on to activate the targeted BMNs. Note that head nodding movements and backward motions are also elicited.

Video 2. Optogenetic activation of BM-Taste neurons elicits aimed proboscis and ventral head grooming. CsChrimson was expressed in BMNs targeted by the TasteBMN-

spGAL4 driver line. Infrared light in the bottom right corner indicates when the red light was on to activate the targeted BMNs.

Video 3. Optogenetic activation of BM-InOm neurons elicits eye and dorsal head grooming. CsChrimson was expressed in BMNs targeted by the InOmBMN-LexA driver line. Infrared light in the bottom right corner indicates when the red light was on to activate the targeted BMNs. Note that head nodding movements and backward motions are also elicited.

Video 4. Optogenetic activation of posterior head BMNs elicits head nodding. CsChrimson was expressed in BMNs targeted by the pBMN-spGAL4 driver line. Infrared light in the bottom right corner indicates when the red light was on to activate the targeted BMNs. Note that dorsal head grooming movements are also elicited (not shown in video).

Video 5. Optogenetic stimulus in control flies. Control fly was exposed to the same red-light stimulus shown in **Videos 1-3**. The infrared light in the bottom right corner indicates when the red light was on.

References

Abraira VE , Ginty DD (2013) **The sensory neurons of touch** *Neuron* **79**:618–639
<https://doi.org/10.1016/j.neuron.2013.07.051>

Adibi M (2019) **Whisker-Mediated Touch System in Rodents: From Neuron to Behavior** *Frontiers Syst Neurosci* **13**
<https://doi.org/10.3389/fnsys.2019.00040>

Averbeck BB , Chafee MV , Crowe DA , Georgopoulos AP (2002) **Parallel processing of serial movements in prefrontal cortex** *Proceedings of the National Academy of Sciences* **99**:13172–13177
<https://doi.org/10.1073/pnas.162485599>

Bates AS , Manton JD , Jagannathan SR , Costa M , Schlegel P , Rohlfing T , Jefferis GS (2020) **The natverse, a versatile toolbox for combining and analysing neuroanatomical data** *Elife* **9**
<https://doi.org/10.7554/elife.53350>

Berkowitz A , Laurent G (1996) **Central generation of grooming motor patterns and interlimb coordination in locusts** *The Journal of Neuroscience* **16**:8079–8091

Bodenstein D , Miller A , Cooper KW , Poulson DF , Ferris GF , Sonnenblick BP , Spenser WP (1994) **Biology of Drosophila**

Bohland JW , Bullock D , Guenther FH (2010) **BohlandJW, BullockD, GuentherFH. 2010. Neural representations and mechanisms for the performance of simple speech sequences. 22:1504–1529. doi:10.1162/jocn.2009.21306** *Neural representations and mechanisms for the performance of simple speech sequences* **22**:1504–1529
<https://doi.org/10.1162/jocn.2009.21306>

Brown AG , Rose PK , Snow PJ (1977) **The morphology of hair follicle afferent fibre collaterals in the spinal cord of the cat** *J Physiology* **272**:779–797
<https://doi.org/10.1113/jphysiol.1977.sp012073>

Buhmann J , Sheridan A , Malin-Mayor C , Schlegel P , Gerhard S , Kazimiers T , Krause R , Nguyen TM , Heinrich L , Lee W-CA , Wilson R , Saalfeld S , Jefferis GSXE , Bock DD , Turaga SC , Cook M , Funke J (2021) **Automatic detection of synaptic partners in a whole-brain Drosophila electron microscopy data set** *Nat Methods* **18**:771–774
<https://doi.org/10.1038/s41592-021-01183-7>

Bullock D (2004) **Adaptive neural models of queuing and timing in fluent action** *Trends in Cognitive Sciences* **8**:426–433
<https://doi.org/10.1016/j.tics.2004.07.003>

Burg M , Wu C (1986) **Differentiation and central projections of peripheral sensory cells with action-potential block in Drosophila mosaics** *J Neurosci* **6**:2968–2976
<https://doi.org/10.1523/jneurosci.06-10-02968.1986>

Burg MG , Hanna L , Kim YT , Wu CF (1993) **Development and maintenance of a simple reflex circuit in small-patch mosaics of Drosophila: effects of altered neuronal function and developmental arrest** *Journal of Neurobiology* **24**:803–823
<https://doi.org/10.1002/neu.480240608>

Burg MG , Wu CF (1989) **Central projections of peripheral mechanosensory cells with increased excitability in Drosophila mosaics** *Developmental Biology* **131**:505–514
[https://doi.org/10.1016/s0012-1606\(89\)80021-1](https://doi.org/10.1016/s0012-1606(89)80021-1)

Cande J , Namiki S , Qiu J , Korff W , Card GM , Shaevitz JW , Stern DL , Berman GJ (2018) **Optogenetic dissection of descending behavioral control in Drosophila** *Elife* **7**
<https://doi.org/10.7554/elife.34275>

Chen BE , Kondo M , Garnier A , Watson FL , Püettmann-Holgado R , Lamar DR , Schmucker D (2006) **The Molecular Diversity of Dscam Is Functionally Required for Neuronal Wiring Specificity in Drosophila** *Cell* **125**:607–620
<https://doi.org/10.1016/j.cell.2006.03.034>

Corfas G , Dudai Y (1990) **Adaptation and fatigue of a mechanosensory neuron in wild-type Drosophila and in memory mutants** *Journal of Neuroscience* **10**:491–499

Corfas G , Dudai Y (1989) **Habituation and dishabituation of a cleaning reflex in normal and mutant Drosophila** *The Journal of Neuroscience* **9**:56–62

Costa M , Manton JD , Ostrovsky AD , Prohaska S , Jefferis GSXE (2016) **NBLAST: Rapid, Sensitive Comparison of Neuronal Structure and Construction of Neuron Family Databases** *Neuron*
<https://doi.org/10.1016/j.neuron.2016.06.012>

Dawkins R , Dawkins M (1976) **Hierarchical organization and postural facilitation: Rules for grooming in flies** *Animal Behaviour* **24**:739–755
[https://doi.org/10.1016/s0003-3472\(76\)80003-6](https://doi.org/10.1016/s0003-3472(76)80003-6)

Dionne H , Hibbard K , Cavallaro A , Kao JC , bioRxiv GR (2017) **2017. Genetic reagents for making split-GAL4 lines in Drosophila** *bioRxiv.org*
<https://doi.org/10.1101/197509>

Dorkenwald S , McKellar CE , Macrina T , Kemnitz N , Lee K , Lu R , Wu J , Popovych S , Mitchell E , Nehoran B , Jia Z , Bae JA , Mu S , Ih D , Castro M , Ogedengbe O , Halageri A , Kuehner K , Sterling AR , Ashwood Z , Zung J , Brittain D , Collman F , Schneider-Mizell C , Jordan C , Silversmith W , Baker C , Deutsch D , Encarnacion-Rivera L , Kumar S , Burke A , Bland D , Gager J , Hebditch J , Koolman S , Moore M , Morejohn S , Silverman B , Willie K , Willie R , Yu S , Murthy M , Seung HS (2022) **FlyWire: online community for whole-brain connectomics** *Nat Methods* **19**:119–128
<https://doi.org/10.1038/s41592-021-01330-0>

Eckstein N , Bates AS , Du M , Hartenstein V , Jefferis GSXE , Funke J. (2020) **Neurotransmitter Classification from Electron Microscopy Images at Synaptic Sites in Drosophila** *Biorxiv*
<https://doi.org/10.1101/2020.06.12.148775>

Engert S , Sterne GR , Bock DD , Scott K (2022) **Drosophila gustatory projections are segregated by taste modality and connectivity** *Elife* **11**
<https://doi.org/10.7554/elife.78110>

Falk R , Bleiser-Avivi N , Atidia J (1976) **Labellar taste organs of Drosophila melanogaster** *J Morphol* **150**:327–341
<https://doi.org/10.1002/jmor.1051500206>

Feng L , Zhao T , Kim J (2015) **neuTube 1.0: A New Design for Efficient Neuron Reconstruction Software Based on the SWC Format** *eNeuro* **2**
<https://doi.org/10.1523/eneuro.0049-14.2014>

Galili DS , Jefferis GS , Costa M (2022) **Connectomics and the neural basis of behaviour** *Curr Opin Insect Sci* **54**
<https://doi.org/10.1016/j.cois.2022.100968>

Ghysen A (1980) **The projection of sensory neurons in the central nervous system of Drosophila: choice of the appropriate pathway** *Developmental Biology* **78**:521–541
[https://doi.org/10.1016/0012-1606\(80\)90351-6](https://doi.org/10.1016/0012-1606(80)90351-6)

Guo L , Zhang N , Simpson JH (2022) **Descending neurons coordinate anterior grooming behavior in Drosophila** *Curr Biol*
<https://doi.org/10.1016/j.cub.2021.12.055>

Hagedorn J , Hailpern J , Karahalios KG (2008) **VCode and VData: illustrating a new framework for supporting the video annotation workflow** *Extended Abstracts of AVI* **317**
<https://doi.org/10.1145/1385569.1385622>

Hampel S , Chung P , McKellar CE , Hall D , Looger LL , Simpson JH (2011) **Drosophila Brainbow: a recombinase-based fluorescence labeling technique to subdivide neural expression patterns** :253–259
<https://doi.org/10.1038/nmeth.1566>

Hampel S , Eichler K , Yamada D , Bock D , Kamikouchi A , Seeds AM (2020) **Distinct subpopulations of mechanosensory chordotonal organ neurons elicit grooming of the fruit fly antennae** *Elife* 9
<https://doi.org/10.7554/elife.59976>

Hampel S , Eichler K , Yamada D , Kim H , Horigome M , Franconville R , Bock DD , Kamikouchi A , Seeds AM (2020) **Convergence of distinct subpopulations of mechanosensory neurons onto a neural circuit that elicits grooming** *Biorxiv*
<https://doi.org/10.1101/2020.06.08.141341>

Hampel S , Franconville R , Simpson JH , Seeds AM (2015) **A neural command circuit for grooming movement control** *eLife* 4
<https://doi.org/10.7554/elife.08758>

Hampel S , McKellar CE , Simpson JH , Seeds AM (2017) **Simultaneous activation of parallel sensory pathways promotes a grooming sequence in Drosophila** *eLife* 6
<https://doi.org/10.7554/elife.28804>

Heinrich L , Funke J , Pape C , Nunez-Iglesias J , Saalfeld S (2018) **Synaptic Cleft Segmentation in Non-isotropic Volume Electron Microscopy of the Complete Drosophila Brain., Medical Image Computing and Computer Assisted Intervention – MICCAI 2018. MICCAI 2018 Lecture Notes in Computer Science**
https://doi.org/10.1007/978-3-030-00934-2_36

Homberg U , Christensen TA , Hildebrand JG (1989) **Structure and function of the deutocerebrum in insects** *Annual Review of Entomology* 34:477–501
<https://doi.org/10.1146/annurev.en.34.010189.002401>

Honegger H-W (1977) **Interommatidial hair receptor axons extending into the ventral nerve cord in the cricket *Gryllus campestris*** *Cell and tissue research*

Honegger HW , Reif H , Müller W (1979) **Sensory mechanisms of eye cleaning behavior in the cricket *Gryllus campestris*** *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology* 129:247–256

Houghton G , Hartley T (1995) **Parallel models of serial behavior: Lashley revisited** *Psyche* 2:1–25

Ishikawa Y , Okamoto N , Nakamura M , Kim H , Kamikouchi A (2017) **Anatomic and Physiologic Heterogeneity of Subgroup-A Auditory Sensory Neurons in Fruit Flies** *Frontiers in neural circuits* 11
<https://doi.org/10.3389/fncir.2017.00046>

Jefferis GSXE , Potter CJ , Chan AM , Marin EC , Rohlfsing T , Luo L (2007) **Comprehensive Maps of Drosophila Higher Olfactory Centers: Spatially Segregated Fruit and Pheromone Representation** *Cell* **128**:1187–1203
<https://doi.org/10.1016/j.cell.2007.01.040>

Jenett AA , Rubin GM , Ngo T-TB , Shepherd DD , Murphy CC , Dionne HH , Pfeiffer BD , Cavallaro AA , Hall DD , Jeter JJ , Iyer NN , Fetter DD , Hausenfluck JH , Peng HH , Trautman ET , Svirskas RR , Myers EW , Iwinski ZR , Aso YY , Depasquale GM , Enos AA , Hulamm PP , Lam SCB , Li H-H , Lavery TR , Long FF , Qu LL , Murphy SD , Rokicki KK , Safford TT , Shaw KK , Simpson JH , Sowell AA , Tae SS , Yu YY , Zugates CT (2012) **A GAL4-driver line resource for Drosophila neurobiology** *Cell Reports* **2**:991–1001
<https://doi.org/10.1016/j.celrep.2012.09.011>

Jeong YT , Oh SM , Shim J , Seo JT , Kwon JY , Moon SJ (2016) **Mechanosensory neurons control sweet sensing in Drosophila** *Nature communications* **7**
<https://doi.org/10.1038/ncomms12872>

Johnson SE , Murphey RK (1985) **The afferent projection of mesothoracic bristle hairs in the cricket, Acheta domesticus** *J Comp Physiology* **156**:369–379
<https://doi.org/10.1007/bf00610729>

Kaas JH (1997) **Topographic maps are fundamental to sensory processing** *Brain Research Bulletin* **44**:107–112
[https://doi.org/10.1016/s0361-9230\(97\)00094-4](https://doi.org/10.1016/s0361-9230(97)00094-4)

Kamikouchi A , Inagaki HK , Effertz T , Hendrich O , Fiala A , Göpfert MC , Ito K (2009) **The neural basis of Drosophila gravity-sensing and hearing** *Nature* **458**:165–171
<https://doi.org/10.1038/nature07810>

Kamikouchi A , Shimada T , Ito K (2006) **Comprehensive classification of the auditory sensory projections in the brain of the fruit fly Drosophila melanogaster** *The Journal of Comparative Neurology* **499**:317–356
<https://doi.org/10.1002/cne.21075>

Kays I , Cvetkovska V , Chen BE (2014) **Structural and functional analysis of single neurons to correlate synaptic connectivity with grooming behavior** *Nature protocols* **9**:1–10
<https://doi.org/10.1038/nprot.2013.157>

Kent KS , Levine RB (1988) **Neural control of leg movements in a metamorphic insect: Sensory and motor elements of the larval thoracic legs in Manduca sexta** *J Comp Neurol* **271**:559–576
<https://doi.org/10.1002/cne.902710407>

Kim H , Horigome M , Ishikawa Y , Li F , Lauritzen JS , Card G , Bock DD , Kamikouchi A (2020) **Wiring patterns from auditory sensory neurons to the escape and song-relay pathways in fruit flies** *J Comp Neurol*
<https://doi.org/10.1002/cne.24877>

Klapoetke NC , Murata Y , Kim SS , Pulver SR , Birdsey-Benson A , Cho YK , Morimoto TK , Chuong AS , Carpenter EJ , Tian Z , Wang J , Xie Y , Yan Z , Zhang Y , Chow BY , Surek B , Melkonian M , Jayaraman V , Constantine-Paton M , Wong GK-S , Boyden ES (2014) **KlapoetkeNC, MurataY, KimSS, PulverSR, Birdsey-BensonA, ChoYK, MorimotoTK, ChuongAS, CarpenterEJ, TianZ, WangJ, XieY, YanZ, ZhangY, ChowBY, SurekB, MelkonianM, JayaramanV, Constantine-PatonM, WongGK-S, BoydenES. 2014. Independent optical excitation of distinct neural populations** *11*:338–346
doi:10.1038/nmeth.2836 *Independent optical excitation of distinct neural populations* **11**:338–346
<https://doi.org/10.1038/nmeth.2836>

Lashley KS (1951) **Cerebral mechanisms in behavior**

Lehnert BP , Baker AE , Gaudry Q , Chiang A-S , Wilson RI (2013) **Distinct roles of TRP channels in auditory transduction and amplification in *Drosophila*** *Neuron* **77**:115–128
<https://doi.org/10.1016/j.neuron.2012.11.030>

Li J , Zhang W , Guo Z , Wu S , Jan LY , Jan YN (2016) **A Defensive Kicking Behavior in Response to Mechanical Stimuli Mediated by *Drosophila* Wing Margin Bristles** *Journal of Neuroscience* **36**:11275–11282
<https://doi.org/10.1523/jneurosci.1416-16.2016>

Luan H , Peabody NC , Vinson CR , White BH (2006) **Refined spatial manipulation of neuronal function by combinatorial restriction of transgene expression** *Neuron* **52**:425–436
<https://doi.org/10.1016/j.neuron.2006.08.028>

Mamiya A , Dickinson MH (2015) **Antennal mechanosensory neurons mediate wing motor reflexes in flying *Drosophila*** *Journal of Neuroscience* **35**:7977–7991
<https://doi.org/10.1523/jneurosci.0034-15.2015>

Mamiya A , Straw AD , Tómasson E , Dickinson MH (2011) **Active and passive antennal movements during visually guided steering in flying *Drosophila*** *Journal of Neuroscience* **31**:6900–6914
<https://doi.org/10.1523/jneurosci.0498-11.2011>

Matheson T (1997) **Hindleg targeting during scratching in the locust** *The Journal of experimental biology* **200**:93–100

Matsuo E , Yamada D , Ishikawa Y , Asai T , Ishimoto H , Kamikouchi A (2014) **Identification of novel vibration- and deflection-sensitive neuronal subgroups in Johnston's organ of the fruit fly** *Frontiers in Physiology* **5**
<https://doi.org/10.3389/fphys.2014.00179>

Melzig J , Buchner S , Wiebel F , Wolf R , Buchner E , Burg M , Pak WL (1996) **Genetic depletion of histamine from the nervous system of *Drosophila* eliminates specific visual and mechanosensory behavior** *Journal of Comparative Physiology A* **179**:763–773

- Merritt DJ , Murphey RK (1992) **Projections of leg proprioceptors within the CNS of the fly *Phormia* in relation to the generalized insect ganglion** *The Journal of Comparative Neurology* **322**:16–34
<https://doi.org/10.1002/cne.903220103>
- Mitchell BK , Itagaki H , Rivet MP (1999) **Peripheral and central structures involved in insect gustation** *Microscopy Research and Technique* **47**:401–415
[https://doi.org/10.1002/\(sici\)1097-0029\(19991215\)47](https://doi.org/10.1002/(sici)1097-0029(19991215)47)
- Miyazaki T , Ito K (2010) **Neural architecture of the primary gustatory center of *Drosophila melanogaster* visualized with GAL4 and LexA enhancer-trap systems** *Journal of comparative neurology* **518**:4147–4181
<https://doi.org/10.1002/cne.22433>
- Mueller JM , Ravbar P , Simpson JH , Carlson JM (2019) ***Drosophila melanogaster* grooming possesses syntax with distinct rules at different temporal scales** *Plos Comput Biol* **15**
<https://doi.org/10.1371/journal.pcbi.1007105>
- Murphey RK , Possidente D , Pollack G , Merritt DJ (1989) **Modality-specific axonal projections in the CNS of the flies *Phormia* and *Drosophila*** *The Journal of Comparative Neurology* **290**:185–200
<https://doi.org/10.1002/cne.902900203>
- Murphey RK , Possidente DR , Vandervorst P , Ghysen A (1989) **Compartments and the topography of leg afferent projections in *Drosophila*** *Journal of Neuroscience* **9**:3209–3217
- Mushiake H , Saito N , Sakamoto K , Itoyama Y , Tanji J (2006) **Activity in the lateral prefrontal cortex reflects multiple steps of future events in action plans** *Neuron* **50**:631–641
<https://doi.org/10.1016/j.neuron.2006.03.045>
- Nayak SV , Singh RN (1983) **Sensilla on the tarsal segments and mouthparts of adult *Drosophila melanogaster* meigen (Diptera : Drosophilidae)** *Int J Insect Morphol Embryology* **12**:273–291
[https://doi.org/10.1016/0020-7322\(83\)90023-5](https://doi.org/10.1016/0020-7322(83)90023-5)
- Nern A , Pfeiffer BD , Rubin GM (2015) **Optimized tools for multicolor stochastic labeling reveal diverse stereotyped cell arrangements in the fly visual system** *Proceedings of the National Academy of Sciences* **112**
<https://doi.org/10.1073/pnas.1506763112>
- Newland PL (1991) **Morphology and somatotopic organisation of the central projections of afferents from tactile hairs on the hind leg of the locust** *J Comp Neurol* **312**:493–508
<https://doi.org/10.1002/cne.903120402>
- Newland PL , Burrows M (1997) **Processing of tactile information in neuronal networks controlling leg movements of the locust** *Journal of insect physiology* **43**:107–123

Newland PL , Rogers SM , Gaaboub I , Matheson T (2000) **Parallel somatotopic maps of gustatory and mechanosensory neurons in the central nervous system of an insect** *The Journal of Comparative Neurology* **425**:82–96
[https://doi.org/10.1002/1096-9861\(20000911\)425](https://doi.org/10.1002/1096-9861(20000911)425)

Page KLK , Matheson TT (2004) **Wing hair sensilla underlying aimed hindleg scratching of the locust** *Journal of Experimental Biology* **207**:2691–2703
<https://doi.org/10.1242/jeb.01096>

Patella P , Wilson RI (2018) **Functional Maps of Mechanosensory Features in the Drosophila Brain** *Current biology : CB* **28**:1189–1203
<https://doi.org/10.1016/j.cub.2018.02.074>

Pfeiffer BD , Jenett A , Hammonds AS , Ngo T-TB , Misra S , Murphy C , Scully A , Carlson JW , Wan KH , Lavery TR , Mungall C , Svirska R , Kadonaga JT , Doe CQ , Eisen MB , Celniker SE , Rubin GM (2008) **Tools for neuroanatomy and neurogenetics in Drosophila** *Proceedings of the National Academy of Sciences* **105**:9715–9720
<https://doi.org/10.1073/pnas.0803697105>

Pfeiffer BD , Ngo T-TB , Hibbard KL , Murphy C , Jenett A , Truman JW , Rubin GM (2010) **Refinement of tools for targeted gene expression in Drosophila** *Genetics* **186**:735–755
<https://doi.org/10.1534/genetics.110.119917>

Pflüger H-J (1980) **The Function of Hair Sensilla on the Locust's Leg: the Role Of tibial hairs** *J Exp Biol* **87**:163–176
<https://doi.org/10.1242/jeb.87.1.163>

Phelps JS , Hildebrand DGC , Graham BJ , Kuan AT , Thomas LA , Nguyen TM , Buhmann J , Azevedo AW , Sustar A , Agrawal S , Liu M , Shanny BL , Funke J , Tuthill JC , Lee W-CA (2021) **Reconstruction of motor control circuits in adult Drosophila using automated transmission electron microscopy** *Cell* **184**:759–774
<https://doi.org/10.1016/j.cell.2020.12.013>

Phillis RW , Bramlage AT , Wotus C , Whittaker A , Gramates LS , Seppala D , Farahanchi F , Caruccio P , Murphey RK (1993) **Isolation of mutations affecting neural circuitry required for grooming behavior in Drosophila melanogaster** *Genetics* **133**:581–592

Ready DF , Hanson TE , Benzer S (1976) **Development of the Drosophila retina, a neurocrystalline lattice** *Dev Biol* **53**:217–240
[https://doi.org/10.1016/0012-1606\(76\)90225-6](https://doi.org/10.1016/0012-1606(76)90225-6)

Salvaterra PM , Kitamoto T (2001) **Drosophila cholinergic neurons and processes visualized with Gal4/UAS-GFP** *Gene expression patterns* **1**:73–82
[https://doi.org/10.1016/s1567-133x\(01\)00011-4](https://doi.org/10.1016/s1567-133x(01)00011-4)

Sánchez-Alcañiz JA , Zappia G , Marion-Poll F , Benton R (2017) **A mechanosensory receptor required for food texture detection in Drosophila** *Nature communications* **8**
<https://doi.org/10.1038/ncomms14192>

Seeds AM , Ravbar P , Chung P , Hampel S , Midgley FM , Mensh BD , Simpson JH (2014) **A suppression hierarchy among competing motor programs drives sequential grooming in *Drosophila*** *eLife* **3**

Shanbhag S , Park S-K , Pikielny C , Steinbrecht R (2001) **Gustatory organs of *Drosophila melanogaster*: fine structure and expression of the putative odorant-binding protein PBPRP2** *Cell Tissue Res* **304**:423–437
<https://doi.org/10.1007/s004410100388>

Shannon P , Markiel A , Ozier O , Baliga NS , Wang JT , Ramage D , Amin N , Schwikowski B , Ideker T (2003) **Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks** *Genome Res* **13**:2498–2504
<https://doi.org/10.1101/gr.1239303>

Singh RN , Nayak SV (1985) **Fine structure and primary sensory projections of sensilla on the maxillary palp of *Drosophila melanogaster* Meigen (Diptera : Drosophilidae)** *Int J Insect Morphol Embryology* **14**:291–306
[https://doi.org/10.1016/0020-7322\(85\)90044-3](https://doi.org/10.1016/0020-7322(85)90044-3)

Smith SA , Shepherd D (1998) **Central afferent projections of proprioceptive sensory neurons in *Drosophila* revealed with the enhancer trap technique** *The Journal of Comparative Neurology* **364**:311–323

Steyskal GC (1976) **The terminology of bristles on the upper back of the head in the higher diptera** *Journal of the Kansas Entomological Society* **49**:55–159

Stocker RF (1994) **The organization of the chemosensory system in *Drosophila melanogaster*: a review** *Cell and tissue research* **275**:3–26

Stocker RF , Schorderet M (1981) **Cobalt filling of sensory projections from internal and external mouthparts in *Drosophila*** *Cell Tissue Res* **216**:513–523
<https://doi.org/10.1007/bf00238648>

Suver MP , Matheson AMM , Sarkar S , Damiata M , Schoppik D , Nagel KI (2019) **Encoding of Wind Direction by Central Neurons in *Drosophila*** *Neuron* **102**:828–842
<https://doi.org/10.1016/j.neuron.2019.03.012>

Szebenyi AL (1969) **Cleaning behaviour in *Drosophila melanogaster*** *Animal Behaviour* **17**:641–651
[https://doi.org/10.1016/s0003-3472\(69\)80006-0](https://doi.org/10.1016/s0003-3472(69)80006-0)

Thei J (1979) **Mechanoreceptive bristles on the head of the blowfly: Mechanics and electrophysiology of the macrochaetae** *Journal of Comparative Physiology A* **132**:55–68
<https://doi.org/10.1007/bf00617732>

Thivierge J-P , Marcus GF (2007) **The topographic brain: from neural connectivity to cognition** *Trends in Neurosciences* **30**:251–259
<https://doi.org/10.1016/j.tins.2007.04.004>

- Tirián L , Dickson B (2017) **The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the Drosophila nervous system** *bioRxiv*
<https://doi.org/10.1101/198648>
- Tootoonian SS , Coen PP , Kawai RR , Murthy MM (2012) **Neural representations of courtship song in the Drosophila brain** *Journal of Neuroscience* **32**:787–798
<https://doi.org/10.1523/jneurosci.5104-11.2012>
- Tsubouchi A , Yano T , Yokoyama TK , Murtin C , Otsuna H , Ito K (2017) **Topological and modality-specific representation of somatosensory information in the fly brain** *Science* **358**:615–623
<https://doi.org/10.1126/science.aan4428>
- Tuthill JC , Wilson RI (2016) **Parallel Transformation of Tactile Signals in Central Circuits of Drosophila** *Cell* **164**:1046–1059
<https://doi.org/10.1016/j.cell.2016.01.014>
- Tuthill JC , Wilson RI (2016) **Mechanosensation and Adaptive Motor Control in Insects** *Current biology : CB* **26**
<https://doi.org/10.1016/j.cub.2016.06.070>
- Tyrer NM , Bacon JP , Davies CA (1979) **Sensory projections from the wind-sensitive head hairs of the locust Schistocerca gregaria. Distribution in the central nervous system** *Cell and tissue research* **203**:79–92
- Usui-Ishihara AA , Ghysen A , Kimura K-I (1995) **Peripheral Axonal Pathway and Cleaning Behavior Are Correlated in Drosophila Microchaetes** *Developmental Biology* **167**:398–401
<https://doi.org/10.1006/dbio.1995.1034>
- Vandervorst P , Ghysen A (1980) **Genetic control of sensory connections in Drosophila** *Nature* **286**:65–67
<https://doi.org/10.1038/286065a0>
- Vaughan AG , Zhou C , Manoli DS , Baker BS (2014) **Neural pathways for the detection and discrimination of conspecific song in D. melanogaster** *Current biology : CB* **24**:1039–1049
<https://doi.org/10.1016/j.cub.2014.03.048>
- Walker RG (2000) **A Drosophila Mechanosensory Transduction Channel** *Science* **287**:2229–2234
<https://doi.org/10.1126/science.287.5461.2229>
- Yang T , Yuan Z , Liu C , Liu T , Zhang W (2021) **A neural circuit integrates pharyngeal sensation to control feeding** *Cell Reports* **37**
<https://doi.org/10.1016/j.celrep.2021.109983>
- Yasuyama K , Salvaterra PM (1999) **Localization of choline acetyltransferase-expressing neurons in Drosophila nervous system** *Microscopy Research and Technique* **45**:65–79
[https://doi.org/10.1002/\(sici\)1097-0029\(19990415\)45](https://doi.org/10.1002/(sici)1097-0029(19990415)45)

Yorozu S , Wong A , Fischer BJ , Dankert H , Kernan MJ , Kamikouchi A , Ito K , Anderson DJ (2009) **Distinct sensory representations of wind and near-field sound in the *Drosophila* brain** *Nature* **458**:201–205
<https://doi.org/10.1038/nature07843>

Zack SS , Bacon JJ (1981) **Interommatidial sensilla of the praying mantis: their central neural projections and role in head-cleaning behavior** *Journal of Neurobiology* **12**:55–65
<https://doi.org/10.1002/neu.480120105>

Zhang N , Guo L , Simpson JH (2020) **Spatial Comparisons of Mechanosensory Information Govern the Grooming Sequence in *Drosophila*** *Curr Biol* **30**:988–1001
<https://doi.org/10.1016/j.cub.2020.01.045>

Zhang YV , Aikin TJ , Li Z , Montell C (2016) **The Basis of Food Texture Sensation in *Drosophila*** *Neuron* **91**:863–877
<https://doi.org/10.1016/j.neuron.2016.07.013>

Zheng Z , Lauritzen JS , Perlman E , Robinson CG , Nichols M , Milkie D , Torrens O , Price J , Fisher CB , Sharifi N , Calle-Schuler SA , Kmecova L , Ali IJ , Karsh B , Trautman ET , Bogovic JA , Hanslovsky P , Jefferis GSXE , Kazhdan M , Khairy K , Saalfeld S , Fetter RD , Bock DD (2018) **A Complete Electron Microscopy Volume of the Brain of Adult *Drosophila melanogaster*** *Cell* **174**:730–743
<https://doi.org/10.1016/j.cell.2018.06.019>

Zhou Y , Cao L-H , Sui X-W , Guo X-Q , Luo D-G (2019) **Mechanosensory circuits coordinate two opposing motor actions in *Drosophila* feeding** *Science advances* **5**
<https://doi.org/10.1126/sciadv.aaw5141>

Author information

Katharina Eichler

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico
ORCID iD: [0000-0002-7833-8621](https://orcid.org/0000-0002-7833-8621)

Stefanie Hampel

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico

For correspondence: seeds.andrew@gmail.com

ORCID iD: [0000-0001-8287-549X](https://orcid.org/0000-0001-8287-549X)

Adrián Alejandro-García

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico

Steven A. Calle-Schuler

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico

ORCID iD: [0000-0003-1304-585X](https://orcid.org/0000-0003-1304-585X)

Alexis Santana-Cruz

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico

Lucia Kmecova**Jonathan M. Blagburn**

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico

ORCID iD: [0000-0002-7681-2764](https://orcid.org/0000-0002-7681-2764)

Eric D. Hoopfer

Neuroscience Program, Carleton College, Northfield, Minnesota

ORCID iD: [0000-0002-5888-0173](https://orcid.org/0000-0002-5888-0173)

Andrew M. Seeds

Institute of Neurobiology, University of Puerto Rico-Medical Sciences Campus, San Juan, Puerto Rico

For correspondence: seeds.andrew@gmail.com

ORCID iD: [0000-0002-4932-6496](https://orcid.org/0000-0002-4932-6496)

Editors

Reviewing Editor

Sonia Sen

Tata Institute for Genetics and Society, India

Senior Editor

K VijayRaghavan

National Centre for Biological Sciences, Tata Institute of Fundamental Research, India

Reviewer #1 (Public Review):

Sensory neurons of the mechanosensory bristles on the head of the fly project to the subesophageal ganglion (SEZ). In this manuscript, the authors have built on a large body of previous work to comprehensively classify and quantify the head bristles. They broadly identify the nerves that various bristles use to project to the SEZ and describe their region-specific innervation in the SEZ. They use dye-fills, clonal labelling, and electron microscopic reconstructions to describe in detail the phenomenon of somatotopy - conserved peripheral representations within the central brain - within the innervation of these neurons. In the process they develop novel tools to access subsets of these neurons. They use these to demonstrate that groups of bristles in different parts of the head control different aspects of the grooming sequence.

Reviewer #2 (Public Review):

The authors combine genetic tools, dye fills and connectome analysis techniques to generate a "first-of-its-kind", near complete, synaptic resolution map of the head bristle neurons of

Drosophila. While some of the BMN anatomy was already known based on previous work by the authors and other researchers, this is the first time a near complete map has been created for the head BMNs at electron microscopy resolution.

Strengths:

1. The authors cleverly use techniques that allow moving back and forth between periphery (head bristle location) and brain, as well as moving between light microscopy and electron microscopy data. This allows them to first characterize the pathways taken by different head BMNs to project to the brain and also characterize anatomical differences among individual neurons at the level of morphology and connectivity.
2. The work is very comprehensive and results in a near complete map of all head BMNs.
3. Authors also complement this anatomical characterization with a first-level functional analysis using optogenetic activation of BMNs that results in expected directed grooming behavior.

Weaknesses:

1. The clustering analysis is compelling but cluster numbers seem to be arbitrarily chosen instead of by using some informed metrics.
2. It could help provide context if authors revealed some of the important downstream pathways that could explain optogenetics behavioral phenotypes and previously shown hierarchical organization of grooming sequences.
3. In contrast to the rigorous quantitative analysis of the anatomical data, the behavioral data is analyzed using much more subjective methods. While I do not think it is necessary to perform a rigorous analysis of behaviors in this anatomy focused manuscript, the conclusions based on behavioral analysis should be treated as speculative in the current form e.g. calling "nodding + backward walking" as an avoidance response is not justified as it currently stands. Strong optogenetic activation could lead to sudden postural changes that due to purely biomechanical constraints could lead to a couple of backward steps as seen in the example videos. Moreover since the quantification is manual, it is not clear what the analyst interprets as backward walking or nodding. Interpretation is also concerning because controls show backward walking (although in fewer instances based on subjective quantification).

Summary:

The authors end up generating a near-complete map of head BMNs that will serve as a long-standing resource to the *Drosophila* research community. This will directly shape future experiments aimed at modeling or functionally analyzing the head grooming circuit to understand how somatotopy guides behaviors.

Reviewer #3 (Public Review):

Eichler et al. set out to map the locations of the mechanosensory bristles on the fly head, examine the axonal morphology of the bristle mechanosensory neurons (BMNs) that innervate them, and match these to electron microscopy reconstructions of the same BMNs in a previously published EM volume of the female adult fly brain. They used BMN synaptic connectivity information to create clusters of BMNs that they show occupy different regions of the subesophageal zone brain region and use optogenetic activation of subsets of BMNs to support the claim that the morphological projections and connectivity of defined groups of BMNs are consistent with the parallel model for behavioral sequence generation.

The authors have beautifully cataloged the mechanosensory bristles and the projection paths and patterns of the corresponding BMN axons in the brain using detailed and painstaking methods. The result is a neuroanatomy resource that will be an important community resource. To match BMNs reconstructed in an electron microscopy volume of the

adult fly brain, the authors matched clustered reconstructed BMNs with light-level BMN classes using a variety of methods, but evidence for matching is only summarized and not demonstrated in a way that allows the reader to evaluate the strength of the evidence. The authors then switch from morphology-based categorization to non-BMN connectivity as a clustering method, which they claim demonstrates that BMNs form a somatotopic map in the brain. This map is not easily appreciated, and although contralateral projections in some populations are clear, the distinct projection zones that are mentioned by the authors are not readily apparent. Because of the extensive morphological overlap between connectivity-based clusters, it is not clear that small projection differences at the projection level are what determines the post-synaptic connectivity of a given BMN cluster or their functional role during behavior. The claim the somatotopic organization of BMN projections is preserved among their postsynaptic partners to form parallel sensory pathways is not supported by the result that different connectivity clusters still have high cosine similarity in a number of cases (i.e. Clusters 1 and 3, or Clusters 1 and 2). Finally, the authors use tools that were generated during the light-level characterization of BMN projections to show that specifically activating BMNs that innervate different areas of the head triggers different grooming behaviors. In one case, activation of a single population of sensory bristles (InOm) triggers two different behaviors, both eye and dorsal head grooming. This result does not seem consistent with the parallel model, which suggests that these behaviors should be mutually exclusive and rely on parallel downstream circuitry.

This work will have a positive impact on the field by contributing a complete accounting of the mechanosensory bristles of the fruit fly head, describing the brain projection patterns of the BMNs that innervate them, and linking them to BMN sensory projections in an electron microscopy volume of the adult fly brain. It will also have a positive impact on the field by providing genetic tools to help functionally subdivide the contributions of different BMN populations to circuit computations and behavior. This contribution will pave the way for further mechanistic study of central circuits that subserve grooming circuits.