

ROM1 is redundant to PRPH2 as a molecular building block of photoreceptor disc rims

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

August 24, 2023 (this version)

Posted to bioRxiv

July 2, 2023

Sent for peer review

June 13, 2023

Taylor R. Lewis, Mustafa S. Makia, Carson M. Castillo, Ying Hao, Muayyad R. Al-Ubaidi, Nikolai P. Skiba, Shannon M. Conley, Vadim Y. Arshavsky , Muna I. Naash 

Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710 • Department of Biomedical Engineering, University of Houston, Houston, TX, USA, 77204 • College of Optometry, University of Houston, Houston, TX, USA, 77204 • Department of Cell Biology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA, 73104 • Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC, USA, 27710

 https://en.wikipedia.org/wiki/Open_access

 <https://creativecommons.org/licenses/by/4.0/>

Abstract

Visual signal transduction takes place within a stack of flattened membranous “discs” enclosed within the light-sensitive photoreceptor outer segment. The highly curved rims of these discs, formed in the process of disc enclosure, are fortified by large hetero-oligomeric complexes of two homologous tetraspanin proteins, PRPH2 (a.k.a. peripherin-2 or rds) and ROM1. While mutations in PRPH2 affect the formation of disc rims, the role of ROM1 remains poorly understood. In this study, we found that the knockout of ROM1 causes a compensatory increase in the disc content of PRPH2. Despite this increase, discs of ROM1 knockout mice displayed a delay in disc enclosure associated with a large diameter and lack of incisures in mature discs. Strikingly, further increasing the level of PRPH2 rescued these morphological defects. We next showed that disc rims are still formed in a knockin mouse in which the tetraspanin body of PRPH2 was replaced with that of ROM1. Together, these results demonstrate that, despite its contribution to the formation of disc rims, ROM1 can be replaced by an excess of PRPH2 for timely enclosure of newly forming discs and establishing normal outer segment structure.

eLife assessment

This **valuable** study is focused on the requirement for the photoreceptor-specific tetraspanins, ROM1 and PRPH2, in the formation of the light-sensitive membrane discs. The evidence supporting the claim that deficiency in one of the proteins can be compensated by the other is **convincing**, with both established and modern techniques yielding results that will be of interest to those studying photoreceptor development and membrane curvature.

The light-sensitive outer segment organelle of vertebrate photoreceptor cells is a modified cilium containing a stack of disc-shaped membranes, or “discs”. While initially formed as serial evaginations of the photoreceptor plasma membrane, mature discs are separated from this membrane and are fully enclosed inside the outer segment (Steinberg et al., 1980 [DOI](#); Burgoyne et al., 2015 [DOI](#); Ding et al., 2015 [DOI](#); Volland et al., 2015 [DOI](#)) (reviewed in (Spencer et al., 2020 [DOI](#))). Membrane enclosure involves the formation of highly curved, hairpin-shaped disc rims. Two homologous tetraspanin proteins, PRPH2 (also known as peripherin-2 or rds) and ROM1, localize specifically to disc rims (Connell and Molday, 1990 [DOI](#); Travis et al., 1991 [DOI](#); Bascom et al., 1992 [DOI](#)) (reviewed in (Stuck et al., 2016 [DOI](#))). Within the disc rim, PRPH2 and ROM1 form homo- and heteromeric complexes (Bascom et al., 1992 [DOI](#); Goldberg et al., 1995 [DOI](#); Goldberg and Molday, 1996b [DOI](#); Goldberg and Molday, 1996a [DOI](#)) that are assembled into larger oligomers connected by disulfide bonds (Goldberg et al., 1998 [DOI](#); Loewen and Molday, 2000 [DOI](#); Chakraborty et al., 2009 [DOI](#); Zulliger et al., 2018 [DOI](#)). Three parallel interconnected chains of these oligomers are wrapped around the circumference of the mature disc and support its hairpin-shaped structure (Pöge et al., 2021 [DOI](#)).

Despite a high level of structural homology between PRPH2 and ROM1 (Li et al., 2003 [DOI](#)), the consequences of their mutations for photoreceptor health are quite different. Around 200 mutations of *PRPH2* are shown to cause a heterogeneous set of inherited retinal diseases, including retinitis pigmentosa, cone-rod dystrophy and macular dystrophies (Landrum et al., 2018 [DOI](#); Peeters et al., 2021 [DOI](#)). Yet, mutations in *ROM1* typically cause digenic retinitis pigmentosa in conjunction with mutations in *PRPH2* (Kajiwara et al., 1994 [DOI](#); Dryja et al., 1997 [DOI](#)), except for a handful of reports of mutations in *ROM1* in patients without accompanying *PRPH2* mutations (Bascom et al., 1995 [DOI](#); Sakuma et al., 1995 [DOI](#); Reig et al., 2000 [DOI](#)). With regard to animal models, the *Prph2* knockout mouse, commonly known as the *rds* mouse, has a complete failure of outer segment formation (Cohen, 1983 [DOI](#); Jansen and Sanyal, 1984 [DOI](#)), while the *Rom1* knockout mouse forms outer segments with relatively minor structural abnormalities (Clarke et al., 2000 [DOI](#)). This discrepancy could be explained, at least in part, by the importance of the cytoplasmic C-terminus of PRPH2, which has been shown to retain membranes at the photoreceptor cilium, thereby allowing their remodeling into outer segment discs (Salinas et al., 2017 [DOI](#)). However, most of the disease-associated PRPH2 mutations spare its C-terminus, so the complete explanation of this discrepancy awaits further investigation of the role of ROM1.

In this study, we investigated the role of ROM1 in the formation of photoreceptor disc rims and the process of disc enclosure. We first revisited the phenotype of the *Rom1* knockout (*Rom1*^{-/-}) mouse and made several novel observations. Using quantitative proteomics, we showed that the knockout of ROM1 causes a compensatory increase in the relative disc content of PRPH2. Despite the total tetraspanin content being essentially the same as in WT discs, *Rom1*^{-/-} discs displayed a delay in their maturation and an increase in outer segment diameter. In addition, *Rom1*^{-/-} discs lacked the indentations of their rims, known as incisures, present in WT discs. These changes could be explained, at least in part, by an alteration in tetraspanin oligomerization likely arising from a rearrangement of intramolecular disulfide bridge(s) in PRPH2. Strikingly, these morphological phenotypes can be rescued by transgenic overexpression of PRPH2, suggesting that ROM1 is not absolutely necessary for the formation of normal outer segments. In another set of experiments, we investigated whether disc rims could be formed in a knockin mouse in which the tetraspanin core of PRPH2 was replaced with that of ROM1. While the knockin photoreceptors failed to form orderly disc stacks, outer segment membranes preserved the ability to form hairpin-shaped rims. Together, these results demonstrate that, despite its contribution to disc formation and intrinsic ability to support disc rim structure, ROM1 can be replaced by an excess of PRPH2 for the formation of a normal outer segment.

Loss of ROM1 causes a compensatory increase in the relative outer segment content of PRPH2

Three proteins, rhodopsin, PRPH2 and ROM1, comprise ~98% of the total transmembrane protein material in normal discs (Skiba et al., 2023). Recently, it has been proposed that the overall dimensions of a disc are determined by the molar ratio between the total tetraspanin content (PRPH2 and ROM1) forming the disc rim and rhodopsin forming the disc surface (Lewis et al., 2023). Therefore, interpreting the phenotype of the *Rom1*^{-/-} mouse requires an understanding of how the loss of ROM1 affects the outer segment content of PRPH2. This is particularly important because discs deficient in PRPH2 display a compensatory increase in the relative amount of ROM1 (Lewis et al., 2023), suggesting that the opposite (i.e., a compensatory increase in the amount of PRPH2 when there is a deficiency in ROM1) may also be true.

To address this possibility, we employed a quantitative mass spectrometry approach (Skiba et al., 2023) recently used to determine the molar ratio among PRPH2, ROM1 and rhodopsin in other mutant photoreceptors (Lewis et al., 2023). In these experiments, the outer segment content of PRPH2 was determined as a molar fraction of rhodopsin. Because the packing density of rhodopsin in discs is unaffected by any changes in disc dimensions (Liang et al., 2004), this parameter reflects the relative content of PRPH2 in each disc, independent of any changes in outer segment length or volume. We found that the relative content of PRPH2 in *Rom1*^{-/-} outer segments increased to an ~1:12 molar ratio to rhodopsin from an ~1:18 ratio in WT outer segments (Table 1; see raw data in Supplementary Table 1). Strikingly, this 1:12 ratio is equal to the ratio between total tetraspanin (PRPH2 + ROM1) and rhodopsin in WT outer segments. Therefore, the knockout of ROM1 leads to a compensatory increase in the relative disc content of PRPH2.

Loss of ROM1 delays disc enclosure causing increased outer segment diameter and occasional disc overgrowth

The retinal phenotype of *Rom1*^{-/-} mice was analyzed at postnatal day 30 (P30), as photoreceptor cell degeneration was just beginning at this age (Figs. 1A,B). Interestingly, light microscopy imaging was sufficient to identify that the outer segment layer of these mice appeared disorganized and even shortened (Figs. 1A,C). Please note that this and the three subsequent figures also include data obtained with a mouse overexpressing PRPH2 on the *Rom1*^{-/-} background (PRPH2 OE/*Rom1*^{-/-}), which will be described below, to facilitate a side-by-side comparison of all three phenotypes. Using transmission electron microscopy (TEM), we observed a number of outer segment structural abnormalities (Fig. 2 and Supplementary Fig. 1), consistent with those noted in the original report (Clarke et al., 2000). In general, outer segments appeared to be shorter and wider, with some of them displaying overgrown disc membranes not aligned in a stack (yellow arrows, Fig 2B). The degree of this overgrowth varied, with the extended membranes occasionally wrapping around the entire outer segment.

Because these ultrastructural defects are reminiscent of those from several PRPH2 mutant lines in which photoreceptor disc enclosure is affected (Lewis et al., 2021), we assessed the status of disc enclosure in *Rom1*^{-/-} outer segments. We contrasted retinal tissue with tannic acid and uranyl acetate instead of the traditionally used osmium tetroxide (Ding et al., 2015). This technique yields a darker staining of newly forming, “open” discs exposed to the extracellular space than discs fully enclosed within the outer segment. This approach revealed that even outer segments lacking gross abnormalities in *Rom1*^{-/-} mice have an increased number of open discs at their base (Figs. 3A,B). Whereas WT rods contained a relatively constant number of ~7 open discs (consistent with previous reports (Ding et al., 2015; Volland et al., 2015; Lewis et al., 2021)), *Rom1*^{-/-} rods had an average of ~14 open discs. In addition, the number of open discs in *Rom1*^{-/-}

Protein molar ratios*	WT	<i>Rom1</i> ^{-/-}
PRPH2:rhodopsin	1:18.1 ± 0.5	1:12.2 ± 1.3
ROM1:rhodopsin	1:36.3 ± 3.9	n/a
PRPH2:ROM1	2.0:1 ± 0.3	n/a
(PRPH2 + ROM1):rhodopsin	1:12.1 ± 0.2	1:12.2 ± 1.3

*Values are shown as mean ± s.d. n/a: not applicable. Two outer segment preparations from mice of each genotype were analyzed.

Table 1.

Quantification of molar ratios between tetraspanins and rhodopsin in WT and *Rom1*^{-/-} mice




Figure 1.

Light microscopy of retinas from WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice

(A) Representative light microscopy images of WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} retinas analyzed at P30. OS: outer segment; ONL: outer nuclear layer. Scale bar: 20 μm . **(B)** Quantification of the number of photoreceptor nuclei in a 100 μm segment of the retina at 500 μm increments away from the optic nerve (ON). Three retinas were analyzed for each genotype. Two-way ANOVA revealed statistically significant differences in the nuclear counts across genotypes ($p = 0.0002$). Sidak's multiple comparisons post-hoc test revealed statistically significant differences between the total nuclear count between WT and *Rom1*^{-/-} retinas ($p = 0.0001$) and between *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} retinas ($p = 0.0148$), but not between WT and *PRPH2 OE/Rom1*^{-/-} retinas ($p = 0.3564$). **(C)** Quantification of the outer segment (OS) layer length at 500 μm increments away from the optic nerve. Three retinas were analyzed for each genotype. Two-way ANOVA revealed statistically significant differences in the OS layer lengths across genotypes ($p < 0.0001$). Sidak's multiple comparisons post-hoc test revealed statistically significant differences for the total nuclear count between WT and *Rom1*^{-/-} retinas ($p < 0.0001$) and between WT and *PRPH2 OE/Rom1*^{-/-} retinas ($p = 0.0006$), but not between *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} retinas ($p = 0.2924$).




Figure 2.

Ultrastructural analysis of retinas and rod outer segments from WT, *Rom1*^{-/-} and *PRPH2* OE/*Rom1*^{-/-} mice

(A) Representative low magnification TEM images of WT, *Rom1*^{-/-} and *PRPH2* OE/*Rom1*^{-/-} retinas analyzed at P30. RPE: retinal pigment epithelium; OS: outer segment; IS: inner segment. Scale bar: 10 μ m. **(B)** Representative high magnification TEM images of *Rom1*^{-/-} outer segments. Yellow arrows indicate outer segment structural defects that range from slightly overgrown open discs to membranous whorls. Scale bar: 1 μ m.

rods was much more variable than in WT rods, with up to ~30 open discs occasionally observed. Lastly, we quantified outer segment diameters of *Rom1*^{-/-} rods and found that they were on average ~35% wider than WT rods (Fig. 3C), which is close to the ~43% increase previously reported in (Clarke et al., 2000).

These findings show that ROM1 contributes to the process of disc enclosure and that the compensatory increase in the disc content of PRPH2 does not fully replace ROM1 in this capacity. Accordingly, the increase in outer segment diameter may be explained by prolonged delivery of disc membrane material to each disc before it becomes fully enclosed. On occasion, an expansion of open discs becomes uncontrolled leading to formation of membranous whorls.

Another phenotype of *Rom1*^{-/-} mice was revealed using TEM of tangentially sectioned retinas (Fig. 4). Normal mouse discs contain a single deep indentation of their rims, called an incisure (yellow arrowheads, Fig. 4). In contrast, no incisures were observed in *Rom1*^{-/-} discs. Because the rims of both disc circumference and incisure are built from PRPH2/ROM1, the lack of incisures in *Rom1*^{-/-} discs may be explained by the entire pool of remaining tetraspanin being deposited at the circumference of larger discs.

ROM1 regulates the supramolecular organization of PRPH2

We next sought to address why the compensatory increase in the disc content of PRPH2 in *Rom1*^{-/-} rods is insufficient to support normal disc morphogenesis. In this context, it is important to consider that defects in PRPH2 oligomerization, without an accompanying decrease in PRPH2 levels, also lead to defects in disc enclosure (Lewis et al., 2021). Therefore, we reasoned that loss of ROM1 may modulate the status of PRPH2 oligomerization. In fact, the idea that ROM1 may regulate the formation of high order PRPH2 oligomers has been put forward in a previous study (Loewen and Molday, 2000).

To address the possibility that PRPH2 oligomerization may be disrupted without ROM1, we first performed SDS-PAGE of retinal lysates from WT and *Rom1*^{-/-} mice under non-reducing conditions (i.e., no DTT in samples). Under these conditions, PRPH2 is represented by two bands, one corresponding to ~35 kDa monomers and another to ~75 kDa dimers linked by a disulfide bond (Fig. 5A). Both bands are somewhat diffuse, likely reflecting the heterogeneous status of PRPH2 glycosylation. The majority of PRPH2 in WT retinas is represented by dimers, consistent with its origin from large, disulfide-bound oligomers.

Two differences were observed in *Rom1*^{-/-} retinas. First, there was an ~50% increase in the fraction of PRPH2 that runs in a monomeric state (Figs. 5A,B), indicating that PRPH2 oligomerization may be negatively affected by loss of ROM1. Second, the monomeric form of PRPH2 was represented by a doublet, consistent with a previous observation (Stuck et al., 2015). This doublet was consolidated into a single band under reducing conditions in the presence of DTT (Fig. 5C), which suggests that the two bands in *Rom1*^{-/-} retinas represent pools of PRPH2 molecules with different patterns of internal disulfide bonds. Apart from the C150 residue involved in intermolecular disulfide bonding of a PRPH2 molecule to another PRPH2 or ROM1, there are six other conserved cysteines in PRPH2 that are involved in intramolecular disulfide bonding (Goldberg et al., 1998). Therefore, it is possible that ROM1 contributes to PRPH2 assuming the intermolecular disulfide bond conformation optimal for tetraspanin oligomerization at the disc rim.

To further investigate the status of PRPH2 oligomerization in *Rom1*^{-/-} mice, we utilized a technique typically used to describe PRPH2 and ROM1 complexes of various sizes. In this approach, complexes are extracted from disc membranes in the presence of a non-ionic detergent and separated by velocity sedimentation on a sucrose gradient under non-reducing conditions (Clarke et al., 2000; Goldberg et al., 2001; Ding et al., 2004; Chakraborty et al., 2008; Chakraborty




Figure 3.

Analysis of disc enclosure in rod outer segments of WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice

(A) Representative high magnification TEM images of tannic acid/uranyl acetate-stained retinas of WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice analyzed at P30. This approach stains newly forming “open” discs more intensely than mature enclosed discs. Yellow arrows point to darkly stained, unenclosed discs; yellow arrowheads point to lightly stained, enclosed discs. Scale bar: 0.5 μ m. **(B)** Quantification of the number of darkly stained open discs at the rod outer segment base. Each data point represents a single outer segment. For each genotype, three retinas were analyzed with at least 35 outer segments analyzed per retina. Data were plotted with samples separated, while statistical analysis was performed on the averages within each retina (n=3 for each genotype). One-way ANOVA revealed statistically significant differences in the number of open discs across genotypes (p < 0.0001). Tukey’s multiple comparisons post-hoc test revealed statistically significant differences in the number of open discs between WT and *Rom1*^{-/-} (p < 0.0001) and *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} (p < 0.0001) mice, but not between WT and *PRPH2 OE/Rom1*^{-/-} mice (p = 0.2686). **(C)** Quantification of the outer segment (OS) diameter. For each genotype, three retinas were analyzed with at least 66 outer segments analyzed per retina. One-way ANOVA revealed statistically significant differences in the OS diameters across genotypes (p = 0.0074). Tukey’s multiple comparisons post-hoc test revealed statistically significant differences in the OS diameters between WT and *Rom1*^{-/-} (p = 0.0083) and *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} (p = 0.0197) mice, but not between WT and *PRPH2 OE/Rom1*^{-/-} mice (p = 0.7198). Note that these diameters were measured in longitudinal sections, in which outer segment are not always sectioned across their widest part; therefore, these values are likely under-representations of the true OS diameters. However, this does not affect the comparison across genotypes.




Figure 4.

Ultrastructural analysis of disc incisures in WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice

Representative TEM images of retinas, tangentially sectioned through the outer segment layer, from WT, *Rom1*^{-/-} and *PRPH2 OE/Rom1*^{-/-} mice analyzed at P30. Yellow arrowheads indicate incisures observed in WT and *PRPH2 OE/Rom1*^{-/-} discs, but not *Rom1*^{-/-} discs. Scale bar: 1 μ m.




Figure 5.

Loss of ROM1 alters the oligomerization status of PRPH2

(A) Western blot probed for PRPH2 after protein separation by SDS-PAGE under non-reducing conditions. Each sample contained 10 μ g of lysate obtained from eyecups of WT or *Rom1*^{-/-} mice at P30. Under these conditions, PRPH2 runs as disulfide-bound dimers (~75 kDa) and monomers (~37 kDa). The monomer band of PRPH2 runs as a doublet in *Rom1*^{-/-} but not WT eyecups. (B) Quantification of the ratio between monomer and disulfide-bound (dimer) bands of PRPH2 was performed using densitometry of three independent lysates. For the doublet of monomer bands in *Rom1*^{-/-} lysates, both bands were used for quantification. Unpaired t-test revealed a statistically significant difference in the PRPH2 monomer:dimer ratio between WT and *Rom1*^{-/-} retinas ($p = 0.0071$). (C) Western blot probed for PRPH2 after protein separation by SDS-PAGE under reducing conditions. Each sample contained 10 μ g of lysate obtained from eyecups of WT or *Rom1*^{-/-} mice. (D) Lysates obtained under non-reducing conditions from eyecups of WT and *Rom1*^{-/-} mice were subjected to velocity sedimentation on 5-20% sucrose gradients. Twelve fractions were collected with fraction #1 corresponding to 20% sucrose and fraction #12 to 5% sucrose. Proteins from each fraction were subjected to non-reducing SDS-PAGE and Western blotting for PRPH2. The distribution of molecular mass standards across fraction, as determined in (Chakraborty et al., 2008), is shown above the panels. (E) Quantification of both the monomeric and disulfide-bound bands of PRPH2 in each fraction was performed using densitometry of at least four independent samples and normalized to the total PRPH2 content across all fractions. Two-way ANOVA revealed statistically significant differences in the PRPH2 content across genotypes and fractions ($p < 0.0001$ for disulfide-bound, $p = 0.9489$ for monomer). Sidak's multiple comparisons post-hoc test revealed statistically significant differences between genotypes for the disulfide-bound form in fractions #1 ($p = 0.0022$), #4 ($p = 0.0002$) and #5 ($p = 0.0061$).

et al., 2009 [↗](#); Stuck et al., 2014 [↗](#); Zulliger et al., 2018 [↗](#); Milstein et al., 2020 [↗](#); Lewis et al., 2021 [↗](#)). We performed velocity sedimentation of lysates from both WT and *Rom1*^{-/-} retinas followed by SDS-PAGE of gradient fractions under non-reducing conditions (**Figs. 5D,E** [↗](#)). The distribution of the PRPH2 monomeric form across fractions was unaffected by the ROM1 knockout. In both WT and *Rom1*^{-/-} retinas, it was most abundant in fractions #6-8, shown to correspond to PRPH2 core complexes (Chakraborty et al., 2008 [↗](#)). However, the loss of ROM1 shifted the distribution of the disulfide-bound PRPH2 dimers to smaller oligomeric forms, consistent with a previous report (Conley et al., 2019 [↗](#)). Taken together, these data show that loss of ROM1 does indeed affect the status of PRPH2 oligomerization and suggest that abnormal PRPH2 oligomerization underlies the defects in disc enclosure in *Rom1*^{-/-} mice.

Transgenic overexpression of PRPH2 can compensate for the loss of ROM1

We next investigated whether a further increase in the level of PRPH2 could improve the morphological defects observed in *Rom1*^{-/-} outer segments. We employed a transgenic line that expresses PRPH2 in both rods and cones under control of the human IRBP promoter (Nour et al., 2004 [↗](#)). While originally called NMP, we refer to this line as *PRPH2 OE* (PRPH2 overexpressor). In the current study, we used *PRPH2 OE* mice with a single copy of this transgene, which has been shown to express ~30% excess PRPH2 over WT levels (Nour et al., 2004 [↗](#)). *PRPH2 OE* mice were crossed with *Rom1*^{-/-} mice and retinas were analyzed as for WT and *Rom1*^{-/-} mice (**Figs. 1** [↗](#)**3** [↗](#)). Strikingly, the gross morphological defects observed in *Rom1*^{-/-} mice were significantly improved by transgenic overexpression of PRPH2 (**Fig. 2A** [↗](#)). The outer segment length was somewhat restored (**Figs. 1A,C** [↗](#)), although the degree of this effect did not reach statistical significance. Furthermore, PRPH2 overexpression rescued the disc enclosure defect of *Rom1*^{-/-} outer segments by restoring the number of open discs to the WT level (**Figs. 3A,B** [↗](#)). In addition, *PRPH2 OE/Rom1*^{-/-} outer segments had normal diameters (**Figs. 3A,C** [↗](#)) and their discs contained incisures (**Fig. 4** [↗](#)). These data indicate that, whereas ROM1 contributes to disc formation in WT rods, it can be functionally replaced by a sufficient excess of PRPH2.

ROM1 is able to form disc rims in the absence of PRPH2

While we have shown that ROM1 contributes to the formation of disc rims, it is unclear whether it can do so in the absence of PRPH2. Because disc formation is completely abolished in the absence of PRPH2, we could not address this question using PRPH2 knockout mice. Instead, we utilized a knockin mouse model, called RRCT (Conley et al., 2019 [↗](#)), in which the tetraspanin core of PRPH2 was replaced with that of ROM1 (**Fig. 6A** [↗](#)). This chimeric protein contains the C-terminus of PRPH2 required for retaining disc membranes at the outer segment base (Salinas et al., 2017 [↗](#)), while replacing the tetraspanin core required for its oligomerization and functioning in disc rim formation (Salinas et al., 2017 [↗](#); Conley et al., 2019 [↗](#)).

As previously reported (Conley et al., 2019 [↗](#)), homozygous RRCT photoreceptors formed rudimentary outer segments (**Fig. 6B** [↗](#)). The small number and size of these outer segment-like structures is likely explained by a very low expression of the chimeric protein (Conley et al., 2019 [↗](#)), which may lead to a low efficiency of disc membrane retention. We now show that some of these structures contain stacked disc membranes that are partially enclosed within the outer segment plasma membrane. Importantly, these disc membranes have discernible rims with a characteristic hairpin-like shape. This finding indicates that ROM1 alone is, in principle, able to support the formation of disc rims in the absence of PRPH2.

In an additional set of experiments, we explored the oligomerization status of ROM1 and the RRCT chimera in the absence of PRPH2. Unfortunately, the severe disruptions of outer segments in homozygous RRCT mice precluded us from obtaining a sufficient amount of retinal lysate to perform this analysis. Instead, we used an *in vitro* system in which we co-transfected equal amounts of RRCT and ROM1 in COS7 cells (**Fig. 6C** [↗](#)). In these cells, RRCT and ROM1 co-localized




Figure 6.

The tetraspanin core of ROM1 is sufficient for disc rim formation

(A) Cartoon schematic of the RRCT chimeric tetraspanin protein. In the RRCT mouse, the *Prph2* gene has a knockin mutation that replaces it with a DNA sequence encoding a chimeric protein consisting of the tetraspanin body of ROM1 while retaining the C-terminal tail of PRPH2 that is essential for disc formation. **(B)** Representative TEM images of homozygous RRCT mice analyzed at P30. The boxed inset (left) is shown at a higher magnification (right) that reveals the presence of disc rims. Scale bars: 1 μ m (left); 0.1 μ m (right). **(C)** Immunofluorescent images of COS7 cells co-transfected with RRCT-FLAG and ROM1 constructs. Cells were stained with antibodies against RRCT (red), ROM1 (blue) and calreticulin (green) to label ER membranes. Nuclei were counterstained with DAPI (grey). Scale bar: 10 μ m. **(D)** Lysates obtained under non-reducing conditions from COS7 cells co-transfected with RRCT-FLAG and ROM1 constructs were subjected to velocity sedimentation on 5-20% sucrose gradients. Twelve fractions were collected with fraction #1 corresponding to 20% sucrose and fraction #12 to 5% sucrose. Proteins from each fraction were subjected to reducing SDS-PAGE and Western blotting for ROM1 and RRCT. **(E)** Quantification of ROM1 and RRCT in each fraction was performed using densitometry of three independent lysates and normalized to the total content across all fractions.

in structures that were distinct from the endoplasmic reticulum, as evident from the lack of co-localization with the endoplasmic reticulum marker, calreticulin, suggesting that they were normally processed in the biosynthetic membranes. Velocity sedimentation of lysates from these cells revealed that both RRCT and ROM1 sediment in fractions #6-8 (**Figs. 6D-E**), which is the same as for PRPH2/ROM1 core complexes from WT photoreceptors. These data indicate that heterologously expressed RRCT and ROM1 can form core tetraspanin complexes, even in the absence of PRPH2, that are able to support disc rim formation.

Disc rims can be formed without disulfide bonds between tetraspanin molecules

In a final experiment, we investigated whether the formation of intermolecular disulfide bonds between tetraspanins is dispensable for disc rim formation. We have previously shown that disulfide bonds between PRPH2 molecules are not absolutely required for this function because disc rims were formed in *Prph2*^{C150S/C150S} knockin mice lacking the cysteine residue forming this bond (Lewis et al., 2020). Yet ROM1 also forms intermolecular disulfide bonds (Chakraborty et al., 2008), which raises the question of whether disc rim formation in *Prph2*^{C150S/C150S} knockin mice was driven by disulfide-linked ROM1 molecules.

To address this question, we analyzed disc rim structure in *Rom1*^{-/-} mice bearing the *Prph2*^{C150S/C150S} mutation. Outer segments of these mice had morphological defects comparable to those of *Prph2*^{C150S/C150S} mice (**Fig. 7A**). Nonetheless, orderly disc stacks were occasionally observed in these retinas. Higher magnification images of these stacks revealed the presence of disc rims which looked comparable to those in *Prph2*^{C150S/C150S} retinas (**Fig. 7B**). This finding demonstrates that, whereas disulfide bonds connecting tetraspanin molecules are essential for maintaining the overall outer segment structure, these bonds are not absolutely required for the formation of disc rims.

In addition, staining with tannic acid/uranyl acetate revealed that the outer segments of *Prph2*^{C150S/C150S}/*Rom1*^{-/-} mice contained both open (yellow arrows, **Fig. 7A**) and enclosed discs (yellow arrowheads, **Fig. 7A**). This indicates that disulfide bonds among PRPH2 and ROM1 molecules are also dispensable for disc enclosure.

Discussion

In this study, we analyzed the role of ROM1 in supporting the formation of outer segment discs. In addition to the previously described morphological defects of *Rom1*^{-/-} outer segments, we showed that these outer segments experience a delay in disc enclosure. This delay leads to an increase in disc diameter, loss of incisures and occasional uncontrolled disc membrane outgrowth. Remarkably, these defects can be reversed by sufficient overexpression of PRPH2, indicating a functional redundancy between these two tetraspanin proteins.

The idea of redundancy between PRPH2 and ROM1 is supported by the finding that mutations in both genetically interact to modulate disease (Kajiwara et al., 1994; Dryja et al., 1997). While individual PRPH2 mutations have greater phenotypic consequences, this could relate to both the unique role of the C-terminal tail of PRPH2 in the initial stages of disc formation (Salinas et al., 2017) as well as the fact that there is twice as much PRPH2 as ROM1 in outer segments, at least in mice (Kedzierski et al., 1999; Loewen and Molday, 2000; Skiba et al., 2023).

While evidence that *ROM1* mutations alone may cause human visual pathology is limited (Bascom et al., 1995; Sakuma et al., 1995; Reig et al., 2000), photoreceptors of *Rom1*^{-/-} mice eventually degenerate (Clarke et al., 2000). We propose that the delay in disc enclosure is the primary defect underlying this degeneration. Because discs remain open longer than normal, they




Figure 7.

Disc rims can be formed in the absence of intermolecular disulfide bonds between PRPH2 and ROM1

(A) Representative TEM images of tannic acid/uranyl acetate-stained retinas of *Prph2*^{C150S/C150S} and compound *Prph2*^{C150S/C150S}/*Rom1*^{-/-} mice analyzed at P30. Photoreceptor outer segments of each mouse have severely perturbed outer segment structure. Yet, disc enclosure is not completely prevented as there are both darkly-stained, “open” discs (yellow arrows) and lightly-stained, “enclosed” discs (yellow arrowheads) in each genotype. The boxed region of each image depicts an area in which disc stacking appears relatively normal and permits the analysis of disc rim structure. Scale bar: 1 μ m. **(B)** Higher magnification images of the regions in which disc stacking appears normal in both *Prph2*^{C150S/C150S} and *Prph2*^{C150S/C150S}/*Rom1*^{-/-} mice. Disc rims are formed in each genotype. Scale bar: 0.1 μ m.

may incorporate more than a normal amount of membranous material, which is continuously delivered to the outer segment. This can lead to disc membrane overgrowth and, eventually, the formation of highly dysmorphic membrane whorls. This progressive collapse of outer segment ultrastructure would eventually lead to cell death. Notably, this same progression of structural defects occurs in certain mice in which PRPH2 oligomerization is affected (Lewis et al., 2021 [DOI](#)).

But why is the phenotype of the heterozygous *rd*s mouse containing ~50% of normal PRPH2 protein (Hawkins et al., 1985 [DOI](#); Cheng et al., 1997 [DOI](#)) much more severe than the *Rom1*^{-/-} mouse lacking 100% of ROM1? This difference is not intuitive because the molar ratio between PRPH2 and ROM1 is ~2:1 and, therefore, each mouse is expected to have ~2/3 of the normal tetraspanin content. However, there is a compensatory increase in outer segment tetraspanin in each of these two mice. We now show that relative content of total tetraspanin in *Rom1*^{-/-} discs is essentially the same as in WT discs (~1:12 molar ratio to rhodopsin) due to a compensatory increase in PRPH2. On the other hand, there is only a partial compensatory increase of ROM1 in heterozygous *rd*s mice (~1:15 molar ratio of PRPH2+ROM1 to rhodopsin) (Lewis et al., 2023 [DOI](#)). The mechanism behind these increases remains to be elucidated, but regardless, the lesser severity of the *Rom1*^{-/-} phenotype could be explained, at least in part, by a larger degree of this compensatory tetraspanin increase.

Yet, the complete restoration of the total tetraspanin content in *Rom1*^{-/-} outer segments does not fully rescue their morphological phenotype. This implies that, in normal photoreceptors, ROM1 contributes some unique functional features to the properties of tetraspanin oligomers. In fact, our data suggest that the loss of ROM1 alters both the intra- and intermolecular properties of the remaining PRPH2. We found that ROM1 likely facilitates an optimal internal disulfide bond arrangement of PRPH2 in addition to promoting the formation of larger oligomeric chains of PRPH2. One possible explanation for the ability of ROM1 to potentially facilitate the optimal disulfide bond arrangement of PRPH2 may relate to the intracellular trafficking of these two proteins. Whereas the majority of PRPH2 is normally trafficked through the unconventional secretory pathway from the endoplasmic reticulum that bypasses the Golgi (Tian et al., 2014 [DOI](#); Zulliger et al., 2015 [DOI](#); Pearing et al., 2021 [DOI](#)), ROM1 was shown to direct a small fraction of PRPH2 through the conventional secretory pathway (Conley et al., 2019 [DOI](#)). It is intriguing to speculate that a shift in the intramolecular disulfide bonds of PRPH2 in the absence of ROM1 may arise from a complete loss of PRPH2 trafficking through the conventional pathway. Nonetheless, the outer segment deficiencies arising from loss of ROM1 were nearly completely overcome by a sufficient excess of PRPH2.

Of particular interest is our finding that discs of *Rom1*^{-/-} mice lack incisures. A recent study put forth a model in which the total length of the disc rim, including the incisure, is determined by the molar ratio between the total tetraspanin protein and rhodopsin (Lewis et al., 2023 [DOI](#)). The lack of incisures in *Rom1*^{-/-} discs may not appear intuitive since the total tetraspanin content in these discs is essentially the same as in WT discs. However, *Rom1*^{-/-} discs also have an increased diameter, which allows the total rim length of WT and *Rom1*^{-/-} discs to be about the same. The reason why such a tradeoff between the disc diameter and the presence/absence of an incisure exists remains to be determined, although it is likely to relate to the kinetics of disc enclosure.

In conclusion, our study shows that ROM1 makes distinct contributions to the formation of disc rims and the process of disc enclosure. Nonetheless, a sufficient excess of PRPH2 is able to compensate for the loss of ROM1, indicating that ROM1 is redundant to PRPH2 as a molecular building block of photoreceptor disc rims. Given that excess PRPH2 can rescue defects associated with loss of ROM1, it is exciting to speculate that PRPH2 overexpression can prevent the long-term photoreceptor degeneration associated with either loss of ROM1 or deficiencies arising from mutations in PRPH2 itself. It also remains to be seen whether the opposite is also true, whereby an excess ROM1 may rescue certain defects associated with mutations in PRPH2.

Mouse (*Mus musculus*) husbandry

Animal maintenance and experiments were approved by the local Institutional Animal Care and Use Committee (IACUC; University of Houston, TX, USA) and guidelines as stated by the Association for Research in Vision and Ophthalmology (Rockville, MD). The generation of the *Prph2*^{C150S} mouse was previously described in (Stuck et al., 2014 [DOI](#)). The generation of the RRCT mouse was previously described in (Conley et al., 2019 [DOI](#)). The generation of the *PRPH2* OE mouse was previously described in (Nour et al., 2004 [DOI](#)). The *Rom1*^{-/-} mouse, described in (Clarke et al., 2000 [DOI](#)), was generously provided by Roderick R McInnes (McGill University). All mice were genotyped to ensure that they did not contain either the *rd8* (Mattapallil et al., 2012 [DOI](#)) or *rd1* (Pittler et al., 1993 [DOI](#)) mutations commonly found in inbred mouse strains. All mice were on a C57BL/6 genetic background bearing the RPE65 L450 variant and housed under a 12/12 hour diurnal light (~30 lux) cycle.

Transmission electron microscopy (TEM)

For all main figures, fixation and processing of mouse eyes for TEM was performed as described previously (Ding et al., 2015 [DOI](#)). In the afternoon, anesthetized mice were transcardially perfused with 2% paraformaldehyde, 2% glutaraldehyde and 0.05% calcium chloride in 50 mM MOPS (pH 7.4) resulting in exsanguination. Eucleated eyes were fixed for an additional 2 h in the same fixation solution at room temperature. Eyecups were dissected from fixed eyes, embedded in 2.5% low-melt agarose (Precisionary, Greenville, NC) and cut into 200 µm thick slices on a Vibratome (VT1200S; Leica, Buffalo Grove, IL). Agarose sections were stained with 1% tannic acid (Electron Microscopy Sciences, Hatfield, PA) and 1% uranyl acetate (Electron Microscopy Sciences), gradually dehydrated with ethanol and infiltrated and embedded in Spurr's resin (Electron Microscopy Sciences). 70 nm sections were cut, placed on copper grids and counterstained with 2% uranyl acetate and 3.5% lead citrate (19314; Ted Pella, Redding, CA). The samples were imaged on a JEM-1400 electron microscope (JEOL, Peabody, MA) at 60 kV with a digital camera (BioSprint; AMT, Woburn, MA). Image analysis and processing was performed with ImageJ. For **Supplementary Figure 1** [DOI](#), fixation, processing, and imaging of mouse eyes for TEM was performed as described previously (Stricker et al., 2005 [DOI](#)).

Light microscopy of histological sections

Plastic embedded blocks generated for TEM were sectioned through the optic nerve in 500 nm sections and stained with methylene blue for light microscopy as previously described (Lobanova et al., 2008 [DOI](#)). Images were taken with a confocal microscope (Eclipse 90i and A1 confocal scanner; Nikon) with a 60× objective (1.4 NA Plan Apochromat VC; Nikon) using Nikon NIS-Elements software. Image analysis and processing was performed with ImageJ.

Preparation of retinal lysates

Eyecups were dissected and immediately frozen using liquid nitrogen and stored at -80°C prior to processing. Lysates were prepared essentially as previously described (Stuck et al., 2014 [DOI](#)). In short, individual eyecups were lysed in 200 µl solubilization buffer (PBS, pH 7.0, containing 1% Triton X-100, 5 mM EDTA, 5 mg/ml NEM and protease inhibitors (Roche, Mannheim, Germany)). Samples were incubated on ice for 1 h prior to being centrifuged for 30 min at 20,000g at 4°C. The supernatant was collected and subjected to either Western blot analysis directly or velocity sedimentation.

Western blotting of retinal lysates

Protein concentration was assayed by using a colorimetric Bradford assay (Bio-Rad, Hercules, CA, USA). Western blotting was performed essentially as previously described (Spencer et al., 2016). In short, lysates were incubated with Laemmli sample buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol and 1% Bromophenol Blue) with or without 100 mM DTT for reducing or non-reducing blots respectively. Samples containing 10 µg of total protein were incubated at 90°C for 5 minutes and run on a 10-20% Tris-HCl gel. Gels were transferred onto PVDF membrane and blocked with Intercept (PBS) Blocking Buffer (Li-Cor, Lincoln, Nebraska) with 0.25% Tween-20. Blots were incubated overnight at 4°C with 1:1000 dilution of polyclonal rabbit anti-PRPH2 C-terminal antibody (Kedzierski et al., 1999). After primary antibody incubation, blots were washed and incubated with 1:10000 dilution of donkey anti-rabbit DyLight 800 (Invitrogen, Carlsbad, CA) for 2 h. All experiments were repeated at least three times. Blots were imaged using Odyssey CLx imaging system (Li-Cor).

Velocity sedimentation using sucrose gradients

Sucrose gradients were prepared as previously described (Chakraborty et al., 2009). In short, gradients of 5–20% sucrose were prepared by sequentially layering 0.5 ml each of 20, 15, 10 and 5% sucrose solutions in PBS with 0.1% Triton X-100 and 10 mM NEM and allowing them to sit at room temperature for 1 h to equilibrate. Gradients were then chilled on ice for 30 minutes prior to loading lysate and centrifugation at 40,000 rpm on a TLS-55 swinging bucket rotor (Beckman Coulter, Brea, CA) for 16 hours at 4°C. The bottom of each tube was pierced with a 21G needle and 12 fractions of ~180 µl each were collected. Western blotting was performed as described above using 15 µl samples for each fraction.

Cell culture experiments

COS7 cells were transfected with 7.5 µg of RRCT-FLAG and ROM1 mammalian expression constructs that were previously described (Conley et al., 2010; Conley et al., 2019) using a calcium phosphate transfection protocol as previously described (Conley et al., 2019). After 48 hours, cells were processed for immunofluorescent staining or velocity sedimentation as described below. Experiments were repeated three independent times.

Immunofluorescent staining was performed as described in (Conley et al., 2010). After blocking, coverslips were incubated overnight in primary antibodies to label RRCT, ROM1 and ER membranes as follows: 1) 1:2 dilution of monoclonal mouse anti-RRCT (2E7; described in (Zulliger et al., 2015)), 2) 1:400 dilution of polyclonal rabbit anti-ROM1 (2Rom1; described in (Ding et al., 2005)), and 3) 1:250 dilution of polyclonal chicken anti-calreticulin at 1:250 (ab2908; Abcam, Cambridge, UK). After washing, cells were incubated with 1:000 dilution of donkey anti-mouse Alexa 568 (Invitrogen), donkey anti-rabbit Alexa 647 (Invitrogen) and goat anti-chicken Alexa 488 (Invitrogen) secondary antibodies for 1 h. Cells were mounted in ProlongGold with DAPI (ThermoFisher). Images were captured on a BX-62 spinning disk confocal microscope equipped with an ORCA-ER camera (Olympus, Japan) and analyzed with Slidebook™ 5.2 software (Intelligent Imaging Innovations, Denver, CO). Images from the spinning disk confocal microscope were deconvolved using the nearest neighbors paradigm. Images were captured with 100x/1.40 oil objectives, and exposure times and display settings (brightness and contrast) for all images were normalized to a control section where primary antibody was omitted during processing. No gamma adjustments were made to immunofluorescent images.

For velocity sedimentation, scraped cells were lysed in the same solubilization buffer used to prepare retinal extracts. Lysates were separated as described above (200 µg protein lysate/gradient). Resulting gradient fractions were separated by SDS-PAGE under reducing conditions and blots were probed with either 1:500 dilution of monoclonal mouse anti-RRCT (2B7; described in (Conley et al., 2014)) or monoclonal mouse anti-ROM1 (2H5, described in (Conley et

al., 2014 [\[4\]](#)). Blots were probed with goat anti-mouse and anti-rabbit HRP secondaries (SeraCare, Milford, MA), imaged on a Bio-Rad ChemiDoc imager and analyzed in Image Lab Software version 6.0.1 (Bio-Rad).

Quantitative mass spectrometry

A crude preparation of rod outer segments was obtained as described in (Lewis et al., 2023 [\[4\]](#)). Dissected mouse retinas were vortexed in 8% OptiPrep in mouse Ringer's solution (containing 130 mM NaCl, 3.6 mM KCl, 2.4 mM MgCl₂, 1.2 mM CaCl₂, and 10 mM HEPES, pH 7.4) that was adjusted to 314 mOsm. The preparation was briefly left on ice to allow the remaining retinal tissue to settle. The supernatant was removed and centrifuged at 20,000 x g. Pelleted outer segments were gently washed with mouse Ringer's solution before lysis with 2% SDS in PBS. Protein concentration was measured with the Bio-Rad Protein Assay kit (Bio-Rad). Samples containing 5–10 µg of protein were mixed with 0.25–0.5 µg BSA (used as an internal standard in this analysis) and cleaved with 1 µg trypsin/LysC mix (Promega, Madison, WI) using the SP3 beads protocol described in (Hughes et al., 2014 [\[4\]](#)). The combined digest of outer segments and BSA was mixed with the digest of a chimeric protein consisting of concatenated tryptic peptides of outer segment proteins, including rhodopsin, PRPH2 and ROM1, which is described in (Skiba et al., 2023 [\[4\]](#)). Mass spectrometry, data processing and data analysis were also performed as described in (Skiba et al., 2023 [\[4\]](#)). For each genotype, a total of two biological replicates were analyzed.

Experimental design and statistical analysis

For the quantification of the number of photoreceptor nuclei (**Fig. 1B** [\[4\]](#)), photoreceptor nuclei were counted in 100 µm boxes at 500 µm intervals from the optic nerve spanning 2000 µm in each direction for three mice of each genotype, as previously described (Lobanova et al., 2018 [\[4\]](#)). Two-way ANOVA was performed to determine statistical significance across genotype and location.

For the quantification of the number of open discs (**Fig. 3B** [\[4\]](#)), darkly-stained new discs at the base of the rod outer segment were counted until the first lightly-stained enclosed disc. 3 retinas were collected of each genotype with the number of outer segments analyzed as follows: WT1, 44; WT2, 40; WT3, 68; *Rom1*^{-/-}1, 87; *Rom1*^{-/-}2, 42; *Rom1*^{-/-}3, 44; *PRPH2* OE/*Rom1*^{-/-}1, 48; *PRPH2* OE/*Rom1*^{-/-}2, 35; *PRPH2* OE/*Rom1*^{-/-}3, 59. Data were plotted with samples separated, while statistical analysis was performed on the averages within each retina (n=3 for each genotype). One-way ANOVA with Tukey's multiple comparisons test was performed to determine statistical significance across genotypes. For the quantification of outer segment diameter (**Fig. 3C** [\[4\]](#)), 3 retinas were collected of each genotype with the number of outer segments analyzed as follows: WT1, 88; WT2, 97; WT3, 92; *Rom1*^{-/-}1, 83; *Rom1*^{-/-}2, 81; *Rom1*^{-/-}3, 85; *PRPH2* OE/*Rom1*^{-/-}1, 75; *PRPH2* OE/*Rom1*^{-/-}2, 66; *PRPH2* OE/*Rom1*^{-/-}3, 78. Data were plotted with samples separated, while statistical analysis was performed on the averages within each retina (n=3 for each genotype). One-way ANOVA with Tukey's multiple comparisons test was performed to determine statistical significance across genotypes.

Densitometric analysis of non-saturated bands from Western blots (**Figs. 5** [\[4\]](#), **6** [\[4\]](#)) was performed using Image Lab software v4.1 (Bio-Rad) and ImageJ. For quantification of the PRPH2 monomer:dimer ratio (**Fig. 5B** [\[4\]](#)), 3 samples were used for both WT and *Rom1*^{-/-} mice. Unpaired t-test was performed to determine statistical significance of the PRPH2 monomer:dimer ratio between WT and *Rom1*^{-/-} retinas. For the velocity sedimentation analysis of PRPH2 supramolecular organization (**Fig. 5E** [\[4\]](#)), 11 samples were used for WT and 4 samples were used for *Rom1*^{-/-} mice. Two-way ANOVA was performed to determine statistical significance across genotypes and fractions followed by Sidak's multiple comparisons post-hoc test to determine statistical significance between genotypes in each fraction for the disulfide-bound form. For the quantification of ROM1 and RRCT in each fraction of transfected COS7 cells (**Fig. 6E** [\[4\]](#)), 3 samples were analyzed.

All experiments were performed with mice of randomized sex. Sample sizes were determined based on previous published experiments. Data were graphed with Prism 9 (GraphPad, San Diego, CA) with error bars depicting the S.E.M. Statistical analyses were performed using Prism 9 (GraphPad). Where noted, statistical values are depicted in graphs with asterisks as follows: $p < 0.05$, *; $p < 0.01$, **; $p < 0.001$, ***; $p < 0.0001$, ****.

Acknowledgements


This work was supported by the National Institutes of Health grants EY030451 (VYA), EY005722 (VYA), EY010609 (MIN and MRA), EY034671 (MIN), EY033763 (TRL), AG070915 (SMC) and an Unrestricted Award from Research to Prevent Blindness Inc. (Duke University).

Competing interests

No competing interests declared.

Supplementary Material

Supplementary Table 1. Quantification of molar ratios between tetraspanins and rhodopsin in WT and *Rom1*^{-/-} mice – raw data



Supplementary Figure 1.

Ultrastructural analysis of WT and *Rom1*^{-/-} retinas

Representative low magnification TEM images of WT and *Rom1*^{-/-} retinas contrasted with the conventionally used osmium tetroxide. RPE: retinal pigment epithelium; OS: outer segment; IS: inner segment. Scale bar: 10 μ m.

References

- Bascom R. A., Liu L., Heckenlively J. R., Stone E. M., McInnes R. R (1995) **Mutation analysis of the ROM1 gene in retinitis pigmentosa** *Hum Mol Genet* **4**:1895–902
- Bascom R. A., Manara S., Collins L., Molday R. S., Kalnins V. I., McInnes R. R (1992) **Cloning of the cDNA for a novel photoreceptor membrane protein (rom-1) identifies a disk rim protein family implicated in human retinopathies** *Neuron* **8**:1171–84
- Burgoyne T., Meschede I. P., Burden J. J., Bailly M., Seabra M. C., Futter C. E (2015) **Rod disc renewal occurs by evagination of the ciliary plasma membrane that makes cadherin-based contacts with the inner segment** *Proc Natl Acad Sci U S A* **112**:15922–7
- Chakraborty D., Ding X. Q., Conley S. M., Fliesler S. J., Naash M. I (2009) **Differential requirements for retinal degeneration slow intermolecular disulfide-linked oligomerization in rods versus cones** *Hum Mol Genet* **18**:797–808
- Chakraborty D., Ding X. Q., Fliesler S. J., Naash M. I (2008) **Outer segment oligomerization of Rds: evidence from mouse models and subcellular fractionation** *Biochemistry* **47**:1144–56
- Cheng T., Peachey N. S., Li S., Goto Y., Cao Y., Naash M. I (1997) **The effect of peripherin/rds haploinsufficiency on rod and cone photoreceptors** *J Neurosci* **17**:8118–28
- Clarke G., Goldberg A. F., Vidgen D., Collins L., Ploder L., Schwarz L., Molday L. L., Rossant J., Szel A., Molday R. S., et al. (2000) **Rom-1 is required for rod photoreceptor viability and the regulation of disk morphogenesis** *Nat Genet* **25**:67–73
- Cohen A. I (1983) **Some cytological and initial biochemical observations on photoreceptors in retinas of rds mice** *Invest Ophthalmol Vis Sci* **24**:832–43
- Conley S. M., Stricker H. M., Naash M. I (2010) **Biochemical analysis of phenotypic diversity associated with mutations in codon 244 of the retinal degeneration slow gene** *Biochemistry* **49**:905–11
- Conley S. M., Stuck M. W., Burnett J. L., Chakraborty D., Azadi S., Fliesler S. J., Naash M. I (2014) **Insights into the mechanisms of macular degeneration associated with the R172W mutation in RDS** *Hum Mol Genet* **23**:3102–14
- Conley S. M., Stuck M. W., Watson J. N., Zulliger R., Burnett J. L., Naash M. I (2019) **Prph2 initiates outer segment morphogenesis but maturation requires Prph2/Rom1 oligomerization** *Hum Mol Genet* **28**:459–475
- Connell G. J., Molday R. S (1990) **Molecular cloning, primary structure, and orientation of the vertebrate photoreceptor cell protein peripherin in the rod outer segment disk membrane** *Biochemistry* **29**:4691–8
- Ding J. D., Salinas R. Y., Arshavsky V. Y (2015) **Discs of mammalian rod photoreceptors form through the membrane evagination mechanism** *J Cell Biol* **211**:495–502

- Ding X. Q., Nour M., Ritter L. M., Goldberg A. F., Fliesler S. J., Naash M. I (2004) **The R172W mutation in peripherin/rds causes a cone-rod dystrophy in transgenic mice** *Hum Mol Genet* **13**:2075–87
- Ding X. Q., Stricker H. M., Naash M. I (2005) **Role of the second intradiscal loop of peripherin/rds in homo and hetero associations** *Biochemistry* **44**:4897–904
- Dryja T. P., Hahn L. B., Kajiwarra K., Berson E. L (1997) **Dominant and digenic mutations in the peripherin/RDS and ROM1 genes in retinitis pigmentosa** *Invest Ophthalmol Vis Sci* **38**:1972–82
- Goldberg A. F., Fales L. M., Hurley J. B., Khattree N (2001) **Folding and subunit assembly of photoreceptor peripherin/rds is mediated by determinants within the extracellular/intradiskal EC2 domain: implications for heterogeneous molecular pathologies** *J Biol Chem* **276**:42700–6
- Goldberg A. F., Loewen C. J., Molday R. S (1998) **Cysteine residues of photoreceptor peripherin/rds: role in subunit assembly and autosomal dominant retinitis pigmentosa** *Biochemistry* **37**:680–5
- Goldberg A. F., Molday R. S (1996) **Defective subunit assembly underlies a digenic form of retinitis pigmentosa linked to mutations in peripherin/rds and rom-1** *Proc Natl Acad Sci U S A* **93**:13726–30
- Goldberg A. F., Molday R. S (1996) **Subunit composition of the peripherin/rds-rom-1 disk rim complex from rod photoreceptors: hydrodynamic evidence for a tetrameric quaternary structure** *Biochemistry* **35**:6144–9
- Goldberg A. F., Moritz O. L., Molday R. S (1995) **Heterologous expression of photoreceptor peripherin/rds and Rom-1 in COS-1 cells: assembly, interactions, and localization of multisubunit complexes** *Biochemistry* **34**:14213–9
- Hawkins R. K., Jansen H. G., Sanyal S (1985) **Development and degeneration of retina in rds mutant mice: photoreceptor abnormalities in the heterozygotes** *Exp Eye Res* **41**:701–20
- Hughes C. S., Foehr S., Garfield D. A., Furlong E. E., Steinmetz L. M., Krijgsveld J (2014) **Ultrasensitive proteome analysis using paramagnetic bead technology** *Mol Syst Biol* **10**
- Jansen H. G., Sanyal S (1984) **Development and degeneration of retina in rds mutant mice: electron microscopy** *J Comp Neurol* **224**:71–84
- Kajiwarra K., Berson E. L., Dryja T. P (1994) **Digenic retinitis pigmentosa due to mutations at the unlinked peripherin/RDS and ROM1 loci** *Science* **264**:1604–8
- Kedzierski W., Weng J., Travis G. H (1999) **Analysis of the rds/peripherin.rom1 complex in transgenic photoreceptors that express a chimeric protein** *J Biol Chem* **274**
- Landrum M. J., Lee J. M., Benson M., Brown G. R., Chao C., Chitipiralla S., Gu B., Hart J., Hoffman D., Jang W., et al. (2018) **ClinVar: improving access to variant interpretations and supporting evidence** *Nucleic Acids Res* **46**:D1062–d1067
- Lewis T. R., Makia M. S., Castillo C. M., Al-Ubaidi M. R., Naash M. I., Arshavsky V. Y (2021) **Photoreceptor Disc Enclosure Is Tightly Controlled by Peripherin-2 Oligomerization** *J Neurosci* **41**:3588–3596

- Lewis T. R., Makia M. S., Kakakhel M., Al-Ubaidi M. R., Arshavsky V. Y., Naash M. I (2020) **Photoreceptor Disc Enclosure Occurs in the Absence of Normal Peripherin-2/rds Oligomerization** *Front Cell Neurosci* **14**
- Lewis T. R., Phan S., Castillo C. M., Kim K.-Y., Coppenrath K., Thomas W., Hao Y., Skiba N. P., Horb M. E., Ellisman M. H., et al. (2023) **Photoreceptor disc incisures form as an adaptive mechanism ensuring the completion of disc enclosure** *bioRxiv* **2023**
- Li C., Ding X. Q., O'Brien J., Al-Ubaidi M. R., Naash M. I (2003) **Molecular characterization of the skate peripherin/rds gene: relationship to its orthologues and paralogues** *Invest Ophthalmol Vis Sci* **44**:2433–41
- Liang Y., Fotiadis D., Maeda T., Maeda A., Modzelewska A., Filipek S., Saperstein D. A., Engel A., Palczewski K (2004) **Rhodopsin signaling and organization in heterozygote rhodopsin knockout mice** *J Biol Chem* **279**:48189–96
- Lobanova E. S., Finkelstein S., Herrmann R., Chen Y. M., Kessler C., Michaud N. A., Trieu L. H., Strissel K. J., Burns M. E., Arshavsky V. Y (2008) **Transducin gamma-subunit sets expression levels of alpha- and beta-subunits and is crucial for rod viability** *J Neurosci* **28**:3510–20
- Lobanova E. S., Finkelstein S., Li J., Travis A. M., Hao Y., Klingeborn M., Skiba N. P., Deshaies R. J., Arshavsky V. Y (2018) **Increased proteasomal activity supports photoreceptor survival in inherited retinal degeneration** *Nat Commun* **9**
- Loewen C. J., Molday R. S (2000) **Disulfide-mediated oligomerization of Peripherin/Rds and Rom-1 in photoreceptor disk membranes. Implications for photoreceptor outer segment morphogenesis and degeneration** *J Biol Chem* **275**:5370–8
- Mattapallil M. J., Wawrousek E. F., Chan C. C., Zhao H., Roychoudhury J., Ferguson T. A., Caspi R. R (2012) **The Rd8 mutation of the Crb1 gene is present in vendor lines of C57BL/6N mice and embryonic stem cells, and confounds ocular induced mutant phenotypes** *Invest Ophthalmol Vis Sci* **53**:2921–7
- Milstein M. L., Cavanaugh B. L., Roussey N. M., Volland S., Williams D. S., Goldberg A. F. X (2020) **Multistep peripherin-2/rds self-assembly drives membrane curvature for outer segment disk architecture and photoreceptor viability** *Proc Natl Acad Sci U S A* **117**:4400–4410
- Nour M., Ding X. Q., Stricker H., Fliesler S. J., Naash M. I (2004) **Modulating expression of peripherin/rds in transgenic mice: critical levels and the effect of overexpression** *Invest Ophthalmol Vis Sci* **45**:2514–21
- Pearring J. N., Martínez-Márquez J., Willer J. R., Lieu E. C., Salinas R. Y., Arshavsky V. Y (2021) **The GARP Domain of the Rod CNG Channel's β 1-Subunit Contains Distinct Sites for Outer Segment Targeting and Connecting to the Photoreceptor Disk Rim** *J Neurosci* **41**:3094–3104
- Peeters M., Khan M., Rooijakkers A., Mulders T., Haer-Wigman L., Boon C. J. F., Klaver C. C. W., van den Born L. I., Hoyng C. B., Cremers F. P. M., et al. (2021) **PRPH2 mutation update: In silico assessment of 245 reported and 7 novel variants in patients with retinal disease** *Hum Mutat*
- Pittler S. J., Keeler C. E., Sidman R. L., Baehr W (1993) **PCR analysis of DNA from 70-year-old sections of rodless retina demonstrates identity with the mouse rd defect** *Proc Natl Acad Sci U S A* **90**:9616–9

- Pöge M., Mahamid J., Imanishi S. S., Plitzko J. M., Palczewski K., Baumeister W (2021) **Determinants shaping the nanoscale architecture of the mouse rod outer segment** *bioRxiv* **2021**
- Reig C., Martinez-Gimeno M., Carballo M (2000) **A heterozygous novel C253Y mutation in the highly conserved cysteine residues of ROM1 gene is the cause of retinitis pigmentosa in a Spanish family?** *Hum Mutat* **16**
- Sakuma H., Inana G., Murakami A., Yajima T., Weleber R. G., Murphey W. H., Gass J. D., Hotta Y., Hayakawa M., Fujiki K., et al. (1995) **A heterozygous putative null mutation in ROM1 without a mutation in peripherin/RDS in a family with retinitis pigmentosa** *Genomics* **27**:384–6
- Salinas R. Y., Pearing J. N., Ding J. D., Spencer W. J., Hao Y., Arshavsky V. Y (2017) **Photoreceptor discs form through peripherin-dependent suppression of ciliary ectosome release** *J Cell Biol* **216**:1489–1499
- Skiba N. P., Lewis T. R., Spencer W. J., Castillo C. M., Shevchenko A., Arshavsky V. Y (2023) **Absolute quantification of photoreceptor outer segment proteins** *bioRxiv* **2023**
- Spencer W. J., Lewis T. R., Pearing J. N., Arshavsky V. Y (2020) **Photoreceptor discs: built like ectosomes** *Trends Cell Biol* **30**:904–915
- Spencer W. J., Pearing J. N., Salinas R. Y., Loisel D. R., Skiba N. P., Arshavsky V. Y (2016) **Progressive Rod-Cone Degeneration (PRCD) Protein Requires N-Terminal S-Acylation and Rhodopsin Binding for Photoreceptor Outer Segment Localization and Maintaining Intracellular Stability** *Biochemistry* **55**:5028–37
- Steinberg R. H., Fisher S. K., Anderson D. H (1980) **Disc morphogenesis in vertebrate photoreceptors** *J Comp Neurol* **190**:501–8
- Stricker H. M., Ding X. Q., Quiambao A., Fliesler S. J., Naash M. I (2005) **The Cys214-->Ser mutation in peripherin/rds causes a loss-of-function phenotype in transgenic mice** *Biochem J* **388**:605–13
- Stuck M. W., Conley S. M., Naash M. I (2014) **The Y141C knockin mutation in RDS leads to complex phenotypes in the mouse** *Hum Mol Genet* **23**:6260–74
- Stuck M. W., Conley S. M., Naash M. I (2015) **Retinal Degeneration Slow (RDS) Glycosylation Plays a Role in Cone Function and in the Regulation of RDS-ROM-1 Protein Complex Formation** *J Biol Chem* **290**:27901–13
- Stuck M. W., Conley S. M., Naash M. I (2016) **PRPH2/RDS and ROM-1: Historical context, current views and future considerations** *Prog Retin Eye Res* **52**:47–63
- Tian G., Ropelewski P., Nemet I., Lee R., Lodowski K. H., Imanishi Y. (2014) **Tian, G., Ropelewski, P., Nemet, I., Lee, R., Lodowski, K. H. and Imanishi, Y. (2014). An unconventional secretory pathway mediates the cilia targeting of peripherin/rds. J Neurosci 34, 992-1006. An unconventional secretory pathway mediates the cilia targeting of peripherin/rds. J Neurosci 34**
- Travis G. H., Sutcliffe J. G., Bok D (1991) **The retinal degeneration slow (rds) gene product is a photoreceptor disc membrane-associated glycoprotein** *Neuron* **6**:61–70

Volland S., Hughes L. C., Kong C., Burgess B. L., Linberg K. A., Luna G., Zhou Z. H., Fisher S. K., Williams D. S (2015) **Three-dimensional organization of nascent rod outer segment disk membranes** *Proc Natl Acad Sci U S A* **112**:14870–5

Zulliger R., Conley S. M., Mwoyosvi M. L., Al-Ubaidi M. R., Naash M. I (2018) **Oligomerization of Prph2 and Rom1 is essential for photoreceptor outer segment formation** *Hum Mol Genet* **27**:3507–3518

Zulliger R., Conley S. M., Mwoyosvi M. L., Stuck M. W., Azadi S., Naash M. I (2015) **SNAREs Interact with Retinal Degeneration Slow and Rod Outer Segment Membrane Protein-1 during Conventional and Unconventional Outer Segment Targeting** *PLoS One* **10**

Author information

Tylor R. Lewis

Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710
ORCID iD: [0000-0001-6832-7972](https://orcid.org/0000-0001-6832-7972)

Mustafa S. Makia

Department of Biomedical Engineering, University of Houston, Houston, TX, USA, 77204

Carson M. Castillo

Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710

Ying Hao

Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710

Muayyad R. Al-Ubaidi

Department of Biomedical Engineering, University of Houston, Houston, TX, USA, 77204,
College of Optometry, University of Houston, Houston, TX, USA, 77204

Nikolai P. Skiba

Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710

Shannon M. Conley

Department of Cell Biology, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA, 73104

Vadim Y. Arshavsky

Department of Ophthalmology, Duke University Medical Center, Durham, NC, USA, 27710,
Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC, USA, 27710

For correspondence: vadim.arshavsky@duke.edu

Muna I. Naash

Department of Biomedical Engineering, University of Houston, Houston, TX, USA, 77204,
College of Optometry, University of Houston, Houston, TX, USA, 77204

For correspondence: mnaash@central.uh.edu

Editors

Reviewing Editor

Audrey Bernstein

State University of New York Upstate Medical University, United States of America

Senior Editor

Utpal Banerjee

University of California, Los Angeles, United States of America

Reviewer #1 (Public Review):

Summary:

The precise mechanism of how tetraspanin proteins engage in the generation of discs is still an open question in the field of photoreceptor biology. This question is of significance as the lack of photoreceptor discs or defects in disc morphogenesis due to mutations in tetraspanin proteins is a known cause of vision loss in humans. The authors of this study combine TEM and mouse models to tease out the role of tetraspanin proteins, peripherin, and Rom1 in the genesis of the photoreceptor discs. They show that the absence of Rom1 leads to an increase in peripherin and changes in disc morphology. Further rise in peripherin alleviates some of the defects observed in Rom1 knockout animals leading to the conclusion that peripherin can substitute for the absence of Rom1.

Strengths:

A mouse model of Rom1 generated by the McInnes group in 2000 predicted a role for Rom1 in rim closure. They also showed enlarged discs in the absence of Rom1. This study confirmed this finding and showed the compensatory changes in peripherin, maintaining the total levels of tetraspanin proteins. Lack of Rom1 leads to excessive open disks demonstrated by darkly stained tannic acid-accessible areas in TEM. Interestingly, increased peripherin expression can rescue some morphological defects, including maintaining normal disc diameters and incisures. Overall, these observations lead authors to propose a model that ROM1 can be replaced by peripherin.

Weaknesses:

The compensatory increase in peripherin and morphological rescue in the absence of ROM1 is expected, given the previous work from authors showing i) absence of peripherin showing increased ROM1 and ii) "Eliminating Rom1 also increased levels of Prph2/RRCT: mean Prph2/RRCT levels in P30 Prph2+/R retinas were 34% of WT, while levels in Prph2+/R/Rom1-/- retinas were 59% of WT" from Conley, 2019. The current study provides a comprehensive quantitative analysis. However, the mechanism behind the mechanism is unclear and warrants discussion.

Photoreceptor morphology appears better when peripherin is overexpressed. Is there a rescue of rod function (assessed by ERG or equivalent measures) in peripherin OE/Rom1-/- mice? Given the extensive work in this area and the implications the authors allude to at the end, it is important to investigate this aspect.

Reviewer #2 (Public Review):

In this study, Lewis et al seek to further define the role of ROM1. ROM1 is a tetraspanin protein that oligomerizes with another tetraspanin, PRPH2, to shape the rims of the membrane discs that comprise the light-sensitive outer segment of vertebrate photoreceptors. ROM1 knockout mice and several PRPH2 mutant mice are reexamined. The conclusion

reached is that ROM1 is redundant to PRPH2 in regulating the size of newly forming discs, although excess PRPH2 is required to compensate for the loss of ROM1.

This replicates earlier findings while adding rigor using a mass spectrometry-based approach to quantitate the ratio of ROM1 and PRPH2 to rhodopsin (the protein packed in the body of the disc membranes) and careful analysis of tannic acid labeled newly forming discs using transmission electron microscopy.

In ROM1 knockout mice PRPH2 expression was found to be increased so that the level of PRPH2 in those mice matches the combined amount of PRPH2 and ROM1 in wildtype mice. Despite this, there are defects in disc formation that are resolved when the ROM1 knockout is crossed to a PRPH2 overexpressing line. A weakness of the study is that the molar ratios between ROM1, PRPH2 and rhodopsin were not measured in the PRPH2 overexpressing mice. This would have allowed the authors to be more precise in their conclusion that a 'sufficient' excess of PRPH2 can compensate for defects in ROM1.

Reviewer #3 (Public Review):

In this manuscript, Lewis et al. investigate the role of tetraspanins in the formation of discs - the key structure of vertebrate photoreceptors essential for light reception. Two tetraspanin proteins play a role in this process: PRPH2 and ROM1. The critical contribution of PRPH2 has been well established and loss of its function is not tolerated and results in gross anatomical pathology and degeneration in both mice and humans. However, the role of ROM1 is much less understood and has been considered somewhat redundant. This paper provides a definitive answer about the long-standing uncertainty regarding the contribution of ROM1 firmly establishing its role in outer segment morphogenesis. First, using an ingenious quantitative proteomic technique the authors show PRPH2 compensatory increase in ROM1 knockout explaining the redundancy of its function. Second, they uncover that despite this compensation, ROM1 is still needed, and its loss delays disc enclosure and results in the failure to form incisures. Third, the authors used a transgenic mouse model and show that deficits seen in ROM1 KO could be completely compensated by the overexpression of PRPH2. Finally, they analyzed yet another mouse model based on double manipulation with both ROM1 loss and expression of PRPH2 mutant unable to form dimerizing disulfide bonds further arguing that PRPH2-ROM1 interactions are not required for disc enclosure. To top it off the authors complement their *in vivo* studies by a series of biochemical assays done upon reconstitution of tetraspanins in transfected cultured cells as well as fractionations of native retinas. This report is timely, addresses significant questions in cell biology of photoreceptors, and pushes the field forward in a classical area of photoreceptor biology and mechanics of membrane structure as well. The manuscript is executed at the top level of technical standard, exceptionally well written, and does not leave much more to desire. It also pushes standards of the field- one such domain is the quantitative approach to analysis of the EM images which is notoriously open to alternative interpretations - yet this study does an exceptional job unbiassing this approach.

According to my expertise in photoreceptor biology, there is nothing wrong with this manuscript either technically or conceptually and I have no concerns to express.