

Phosphorylation, disorder, and phase separation govern the behavior of Frequency in the fungal circadian clock

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint posted

November 24, 2023 (this version)

Sent for peer review

August 2, 2023

Posted to bioRxiv

June 30, 2023

Daniyal Tariq, Nicole Maurici, Bradley M. Bartholomai, Siddarth Chandrasekaran, Jay C. Dunlap, Alaji Bah , Brian R. Crane 

Department of Chemistry & Chemical Biology, Cornell University, Ithaca, NY, 14853 • Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, NY, 13210; • Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth, Hanover, NH 03755

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Circadian clocks are composed from molecular oscillators that pace rhythms of gene expression to the diurnal cycle. Therein, transcriptional-translational negative feedback loops (TTFLs) generate oscillating levels of transcriptional repressor proteins that regulate their own gene expression. In the filamentous fungus *Neurospora crassa*, the proteins Frequency (FRQ), the FRQ-interacting RNA helicase (FRH) and Casein-Kinase I (CK1) form the FFC complex that represses expression of genes activated by the White-Collar complex (WCC). A key question concerns how FRQ orchestrates molecular interactions at the core of the clock despite containing little predicted tertiary structure. We present the reconstitution and biophysical characterization of FRQ and the FFC in unphosphorylated and highly phosphorylated states. Site-specific spin labeling and pulse-dipolar ESR spectroscopy provides domain-specific structural details on the full-length, 989-residue intrinsically disordered FRQ and the FFC. FRQ contains a compact core that associates and organizes FRH and CK1 to coordinate their roles in WCC repression. FRQ phosphorylation increases conformational flexibility and alters oligomeric state but the changes in structure and dynamics are non-uniform. Full-length FRQ undergoes liquid-liquid phase separation (LLPS) to sequester FRH and CK1 and influence CK1 enzymatic activity. Although FRQ phosphorylation favors LLPS, LLPS feeds back to reduce FRQ phosphorylation by CK1 at higher temperatures. Live imaging of *Neurospora* hyphae reveals FRQ foci characteristic of condensates near the nuclear periphery. Analogous clock repressor proteins in higher organisms share little position-specific sequence identity with FRQ; yet, they contain amino-acid compositions that promote LLPS. Hence, condensate formation may be a conserved feature of eukaryotic circadian clocks.

eLife assessment

This manuscript provides a **fundamental** contribution to the understanding of the role of intrinsically disordered proteins in circadian clocks and the potential involvement of phase separation mechanisms. The authors **convincingly** report on the structural and biochemical aspects and the molecular interactions of the intrinsically disordered protein FRQ. This paper will be of interest to scientists focusing on circadian clock regulation, liquid-liquid phase separation, and phosphorylation.

Introduction

Eukaryotic circadian rhythms of biological phenomena arise from periodic gene expression patterns that oscillate independent of external cues but can be entrained to light and temperature (Hurley, et al., 2016). At the molecular level, circadian clocks function as transcription-translation negative-feedback loops (TTFLs) that consist of core positive and negative elements (Fig. 1A (Crane, et al., 2014)). In the long-standing model clock of *Neurospora crassa*, the transcriptional repressor complex is known as the FFC, named for its three principal components: Frequency (FRQ, Fig. 1B), the FRQ-interacting RNA helicase (FRH) and Casein Kinase 1 (CK1) (Cha, et al., 2014; Hurley, et al., 2016). FRQ is functionally analogous to Period (PER) in insects and mammals, whereas CK1 is an ortholog of the mammalian CK1 and the *Drosophila* kinase Doubletime (DBT) (Dunlap, et al., 2017). FRH is found only in the core repressor of the fungal clock. Although FRH is homologous to the yeast helicase MTR4, FRH helicase activity does not appear essential to clock function (Conrad, et al., 2016; Hurley, et al., 2013). The FFC represses the positive arm of the cycle, which involves two zinc-finger transcription factors, White-Collar 1 (WC-1) and White-Collar 2 (WC-2) assembled into the White-Collar Complex (WCC).

WCC binding to the clock-box (c-box) activates *frq* expression (Fig. 1A (Froehlich, et al., 2002)). FRQ production is also highly regulated at the post-transcriptional level through the use of rare codons to retard translation, the expression of an anti-sense transcript and alternative splicing mechanisms (Cha, et al., 2014; Dunlap, et al., 2017; Zhou, et al., 2013). The latter leads to the production of either a long (L-FRQ) or short (S-FRQ) isoform, and the ratio of these isoforms fine tunes period length in response to ambient temperatures (Diernfellner, et al., 2007; Liu, Y., et al., 1997; Zhou, et al., 2013). Translated FRQ binds to FRH via its FFD domain and CK1 via its two FCD domains (1 and 2) to form the repressive FFC (Cha, et al., 2014; Dunlap, et al., 2017; Jankowski, et al., 2022; Fig. 1C), although Short Linear Motifs within FRQ may also broadly influence the FRQ-interactome (Pelham, et al., 2021). The FFC enters the nucleus where it directly inhibits the WCC and thereby down-regulates *frq* expression (Hurley, et al., 2013). FRQ-mediated WCC phosphorylation by CK1 (and associated kinases) reduces interactions between the WCC and the c-box to close the feedback loop and restart the cycle (Fig. 1A (Wang, et al., 2019)). FRH facilitates interactions between the FFC and the WCC (Conrad, et al., 2016) and thus, repressor function likely depends upon the FRQ-mediated association of CK1 and FRH. Whereas interaction regions on the FRQ protein sequence have been identified (Fig. 1C), there is little information on the spatial arrangement of the FFC components.

Despite low sequence conservation among the core negative arm proteins across species, they all share a high degree of disorder (Pelham, et al., 2020). FRQ, the analogous *Drosophila* (d) Period (PER) protein and the analogous mammalian (m) PER have > 65% disorder predicted in their sequences with ~86% of the FRQ sequence assumed to have no defined structure (Pelham, et al., 2020). Both FRQ and mPER2 have biochemical properties characteristic of intrinsically




Figure 1

FRQ is a large, dynamic component of the clock oscillator

(A) The TTFL in *Neurospora crassa*. In the core circadian oscillator, FRQ associates with FRH and CK1 to form a repressor complex (the FFC). This complex inhibits the positive-acting transcription-factor known as the white-collar complex WCC, composed of White Collar-1 (WC-1) and White Collar-2 (WC-2). The WCC activates *frq* expression by binding to the c-box region in the *frq* promoter. Stoichiometries of the components are not represented in the schematic. (B) AlphaFold derived model of LFRQ color coded by pLDDT score that highlights the confidence in each predicted region. (C) Functional domains of FRQ showing the start position for the LFRQ isoform (AUG_{LFRQ}) and the four native cysteine residues. Characterized domains include: CC – coiled coil domain, FCD – FRQ:CK1 interacting domain, PEST – proline, glutamic acid, serine, and threonine-rich domain and FFD – FRQ:FRH interacting domain.

disordered proteins (IDPs) (Hurley, et al., 2013 [↗](#); Marzoll, et al., 2022 [↗](#)). These clock repressors also undergo extensive posttranslational modification (PTM), particularly phosphorylation (Dunlap, et al., 2018; Pelham, et al., 2020 [↗](#)). FRQ acquires over 100 phosphorylation sites over the course of the circadian day (similar to PER phosphorylation by DBT), predominantly due to CK1 (Baker, et al., 2009 [↗](#); Tang, et al., 2009 [↗](#)). The phosphorylation events have two consequences: 1) those that weaken the interaction between FRQ and the WCC or CK1 (Cha, et al., 2014 [↗](#); Liu, X., et al., 2019); and 2) those that direct FRQ down the ubiquitin-proteasome pathway (Larrondo, et al., 2015 [↗](#)), although only the former are required for circadian rhythmicity (Larrondo, et al., 2015 [↗](#)). Multi-site phosphorylation of FRQ influences its inhibitory function and clock period length (Larrondo, et al., 2015 [↗](#); Liu, X., et al., 2019). Although the consequences of phosphorylation-driven conformational changes in FRQ have been probed by proteolytic sensitivity, the challenges of producing FRQ have limited direct assessment of its conformational properties (Liu, X., et al., 2019; Pelham, et al., 2020 [↗](#); Querfurth, et al., 2011 [↗](#)). IDPs such as FRQ are known to participate in liquid-liquid phase separation (LLPS), which has emerged as a key mechanism for organizing cellular components, coordinating metabolism and directing signaling through, for example, the formation of membraneless organelles (MLOs) (Dignon, et al., 2020 [↗](#); Wright, et al., 2014). Emerging studies in plants (Jung, et al., 2020 [↗](#)) and flies (Xiao, et al., 2021 [↗](#)) implicate LLPS in circadian clocks, and in *Neurospora* it has recently been shown that the Period-2 (PRD-2) RNA-binding protein influences *frq* mRNA localization through a mechanism potentially mediated by LLPS (Bartholomai, et al., 2022 [↗](#)). Intriguing questions surround why clock repressor proteins are so disordered, undergo extensive post-translational modification and whether LLPS has a role to play in their function.

Clock proteins and their complexes have been challenging to characterize due to their large size (>100 kDa), low sequence complexity, extensive disorder and modification states. Herein, we reconstitute FRQ and the FFC to assess the global and site-specific structural properties of these moieties, as well as the dependence of their properties on multi-site phosphorylation. FRQ undergoes non-uniform conformational change upon phosphorylation and arranges FRH close to CK1 in the FFC. We also demonstrate that FRQ indeed participates in phosphorylation-dependent LLPS in vitro, and as a condensate sequesters FRH and CK1 while altering CK1 activity. Our results suggest how a massively disordered protein such as FRQ organizes the clock core oscillator and reveal the effects of extensive phosphorylation on the conformational and phase-separation properties of a large IDP.

Results

Sequence assessment and general properties of FRQ

Analysis of an AlphaFold 2.0 (Jumper, et al., 2021 [↗](#)) model of full-length FRQ indicates that FRQ lacks a defined arrangement of large folded domains; instead, most of the molecule is unstructured (**Fig. 1B** [↗](#)), in agreement with previous studies predicting that FRQ mostly comprises intrinsically disordered regions (Hurley, et al., 2013 [↗](#)). However, FRQ is also predicted to contain a small, clustered core, with several secondary structural elements that collapse together to form a compact nucleus surrounded by long flexible disordered regions (**Fig. 1B** [↗](#)). The AlphaFold pLDDT scores, which represent the confidence in the structural prediction (Jumper, et al., 2021 [↗](#)), are very low across the model, except within the core (**Fig. 1B** [↗](#), **Fig. S1** [↗](#)), which contains the CK1 binding region. The FRH binding region, FFD, falls within a more flexible region with a lower pLDDT score than that of CK1. Helices compose the majority of secondary structural elements predicted by AlphaFold (**Fig. 1B** [↗](#)). The assignments of the helical regions were well supported by the Agadir program that predicts helical propensity based on the helix/coil transition theory (Muñoz, et al., 1994). Four regions distributed throughout the length of FRQ (residues 194-212, 312-329, 709-719 and 801-817) have at least 10% helical propensity and these helical regions mostly agree with those of the AlphaFold model (**Fig. 1B** [↗](#)). Together, these analyses primarily

indicate that FRQ assumes a broad conformational ensemble containing an ordered nucleus that interacts extensively with partially ordered and flexible regions. However, much of the structural details of the model, including the positioning and interactions between secondary structural elements, are predicted with low confidence, thereby underscoring the need for a multi-pronged biophysical study of the protein.

Reconstitution of FRQ in unphosphorylated and highly phosphorylated states

The size and intrinsic disorder of FRQ have made it challenging to recombinantly express and purify, thereby limiting investigations of its biophysical properties. Previous purifications have characterized FRQ as an IDP, but suffered from low yields and aggregated species (Hurley, et al., 2013 [↗](#); Lauinger, et al., 2014 [↗](#)). We also encountered extensive protein degradation and aggregation with conventional purification methods and affinity tagging strategies; however, methods proven effective for IDPs (Pedersen, et al., 2020 [↗](#)) yielded comparatively well-behaved protein. Briefly, we carried out Nickel affinity chromatography and size exclusion chromatography (SEC) under denaturing conditions to maximize yield and reduce aggregation of full-length L-FRQ (residues 1-989; **Fig. 2A** [↗](#)) (Seal, et al., 2021 [↗](#)). Solubilization with high concentrations of guanidinium-HCl prevented proteolysis, aggregation and limited co-purification of impurities that were otherwise difficult to remove. Once FRQ was isolated, the protein remained soluble, monodisperse and could be concentrated after the denaturant was removed by dialysis (**Fig. 2A,B** [↗](#)). To our knowledge, FRQ is one of the largest native IDPs produced in amounts amenable to structural and biophysical characterization.

To generate phosphorylated FRQ, the protein was co-expressed with (untagged) CK1, which forms a stable complex with FRQ and acts as its primary kinase (Querfurth, et al., 2011 [↗](#)). The same denaturing purification procedure was followed as for FRQ expressed without CK1. Phosphorylation was confirmed visually by altered migration on SDS-PAGE (**Fig. 2A** [↗](#)) and mass spectrometry (**Fig. S2** [↗](#)). The latter revealed a network of > 80 phosphorylation sites (**Fig. S2** [↗](#)), most of which agree with those found on natively expressed FRQ (Baker, et al., 2009 [↗](#); Tang, et al., 2009 [↗](#)). Hereafter, we refer to this preparation as phosphorylated FRQ (p-FRQ) whereas non-phosphorylated FRQ (np-FRQ) refers to FRQ expressed without CK1. In general, we found that co-expression of FRQ with CK1 increased protein yield and stability, while limiting aggregation and degradation.

Global properties of FRQ

Size Exclusion Chromatography coupled to Multi Angle Light Scattering (SEC-MALS) of p-FRQ gives a molecular weight measurement consistent with a primarily monomeric state (MW of ~143 kDa versus a predicted MW of ~110 kDa; **Fig. 2B** [↗](#)). In contrast, the np-FRQ measured by MALS indicates a dimer and its measured mass distribution increases upon concentration (**Fig. 2B** [↗](#)). Hence, purified np-FRQ dimerizes, whereas phosphorylation destabilizes this association. SEC-coupled small-angle x-ray scattering (SAXS) analyses revealed that both p-FRQ and np-FRQ shared characteristics of highly flexible proteins, but that p-FRQ was more extended than np-FRQ (Table S1). The dimensionless Kratky plots of p-FRQ and np-FRQ, which report on compactness and conformational dispersity (Durand, et al., 2010 [↗](#); Martin, Erik W., et al., 2021; Sagar, et al., 2020 [↗](#); Trewthella, et al., 2017 [↗](#)), revealed high degrees of flexibility for both proteins (**Fig. 2C** [↗](#)). The peak values of the plot were shifted from those expected for compact structures but were also not indicative of fully disordered proteins (Durand, et al., 2010 [↗](#); Rambo, et al., 2011; Sagar, et al., 2020 [↗](#)). Furthermore, p-FRQ appears more disordered than np-FRQ, a characteristic also reflected in their pairwise distance distributions (**Fig. S3** [↗](#)). These data then indicate that FRQ generally adopts a relatively monodisperse conformational state that is partially compact but highly flexible and then increases expanse and flexibility upon phosphorylation. This general picture is consistent with the increased susceptibility to proteolysis that FRQ experiences with phosphorylation (Liu, X., et al., 2019; Pelham, et al., 2020 [↗](#); Querfurth, et al., 2011 [↗](#)).




Figure 2

Purification of FRQ in non-phosphorylated and phosphorylated forms reveals IDP behavior

(A) SDS-PAGE gel showing non-phosphorylated (np-FRQ) and phosphorylated FRQ (p-FRQ) alongside a molecular weight ladder. Both samples represent the purified full-length long isoform (L-FRQ, residues 1-989) in non-denaturing buffer. (B) Size exclusion chromatography–multiangle light scattering (SEC-MALS) of phosphorylated FRQ (p-FRQ) and non-phosphorylated (np-FRQ). The molecular weight (MW) of p-FRQ ($\sim 143 \pm 1$ kDa) is somewhat larger than that of an unphosphorylated monomer (~ 120 kDa), but substantially smaller than that of a dimer, whereas the high concentration MW value of np-FRQ ($\sim 261 \pm 1$ kDa) is slightly larger than that expected for a dimer (~ 240 kDa). Note that the p-FRQ MW is affected by >80 phospho sites. The shoulder to the right of the main peak had low light-scattering and a large MW error. The colored lines represent the MW of each sample. (C) Dimensionless Kratky plot of SAXS data for p-FRQ and np-FRQ, in both cases the peak positions of the curves are shifted from that expected for a globular protein ($\sqrt{3}, 1.104$), but neither plateau at 2, which would be characteristic of a flexible, denatured polypeptide (Durand, et al., 2010). See Table S1 for additional SAXS parameters. Total protein concentration was between 50–75 μM for the MALS and SAXS experiments.

Local measures of FRQ structure and dynamics

The properties of FRQ that include its intrinsic disorder and propensity to form heterogeneous ensembles are not well suited to commonly applied structural characterization methods. We applied site-specific spin labeling (**Fig. 3A**) and electron spin resonance spectroscopy (ESR) to monitor FRQ local structural properties. FRQ has four native cysteine residues that provide reactive sites for nitroxide spin-labels (**Fig. 3A,B** (Berliner, et al., 1982)). The Cys residues distribute relatively evenly throughout the protein sequence and fall within areas of varying structural order that include the binding regions for the core clock proteins (FRH, CK1, WC1 and WC2; Table S2-3; **Fig. 3B**). Furthermore, we developed and applied enzymatic SortaseA and AEP1 peptide-coupling methods (Chandrasekaran, et al., 2021; Nguyen, et al., 2015) to spin-label FRQ at its N- and C-termini (**Fig. 3A**; **Fig. S4**).

Line-shape analysis from continuous wave (cw)-ESR spectroscopy reports on local dynamics (Franck, et al., 2015; Freed, 2005). For rapidly rotating nitroxide spin labels (SLs), the characteristic spectrum consists of three narrow lines of equal width and intensity. As the rotational mobility decreases, the lines broaden, especially the high-field line (Bonucci, et al., 2020; Emmanouilidis, et al., 2021). Cw-ESR reveals that FRQ has regions of varying order that agree with computational predictions and that some of these sites undergo phosphorylation-dependent conformational changes (**Fig. 3C**). Spin-labels attached to Cys490 and Cys658 (490SL and 658SL) are predicted to be within an α -helix and a β -strand respectively. These sites correspondingly showed a broad low-field (left-most line) component and a small high-field component (right-most line) indicative of slower tumbling compared to sites predicted to be in flexible regions, such as 746SL (**Fig. 3C**). The 490SL site had a broader profile compared to that of the 658SL site, although both are relatively well ordered. We observed a similar trend with ρ , which is the peak-to-peak width of the central line (Hubbell, et al., 2000). 490SL and 658SL had similar ρ values that were higher than that of 746SL, indicating that they are less flexible than 746SL (Table S4). Furthermore, the spectra also highlighted structural differences between the phospho forms: p-FRQ and np-FRQ gave very different ρ values at 490SL and the C-terminus, thereby demonstrating how phosphorylation can influence conformational properties site-specifically. The C-terminus may have become more dynamic in p-FRQ as compared to np-FRQ, because the C-terminus contains a cluster of phospho-sites (**Fig. 3C**). In contrast, the N-terminus and especially 490SL, which locates to the CK1-interaction region of the FCD₂ domain, are more ordered in p-FRQ. Such differences in mobility between p-FRQ and np-FRQ in the FCD (490SL) may reflect changes in FRQ:CK1 interactions as FRQ becomes progressively phosphorylated across the circadian period (Liu, X., et al., 2019).

Double electron-electron resonance (DEER) spectroscopy provides distance distribution between two nitroxide labels in the range of 15 Å to ~100 Å. We produced 6 cysteine pair variants that each contained only two of the four native FRQ Cys residues to measure pair-wise distances and one single Cys variant to test for oligomerization. Owing to association and aggregation when concentrated, np-FRQ was not amenable to DEER measurements; however, such issues were not prohibitive with p-FRQ. Nevertheless, due to the challenges of spin-labeling appreciable concentrations of p-FRQ, the modulation depths of the DEER data, which reflect the number of spins in range to produce specific dipolar interactions, are small (~2-4%). These values compare to ~10-12% for a well-ordered, highly labeled pair of sites under these instrument conditions (Dunleavy, et al., 2023). The single Cys spin-labeled variant 490SL produced log-linear time domain traces indicative of a uniform background with little evidence of dimerization, consistent with the MALS data showing p-FRQ to be monomeric (**Fig. S5**). A second single Cys variant, 746C likewise produced little dipolar signal, although for this variant, solubility issues limited spin concentration (**Fig. S5**). The double 370SL:746SL p-FRQ variant also gave no dipolar signals and hence no dimerization as reported by those two sites (**Fig. 3D**). When at least one of the two spin labels resides outside of the predicted core, the DEER experiments of the spin pairs generally revealed small, albeit significantly slow decaying time domain traces that deviate from background (**Fig. 3D**; **Fig. S5-6**). Such traces are indicative of long inter-spin distances and




Figure 3

ESR spectroscopy studies of spin-labeled FRQ.

(A) Spin-labeling strategies for FRQ. MTSL was used for internal labeling of cysteine residues and AEP1 and SortaseA were used to label the N and C terminus, respectively. (B) AlphaFold generated model of FRQ with the positions of native spin-labeled cysteine residues highlighted in dark blue. (C) X-band (9.8 GHz) room temperature cw-ESR measurements for various spin-labeled FRQ variants, each containing a single Cys labeling site, as designated. Data represented as first-derivative spectra with the horizontal axes depicting the static field strength (typically 3330 to 3400 G), and the vertical axis depicting the change in magnetic susceptibility with respect to the static field, in arbitrary units. (D) Time-domain data of DEER measurements for double-cysteine labeled FRQ variants. The baseline-corrected time domain dipolar evolution trace fits are shown; details are given in Fig. S5-6. The 490-658 separation is noticeably closer and more ordered than the others. The total protein concentration was between 50-75 μM for all experiments.

conformational flexibility, which would be consistent with the large mobility and intrinsic disorder of these peripheral regions (**Fig. S5** [4-6](#)). Only variant 490SL:658SL gave a fast-decaying time-domain signal, indicative of short inter-spin distances and conformational rigidity (**Fig. S5** [4-6](#)). Despite the overall uncertainty in the AlphaFold model (**Fig. 1B [4](#)), it is notable that 490SL and 658SL both fall within the predicted ordered core.**

FFC reconstitution and component arrangement

The FFC was generated by mixing p-FRQ or np-FRQ with excess FRH and CK1, followed by SEC purification to give p-FFC or np-FFC, respectively. FRHtiN (residues 100-1106), which binds FRQ, was used to improve expression and enhance stability (Conrad, et al., 2016 [4](#); Hurley, et al., 2013 [4](#)). The resulting symmetric SEC traces and SDS-PAGE gels revealed pure and homogenous ternary complexes in each case (**Fig. 4A** [4](#)). As expected, CK1 phosphorylates FRQ in np-FFC (**Fig. 4A** [4](#), **Fig. S7** [4](#)), and this activity does not require priming by other kinases (Marzoll, et al., 2022 [4](#)). SEC-SAXS analysis of the FFC showed that both FRQ phospho-forms produced complexes that were more globular than FRQ alone, but still somewhat flexible, with the p-FFC being more flexible than the np-FFC (**Fig. 4B** [4](#), **Fig. S3** [4](#)). MW estimates of the p-FFC and np-FFC from SAXS indicate a 1:1 stoichiometry of FRQ:FRH, with a likely single component of CK1 also contained within the complex (Table S1).

DEER measurements from selectively spin-labeled components within the complex were used to probe its overall architecture (**Fig. 5** [4](#); **Fig. S8** [4](#)-**S11** [4](#)). An ADP- β SL probe (Muok, et al., 2018 [4](#)) bound to both FRH and CK1 and revealed that the FRH and CK1 ATP-binding pockets are closely positioned within the FFC complex (**Fig. 5A** [4](#); **Fig. S8** [4](#)). Nonetheless, the presence of broad and multiple peaks in the distance distributions indicated conformational heterogeneity. Spin labels placed within the FCD2 and FFD domains of FRQ (490C and 746C), which bind CK1 and FRH respectively, produced a distance distribution similar to that of the labels that target the ATP binding sites of the two enzymes (**Fig. 5A** [4](#)) (Cheng, et al., 2005 [4](#)). These independent observations support the conclusion that FRQ organizes FRH and CK1 in close proximity within the FFC.

Spin-label separations within p-FRQ changed substantially when FRH and CK1 bound (**Fig. 5** [4](#); **Fig. S5** [4](#)-**6** [4](#), **S9** [4](#)-**11** [4](#)). In general, the time domain decays were more rapid in the FFC compared to p-FRQ alone and the resulting distance distributions had smaller mean distances and less breadth (**Fig. 5B-H** [4](#)). However, the changes were not uniform; for example, the 658SL-to-746SL distance distribution increased in the complex compared to free FRQ (**Fig. 5G** [4](#)). To further assess how the distributions p-FRQ conformations changed when the p-FFC formed, the distance data was fit to a model of one near and one far Gaussian functions (**Fig. S11** [4](#); Table S5). Shifts in population between the near and far components were most prominent for the 370-658, 490-658 and 490-746 variants, which contain sites involved in core clock interactions (**Fig. S11** [4](#)). The proportion of the closer component increased from ~30% to ~70% in the 370-658 variant (**Fig. S11** [4](#)) and from ~10% to ~70% in the 490-746 variant (**Fig. 5F** [4](#)) when bound to FRH and CK1. Similar trends were observed when we compared the percentage change of the average separation observed in FRQ alone versus when it is bound to FRH and CK1 (**Fig. 5H** [4](#)). The 490, 658 and 746 sites reside in regions important for WCC interactions (Guo, et al., 2010 [4](#)) and FRH may organize such binding determinants. Notably, the 370SL:746SL spin pair showed little dipolar coupling in free p-FRQ but resolved into an observable mid-range separation in the presence of CK1 and FRH (**Fig. 5C** [4](#)), perhaps because position 746 is within the FRH binding region. Thus, CK1 and FRH order p-FRQ to an appreciable degree and the two enzymes are adjacently positioned within the conformationally heterogeneous complex.




Figure 4

Properties of the FFC.

(A) SEC of the FFC; left inset: SDS-PAGE gel inset of the components; right inset: 32 P-autoradiogram of FRQ phosphorylation by CK1 and CK1 autophosphorylation at 10 and 20 μ M FRQ all in the presence of FRH. Complete autoradiograms are shown in [Fig. S7](#). (B) Overlay of dimensionless Kratky plots of SAXS data from the FFC formed with either np- or p-FRQ (protein concentrations were between 50-75 μ M). The SAXS-derived MW of np-FRQ FFC = 285 kDa, which compares to a predicted value of 273 kDa; MW of p-FFC = 340 kDa; Table S1).




Figure 5

PDS distance distributions of spin-labeled FRQ and the FFC.

(A) DEER-derived distance distributions of the FFC wherein the spins were directed by ADP linkage to target the FRH and CK1 ATP-binding sites (blue) or in the form of MTSL labeling of cysteine residues within the FCD2 and FFD domains of FRQ (red). Time-domain data and distance distributions with error estimates are shown in [Fig. S5](#)–[S10](#). (B–G) DEER-derived distance distributions of cysteine residue pairs labeled with MTSL within p-FRQ, either alone (blue) or within the FFC (red). (H) Percentage change in the average separation of FRQ when it binds FRH and CK1. Errors are derived from the uncertainties in the distance distributions shown in [Fig. S9](#)–[S10](#). In these experiments, only FRQ was labeled with MTSL at the positions noted. Note: Total protein concentrations were between 50–75 μ M for all experiments.

FRQ exhibits behavior consistent with LLPS *in vitro* and *in vivo*

The unstructured regions and multivalent interactions of IDPs can lead to liquid-liquid phase separation (LLPS) (Martin, E. W., et al., 2020; Martin, E. W., et al., 2020). Such de-mixing often manifests in the formation of biomolecular condensates and membrane-less organelles (MLOs) within cells (Posey, et al., 2018 [🔗](#)). As has been previously noted (Pelham, et al., 2020 [🔗](#)), FRQ may be capable of LLPS owing to its extensive disorder. We found that FRQ has a high propensity to undergo LLPS based on a set of computational tools that evaluate charge distribution and clustering, long-range aromatic and non-aromatic $\pi:\pi$ contacts and nucleic acid binding propensities (Bolognesi, et al., 2016 [🔗](#); Vernon, et al., 2018 [🔗](#)). The overall FRQ PScore, which grades LLPS propensity on these properties (Vernon, et al., 2018 [🔗](#)), was 6.39, with three regions scoring above a value of 2, the threshold for favorable LLPS (**Fig. S12** [🔗](#)). The region with the highest PScore also accounted for the majority of LLPS propensity as determined by the catGRANULE prediction algorithm (Bolognesi, et al., 2016 [🔗](#)). Furthermore, we found that these methods also predicted the functional analogs dPER and hPER1 to phase separate (**Fig. 6A** [🔗](#), **Fig. S13** [🔗](#)). The FRQ functional analogs share physicochemical characteristics with FRQ, particularly an enrichment of hydrophilic residues, increases in Gly and Pro content and a depletion of hydrophobic residues, even though their positional sequence conservation with FRQ is low (**Fig. S14** [🔗](#) and **S15**; see also (Jankowski, et al., 2022 [🔗](#); Pelham, et al., 2021 [🔗](#))). Similar amino acid composition patterns among FRQ, dPER and hPER1, include the most common residues in each protein and its UniRef50 cluster (sequences with at least 50% identity with the target protein) being serine, proline and glycine (~30% of total sequence) accompanied by a high depletion in aromatic residues (only ~7.5%) (**Fig. S15** [🔗](#)). In addition, all of the sequences have a reduced proportion of hydrophobic residues such as A, I, L, and V and a slightly acidic isoelectric point. Thus, FRQ, dPER and hPER1 all have low overall hydropathy (**Fig. S14** [🔗](#)). However, in addition, FRQ and its homologs are unusual in their high Arg and Asp content, which tend to separate in charge blocks throughout their sequences (as indicated by high kappa values, **Fig. S15** [🔗](#); see also (Jankowski, et al., 2022 [🔗](#))). Such charge clustering correlates with LLPS (Somjee, et al., 2020 [🔗](#)). Overall, this analysis suggests that like intrinsic disorder, LLPS driven by the physiochemical properties of the constituent residues, may be a conserved feature in the negative arm of the circadian clock.

We tested whether both np-FRQ and p-FRQ can undergo LLPS *in vitro*. In each case, upon exchanging the purified protein into their respective phase separation buffers (which lack any crowding agents), we observed the appearance of a turbid solution that contained microscopic droplets (**Fig. 6B** [🔗](#)). These droplets scaled in size and number with increasing protein concentration, dissolved upon the addition of 1,6-hexanediol, and were shown to dock and fuse to one another, thereby confirming their liquid-like properties (**Fig. 6B** [🔗](#), **Fig. S17** [🔗](#)). UV-Vis turbidity assays as a function of temperature revealed that FRQ undergoes a low critical solution temperature (LCST) phase transition that is concentration and phosphorylation status-dependent. (**Fig. 6C** [🔗](#); **Fig. S17** [🔗](#), Tables S6-7). As expected, phase separation increased as temperature and protein concentration increased as indicated by growing amplitudes and earlier phase separation initiation temperatures (**Fig. 6C** [🔗](#); **Fig. S17** [🔗](#); Table S6-7). Lower salt concentrations favored the LLPS of p-FRQ, whereas higher salt disfavored LLPS, compared to higher salt concentrations favoring LLPS for np-FRQ (**Fig. S17** [🔗](#)). Although the NaCl concentration for np-FRQ LLPS conditions is non-physiological *per se*, the ionic strength conditions and effective solute concentrations in cells are comparatively high (Liu, B. Q., et al., 2017). Furthermore, the higher absorbance values of p-FRQ at lower temperatures relative to that of np-FRQ indicates complex phase behavior for the phosphorylated isoform that cannot adequately be explained by an LCST phase transition alone (**Fig. S17** [🔗](#)). At temperatures below the approximated phase boundary indicated by the temperature transitions (**Fig. 6C** [🔗](#); **Fig. S17** [🔗](#)) some p-FRQ phase separation has already occurred in the phase-separation buffer. Thus, FRQ phosphorylation alters its behavior consistent with LLPS as evidenced by its non-zero, concentration-dependent absorbance at lower




Figure 6

FRQ undergoes LLPS in agreement with predictions from its sequence properties and those of its functional analogs.

(A) LLPS propensity predictions using “Pi-Pi” and IUPred2A disorder predictions are shown for FRQ (purple), dPER1 (teal) and hPER1 (orange) as compared to values for proteins of similar length from their respective proteomes. The mean and standard deviations are represented as horizontal and vertical black bars, respectively. Note: **** denotes a p-value <0.0001 obtained from a Mann-Whitney U-test (B) DIC microscopy images of various concentrations of p-FRQ (in 150 mM NaCl, 25 mM Tris pH 8.0) and np-FRQ (in 500 mM NaCl, 25 mM Tris pH 8.0) at 25°C; images shown at 100x magnification. Scale bar = 2 μ m (C) Temperature vs concentration phase diagram derived from the results of the turbidity assays shown in Fig. S17. Both p-FRQ and np-FRQ undergo an LCST phase transition on or above the line (a second order polynomial fit to the data points). The error bars of each point (about the size of the points) reflect the 95% confidence interval of the mean. The phase boundary for p-FRQ represents the transition apparent in the temperature scans of Fig. S17. Some p-FRQ phase separation already occurs at low temperature under these buffer conditions and thus the region below the line does not represent fully soluble p-FRQ. (D) Live cell fluorescent imaging of FRQ^{mNeonGreen} in nuclei of *Neurospora* hyphae. Upper left panel shows the FRQ channel, represented as a multicolor heat map of fluorescence intensity; upper right panel shows a surface plot derived from the raw FRQ image to emphasize regions of concentration; lower left shows nucleoporin SON-1^{mApple}, which localizes to the cytoplasmic face of the nuclear envelope; bottom right shows the FRQ:SON-1 merge image. The images were acquired on a Zeiss 880 laser scanning confocal microscope and were smoothed by bicubic interpolation during 10-fold enlargement from 42 $\times$ 42 pixels to 420 $\times$ 420 pixels. (E) FRQ[mNeonGreen] nuclear foci shown within an *N. crassa* syncytium mycelium (outlined by white lines), cropped from movie S1. Scale bar = 2 μ m.

temperatures, mostly likely by increasing the relative negative charges, changing the charge patterns, and providing more sites for multivalent interactions within the molecule (**Fig. 6C**, **Fig. S16–17**) (Somjee, et al., 2020; Szabó, et al., 2022).

Live cell imaging of *Neurospora crassa* expressing FRQ tagged with the fluorescent protein mNeonGreen at its C-terminus (FRQ^{mNeonGreen}) revealed heterogeneous patterning of FRQ in nuclei (**Fig. 6D,E**). Live-cell imaging in *Neurospora crassa* offers atypical challenges because the mycelia grow as syncytia, with continuous rapid nuclei motion during the time of imaging. This constant movement of nuclei is compounded by the very low intranuclear abundance of FRQ and the small size of fungal nuclei, making not readily feasible visualization of intranuclear droplet fission/fusion cycles or intranuclear fluorescent photobleaching recovery experiments (FRAP) that could report on liquid-like properties. Nonetheless, bright and dynamic foci-like spots were observed near the nuclear periphery, which is delineated by the cytoplasm facing nucleoporin Son-1 tagged with mApple at its C-terminus (**Fig. 6D,E**, Movie S1). Such foci are characteristic of phase separated IDPs (Bartholomai, et al., 2022; Caragliano, et al., 2022; Gonzalez, A., et al., 2021; Tatavosian, et al., 2019) and share similar patterning to that seen for clock proteins in *Drosophila* (Meyer, et al., 2006; Xiao, et al., 2021).

Phase separated FRQ recruits FRH and CK1 and alters CK1 activity

The PScore and catGRANULE LLPS predictors gave low overall scores for the FRQ partners, FRH and CK1 (**Fig. S13**). This result is unsurprising because these proteins consist of predominantly folded domains. Nonetheless, FRH and CK1 may still act as clients that localize to the FRQ-formed LLPS-like droplets. We chose to investigate whether FRH and CK1 could be recruited into droplets with fluorescent microscopy. Structured proteins are not well tolerated in the temperature dependent turbidity assays because of the large signal that they give upon domain unfolding. We found that fluorescently-labeled FRH and CK1 were recruited into phase separated FRQ droplets, but a control protein labeled with the same fluorophore, *E. coli* CheY, was not (**Fig. 7A**). The ability of FRQ to recruit its partner proteins into phase separated droplets provides a means to sequester them and potentially control their availabilities and activities.

LLPS has been suggested to modulate the activities of enzymes by creating a microenvironment to alter substrate-sequestration and product-release (Shapiro, et al., 2021). We tested the effect of FRQ LLPS on its phosphorylation by CK1, a key reaction for regulating the period of the clock (Liu, X., et al., 2019). We compared the phosphorylation of FRQ by CK1 in phase separation buffer under different temperatures, using the latter as a means to control the degree of LLPS. Both the UV-Vis turbidity assays as well as the DIC micrographs of the protein at different temperatures establish that FRQ LLPS increases substantially with temperature (**Fig. 6C**; **Fig. 7C**). Furthermore, LLPS of p-FRQ and np-FRQ both depend highly buffer composition (**Fig. S17**). CK1 autophosphorylation served as an internal control and its temperature dependence was consistent with previous studies that show increased catalytic activity at higher temperatures (Marzoll, et al., 2022). As temperature increases drove more LLPS, the phosphorylation of FRQ decreased, even as CK1 autophosphorylation increased (**Fig. 7B**; **Fig. S19**). Quantitatively, we observed a reduction in FRQ phosphorylation by two-fold under LLPS conditions relative to conditions under which it does not phase separate (**Fig. 7B**).

Through application of cw-ESR, we probed FRQ dynamics when it undergoes LLPS. We observed a striking rigidification of spin-labeled FRQ when LLPS was promoted by buffer conditions (**Fig. 7D**). The reduction of CK1-mediated FRQ phosphorylation, which likely requires substantial dynamics given the distribution of phosphorylation sites throughout the FRQ sequence, may reflect this change in molecular dynamics. Thus, in addition to affecting the localization of FRQ and its partners, LLPS also impacts the enzymatic activity of the complex, by perhaps modulating molecular encounters and motions essential for large scale phosphorylation.




Figure 7

Conditions favoring LLPS alter the structural and enzymatic properties of the FFC.

(A) Differential Interference Contrast (DIC) and fluorescence microscopy images at 25 °C of phase separated np-FRQ or p-FRQ (5 μ M each) droplets in 500 mM KCl, 20mM Na_2HPO_4 , pH 7.4 with either equimolar Cy5-labeled FRH, CK1 or CheY (control) visualized using Cy5 fluorescence. Scale bar = 2 μ m . (B) (Top) Autoradiography of FRQ (20 μ M) (left) phosphorylated by CK1 in the presence of FRH at increasing temperatures. (Bottom) Schematic showing the nature and results of the autoradiography assay depicted above. The FRQ phosphorylation under LLPS conditions was reduced relative to non-LLPS conditions. The complete autoradiograms are shown in [Fig. S13](#). Quantification of FRQ phosphorylation by CK1 at RT under LLPS and non-LLPS conditions with phosphorylation levels under non-LLPS conditions normalized to 1. Note: *** denotes a p-value $\ll$ 0.05 obtained from a student's t-test. (C) DIC microscopy images of 15 μ M phase separated np-FRQ under the same conditions (i.e. buffer and temperature) as the phosphorylation assay shown in (B). Scale bar = 2 μ m . (D) X-band (9.8GHz) RT cw-ESR spectra of FRQ labeled with MTSL at the 490 site in solubilizing buffer (black) and under conditions that promote LLPS (cyan). Data represented as first-derivative spectra with the horizontal axes depicting the static field strength (typically 3330 to 3400 G), and the vertical axis depicting the change in magnetic susceptibility with respect to the static field, in arbitrary units.

Within the TTFLs that define eukaryotic circadian clocks, large repressor proteins like FRQ compose cellular timing mechanisms by coordinating transient protein-protein interactions, nuclear entry, targeted protein degradation, and transcriptional repression (Pelham, et al., 2020). They compose dynamic macromolecular assemblies that achieve specificity in their functions despite assuming highly variable structures (Pelham, et al., 2020). Our purification scheme enabled a multi-pronged biophysical characterization of FRQ. The majority of previous such studies on IDPs, particularly those that undergo de-mixing, have involved truncated proteins, excised domains or peptides of sizes less than 200 residues (Babinchak, et al., 2019; Emmanouilidis, et al., 2021; Lin, et al., 2021; Lin, et al., 2019; Seal, et al., 2021). FRQ represents a full-length, nearly 1000-residue IDP that can be investigated with respect to post-translational modification, interaction partners and phase separation. As anticipated from its sequence (Hurley, et al., 2013; Pelham, et al., 2020), this direct characterization of structural properties shows that FRQ is indeed a highly dynamic protein, but even in isolation, it is not completely disordered. The SAXS and ESR data indicate that FRQ has compact features that localize to an ordered core interspersed with dynamic regions. Notably the SAXS and DEER data suggest a degree of disorder that is substantially greater than that expected of a molten globule (Kataoka, et al., 1997), but far from that of a completely denatured protein (Kikhney, et al., 2015; Martin, Erik W., et al., 2021). Although FRQ lacks the well-defined PAS domains of the analogous PER proteins (Pelham, et al., 2020), it also maintains a structured core to anchor its more dynamic elements. The regions that bind CK1 and FRQ are the most ordered, and they organize the two partners beside each other; in fact, they are in remarkably close proximity, as judged by spin-labels in their ATP binding pockets or in their binding domains. FRH mediates contacts to the WCC (Conrad, et al., 2016), whereas CK1 phosphorylates the WCC to close the TTFL (Cha, et al., 2014; Larrondo, et al., 2015; Liu, X., et al., 2019; Wang, et al., 2019); hence, CK1 targeting to the WCC likely depends on FRQ associating FRH with CK1 (Wang, et al., 2019).

Previous proteolytic sensitivity studies have suggested that increased phosphorylation across the day transitions FRQ from a “closed” to a more “open” state (Liu, X., et al., 2019; Pelham, et al., 2020; Querfurth, et al., 2011). Our data paint a more nuanced picture. The SAXS data reveal that p-FRQ and p-FFC are more expansive and show greater flexibility than do np-FRQ and np-FFC, respectively. Furthermore, np-FRQ has a greater tendency to self-associate. In the cell, completely unphosphorylated FRQ is likely short-lived, if present at all (Baker, et al., 2009). Phosphorylation may stabilize FRQ against aggregation, yet also provides for enough conformational heterogeneity to poise FRQ near a degradation threshold (Larrondo, et al., 2015). This idea is supported by the cw-ESR data (Fig. 3), which shows that phosphorylation causes non-uniform local structural changes with some regions increasing, while others decreasing, in flexibility. The DEER data also demonstrates that binding to FRH and CK1 increases compactness of p-FRQ, particularly in the central-core that is responsible for binding the partners. In contrast, the FRQ C-terminus gains considerable flexibility in the FFC. FRQ has been suggested to assume a dimeric state, because of a predicted coiled-coil region and formation of mixed dimers formed in cellular pull-down assays with differentially tagged FRQ subunits (Cheng, et al., 2001). Np-FRQ does indeed dimerize when isolated, whereas p-FRQ is primarily monomeric alone. Furthermore, SAXS data indicates there is one copy of FRQ in both the reconstituted np-FFC and p-FFC. Hence phosphorylation has the potential to regulate the FRQ oligomeric state, which may in turn be linked to FFC assembly. Despite being a monomer in isolation, the pull-down data (Cheng, et al., 2001) indicate that association of p-FRQ may be mediated by other cellular factors, including LLPS.

FRQ has physicochemical features embedded within its sequence that give rise to LLPS and the FRQ clock analogs exhibit similar characteristics (Fig. S13-S15). FRQ and negative-arm proteins in general belong to the “flexible disorder” class of IDPs (Shapiro, et al., 2021), wherein amino acid patterns that generate intrinsic disorder are conserved, but amino acid sequence per se, is not (Bellay, et al., 2011). Moreover, although these proteins all clearly contain regions of

low-complexity sequence (Hurley, et al., 2013 [↗](#); Marzoll, et al., 2022 [↗](#); Pelham, et al., 2020 [↗](#)), their conserved features include not only their intrinsic disorder, but also their propensity for LLPS (**Fig. 6A** [↗](#), **Fig. S13** [↗](#)). These two properties, although related, are not directly correlated (Bolognesi, et al., 2016 [↗](#); Martin, E. W., et al., 2020; Szabó, et al., 2022 [↗](#); Vernon, et al., 2018 [↗](#)). In particular, a preponderance of aromatic and non-aromatic π - π planar contacts and charge-dense or charge-repeat regions drive LLPS (Szabó, et al., 2022 [↗](#); Vernon, et al., 2018 [↗](#)) and these features are consistently found in clock repressors, particularly FRQ. The similar amino acid compositions of these proteins may then potentially give rise to their shared function. Moreover, extensive phosphorylation may alter LLPS propensity, because it increases extended states, solvation of the peptide backbone, charge density and intermolecular hydrogen bonding capacity (Bolognesi, et al., 2016 [↗](#); Szabó, et al., 2022 [↗](#); Vernon, et al., 2018 [↗](#)). For FRQ, the sequence predictions are borne out experimentally, with p-FRQ displaying a different phase behavior relative to np-FRQ. In addition to being more soluble, dynamic and less aggregated than np-FRQ, p-FRQ also phase separates at lower temperatures and under lower salt conditions relative to np-FRQ. Increased salt disfavors p-FRQ phase separation, which when compared to np-FRQ, is perhaps consistent with more electrostatically driven de-mixing due to the high degree of phosphorylation.

Live cell imaging of fluorescently-tagged FRQ proteins is consistent with FRQ phase separation in *N. crassa* nuclei. The concentrated foci observed at specific positions in the nuclei indicate condensate behavior and are highly dynamic in size and shape, similar to that observed for other phase separating proteins (Bartholomai, et al., 2022 [↗](#); Caragliano, et al., 2022 [↗](#); Gonzalez, A., et al., 2021; Tatavosian, et al., 2019 [↗](#); Xiao, et al., 2021 [↗](#)). While ongoing experiments are exploring more deeply the spatiotemporal dynamics of FRQ condensates in nuclei, the small size of fungal nuclei as well as their rapid movement with cytoplasmic bulk flow through the hyphal syncytium makes these experiments difficult. Of particular interest is drawing comparisons between FRQ and the *Drosophila* Period protein, which has been observed in similar foci that change in size and subnuclear localization throughout the circadian cycle (Meyer, et al., 2006 [↗](#); Xiao, et al., 2021 [↗](#)).

Phosphorylation is known to modulate LLPS in other systems, but the effects are variable. For example, phosphorylation of the transcriptional silencing factor HP1 α also enhances LLPS, whereas only the non-phosphorylated form of *Xenopus* CPEB4 involved in translation of mRNA poly-A tails undergoes LLPS (Larson, et al., 2017 [↗](#); Seal, et al., 2021 [↗](#)). FRQ, especially p-FRQ, may achieve its inhibitory function by sequestering the FFC and WCC into membraneless organelles. The large, disordered domains and capacity for multivalency of the WC proteins (Pelham, et al., 2020 [↗](#)) may also facilitate their partitioning into such condensates. In the fly clock, PER and the WCC analog CLK produce liquid-like foci in the nuclei that are most prevalent during the repressive phase of the circadian cycle (Xiao, et al., 2021 [↗](#)). These large foci, which are few and localized to the nuclear periphery, may serve to organize clock-controlled genes during repression (Xiao, et al., 2021 [↗](#)). LLPS may also influence the nuclear transport of FRQ by affecting interactions with the nuclear import machinery as has been observed with proteins such as TD43 and FUS (Gasset-Rosa, et al., 2019 [↗](#); Gonzalez, Abner, et al., 2021). Conditions favoring LLPS reduce the ability of FRQ to act as a substrate for CK1 (**Fig. 7B** [↗](#)), and this reactivity correlates with reduced dynamics of FRQ in the phase separated state (**Fig. 7D** [↗](#)). Domains and disordered regions from other spin-labeled proteins show similar reduced dynamics when induced to phase separate and monitored by cw-ESR spectroscopy (Babinchak, et al., 2019 [↗](#); Emmanouilidis, et al., 2021 [↗](#); Lin, et al., 2021 [↗](#); Seal, et al., 2021 [↗](#)), although the temperature and salt dependencies of the ordering behavior can vary considerably, with some species showing little change upon LLPS (Lin, et al., 2019 [↗](#)).

In the cell, the impact of LLPS on CK1 activity may modulate the timing of FRQ phosphorylation, which is closely linked to clock period (Larrondo, et al., 2015 [↗](#); Marzoll, et al., 2022 [↗](#)). Moreover, and collectively, the data from **Fig. 6** [↗](#) and **7** [↗](#) show that as temperature increases, LLPS-like behavior increases, and as LLPS increases, the ability of CK1 to phosphorylate FRQ decreases. The

rate of FRQ phosphorylation dictates period length (Larrondo, et al., 2015 [↗](#)). Therefore, this negative feedback of temperature on period mediated by LLPS formation could provide a mechanism for circadian temperature compensation of period length (**Fig. S19** [↗](#)). Sequestration of CK1 (and FRH) into condensates may also render these factors unavailable to target other cellular substrates. Phosphorylation alters FRQ interactions throughout the circadian cycle, not unlike the workings of the cyanobacterial clock, which relies upon a timed phospho-code to regulate enzymatic components (Rust, et al., 2007 [↗](#); Wang, et al., 2019 [↗](#)). We demonstrate that FRQ likewise responds to phosphorylation with non-uniform conformational changes across its sequence and an increased propensity for LLPS. It is the latter property that appears conserved among negative-arm repressor proteins in eukaryotic clocks and may thus be a critical function of their intrinsic disorder and extensive post-translational modifications.

Materials and Methods

FRQ was obtained by recombinant expression in *E. coli* and purification under denaturing conditions followed by removal of denaturant. FRH and CK1 were produced using more standard protocols. SEC-MALS was performed with Wyatt Technologies DAWN HELEOS-II multi-angle light scattering detector and SEC-SAXS was carried out at the Cornell Synchrotron G1 station. Protein labeling for ESR and microscopy experiments made use of enzymatic labeling and reactive cysteine residue probes. Cw-ESR spectroscopy was carried out on a Bruker E500 ESR spectrometer operating at X-band, whereas DEER spectroscopy utilized a four-pulse protocol on a Bruker E580 spectrometer operating at Q band. Turbidity was as measured by scattering at 300 – 600 nm under different buffer conditions and a Peltier-driven temperature ramp. Microscopy was carried out on a Zeiss Axio Imager Upright Trinocular Fluorescence Microscope at 100x magnification. Phosphorylation assays involved reaction with [γ - 32 P] ATP, protein band resolution by SDS-PAGE and phosphorimage analysis. Mass-spectrometry and phospho-peptide detection was carried out on tryptic peptides analyzed by MS/MS with a CID ion-trap. *Neurospora crassa* strain generation targeted the endogenous *frq* locus with DNA encoding *N. crassa* codon-optimized mNeonGreen appended to the C-terminus of the *frq* ORF. For live-cell imaging a section near the edge of a growing mycelium was removed and inverted on a chambered cover glass. Images were collected with 488 nm laser excitation using a laser scanning confocal microscope equipped with a spinning disk scan head and a CMOS camera. Detailed descriptions of the protocols can be found in the *Supplementary Materials and Methods*.

Author contributions

D.T., N.M., B.M.B., J.C.D., A.B. and B.R.C. designed research; D.T., N.M., B.M.B., and S.C. performed research; D.T., N.M., B.M.B., S.C., J.C.D., A.B. and B.R.C. analyzed data; D.T. and B.R.C. wrote the paper with contributions and editing from all authors.

The authors declare no competing interests.

Data Availability

The mass spectrometry FRQ phosphorylation data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD037938. All other data are included in the article and/or supporting Information.

We thank the Cornell High Energy Synchrotron Source (CHESS) and the National Biomedical Center for Advanced ESR Technologies (ACERT) for access to data collection facilities. This work was supported by grants from the National Institutes of Health: R35GM122535 to BRC, R35GM118021 to JCD; R35GM138097 to AB, 1F31GM143890 to NM, and T32-008704 to BMB and from The Pew Charitable Trusts to AB. ACERT is supported by P41GM103521 and R24GM146107. CHESS is supported by NSF award DMR-1332208 and NIH/NIGMS award P30GM103485.

Supplementary Information

Materials and Methods

Protein Expression and Purification

Full length L-FRQ (Uniprot ID:P19970), CK1 (CK1a; Uniprot ID:V5IR38) and FRH γ N (residues 100–1106) (Uniprot ID:Q1K502) were sub-cloned into a pET28a vector (Novagen) bearing an N-terminal His₆-tag and kanamycin resistance. Note: L-FRQ contained the S513A substitution to improve protein stability as noted in (Querfurth, C., et al., 2007). The plasmids encoding the proteins were transformed into BL21 (DE3) *E. coli* cells, grown in LB media containing 50 $\mu\text{g ml}^{-1}$ kanamycin to an optical density (OD₆₀₀) of ~0.6–0.8 at 37 °C and protein expression was induced with 1 mM IPTG at 17 °C or room temperature. Cells were harvested after 12–18 h by centrifugation at 5,000 rpm and 4 °C. FRH and CK1 were purified by standard Ni-NTA chromatography followed by ion-exchange and size-exclusion chromatography. Briefly, cells were resuspended in lysis buffer (50 mM Tris pH 8.0, 500 mM NaCl, 10% glycerol, 1 mM MgCl₂, 5 mM imidazole) and sonicated on ice. The lysate was clarified by centrifugation at 20,000 rpm for 1 hr. The clarified lysate was applied to a gravity Ni-NTA column and washed with 5 column volumes (CVs) of wash buffer (50 mM Tris pH 8.0, 500 mM NaCl, 10% glycerol, 1 mM MgCl₂, 20 mM imidazole.) The protein was eluted using the elution buffer (50 mM Tris pH 8.0, 500 mM NaCl, 10% glycerol, 1 mM MgCl₂, 500 mM imidazole). The protein was then diluted into IEX (ion-exchange) bufferA (50 mM HEPES pH 7.5, 10% glycerol), applied to an S cation exchange column (GE Healthcare) and eluted over a 20 CV gradient against IEX bufferB (50 mM HEPES pH 7.5, 1 M NaCl, 10% glycerol). Fractions containing the protein were pooled, concentrated, and run over an S200 26/60 gel-filtration column pre-equilibrated in gel-filtration buffer (GFB; 50 mM HEPES, 200 mM NaCl, 10% glycerol.) L-FRQ was purified under denaturing conditions. Cells were resuspended in lysis buffer (50 mM Na₂PO₄ pH 7.42, 300 mM NaCl, 5 mM imidazole, 4 M guanidinium hydrochloride (GuHCl)) and sonicated on ice. The lysate was clarified by centrifugation at 20,000 rpm for 1 hr. The clarified lysate was applied to a gravity Ni-NTA column and washed with 3 CV of lysis buffer. The protein was eluted using the elution buffer (50 mM Na₂PO₄ pH 7.42, 300 mM NaCl, 500 mM imidazole, 4 M GuHCl). The protein was concentrated and applied to a S200 26/60 gel-filtration column pre-equilibrated in 50 mM Na₂PO₄ pH 7.42, 300 mM NaCl, 5mM Imidazole, 4 M GuHCl. Fractions containing the protein were pooled and concentrated. The concentrated protein was flash-frozen and stored at -80 °C. Prior to experiments, the purified L-FRQ was exchanged into a non-denaturing buffer (50 mM Na₂PO₄ pH 7.42, 300 mM NaCl, 200 mM arginine, 10% glycerol). For FFC preparations, GFB (50 mM HEPES pH 7.42, 200 mM NaCl, 10% glycerol) was used.

Sec mals

Protein samples between 50–75 μM were injected onto a Superose-6 (GE Lifesciences), equilibrated with 50 mM Na₂PO₄ pH 7.42, 300 mM NaCl, 200 mM arginine, 10% glycerol. The gel filtration column was coupled to a static 18-angle light scattering detector (DAWN HELEOS-II) and a refractive index detector (Optilab T-rEX) (Wyatt Technology, Goleta, CA). Data were collected every

second at a flow rate of 0.5 mL/min and analysis was carried out using ASTRA VI, yielding the molar mass and mass distribution (polydispersity) of the sample. The monomeric fraction of BSA (Sigma, St. Louis, MO) was used to standardize the light scattering detectors and optimize data quality control.

Sec saxs

X-ray scattering experiments were performed at the Cornell High Energy Synchrotron Source (CHESS) G1 station. Size-exclusion chromatography-coupled SAXS (SEC-SAXS) experiments were performed using a Superdex 200 5/150 GL or Superose-6 5/150 GL (3 mL) column operated by a GE AKTA Purifier at 4 °C with the elution flowing directly into an in-vacuum X-ray sample cell. Samples were centrifuged at 14,000×*g* for 10 min at 4 °C before loading onto a column pre-equilibrated in a matched buffer (50 mM Na₂PO₄ pH 7.42, 300 mM NaCl, 200 mM arginine, 10% glycerol). Samples were eluted at flow rates of 0.05–0.1 mL min^{−1}. For each sample, 2-s exposures were collected throughout elution until the elution profile had returned to buffer baseline, and scattering profiles of the elution buffer were averaged to produce a background-subtracted SEC-SAXS dataset. The scattering data was processed in RAW (Hopkins, et al., 2017 [DOI](#)) and regions with homogenous *R_G* values were chosen for the Guinier and Kratky analyses.

Protein Labeling for ESR Spectroscopy and Microscopy Experiments

Sort/AEP1-tagged FRQ was cloned into a pET28a (Novagen) vector as described in (Chandrasekaran, et al., 2021 [DOI](#)). The production of the SortaseA and AEP1 enzymes and their associated peptides (GGGGC for SortaseA and CNGL for AEP1) is also described in (Chandrasekaran, et al., 2021 [DOI](#)) and (Nguyen, et al., 2015 [DOI](#)), respectively. To enzymatically label FRQ with either SortaseA or AEP1, 1:1 protein and SortaseA/AEP1 as well as 2x excess peptide were incubated overnight to allow the reaction to reach completion. The following day, the reaction was run on a Superpose 6 10/300 analytical SEC column to remove unreacted peptide and enzyme, and the protein containing fractions were pooled and concentrated. To prepare the ADP-β-S-SL samples, ADP-β-S-SL was synthesized as previously reported in (Muok, et al., 2018 [DOI](#)). 100 μL of buffer containing 100 μM of ADP-β-S-SL and 100 μM FRH or CK1 were incubated overnight and then washed with GFB (50 mM HEPES pH 7.42, 200 mM NaCl, 10% glycerol) through 5x dilution followed by re-concentration to remove any unbound ADP-β-S-SL. 30 μL of each sample were used for data collection. For MTSL labeling, the protein was incubated with 5x molar excess of MTSL overnight to allow the reaction to reach completion. The following day, the reaction was run on HiTrap Desalting column (GE Life Sciences) to remove any excess spin-label.

To prepare fluorophore labeled FRH and CK1, the purified proteins were incubated with 2x molar excess Cy5-maleimide dye (GE Life Sciences) overnight to label any accessible cysteines on the target proteins. The excess dye was removed by running the reaction over a HiTrap desalting column equilibrated with GFB (50 mM HEPES pH 7.42, 200 mM NaCl, 10% glycerol).

ESR Spectroscopy

The cw-ESR spectra were obtained on a Bruker E500 ESR spectrometer operating at X-band (~9.4 GHz). The ESR measurements were carried out at room temperature with a modulation amplitude of 0.5–2 G and modulation frequency of 100 kHz. All spectra were digitized to 1024 points with a sweep width of ~100 G and an average of 4–16 scans. The *Q* values were calculated using the display distance function on the Bruker Xepir 2.6b software.

For 4-pulse-DEER, the samples were exchanged into deuterated buffer containing 30% d₈-glycerol, checked by cwESR and plunge frozen into liq. N₂. The pulse ESR measurements were carried out at Q-band (~34 GHz) on a Bruker E580 spectrometer equipped with a 10 W solid state amplifier (150W equivalent TWTA) and an arbitrary waveform generator (AWG). DEER was carried out using four pulses (π/2-τ₁-π-τ₁-π-pump-τ₂-π-τ₂-echo) with 16-step phase cycling at 60 K. The pump

and probe pulses were separated by 56 MHz (~20 G). The DEER data was background subtracted with an exponential background and the distance reconstruction was carried out using denoising and the SVD method (Srivastava, Madhur, et al., 2017; Srivastava, M., et al., 2017). Uncertainty in the distance distributions were calculated using the method described by Srivastava and Freed (Srivastava, M., et al., 2019)

Turbidity Assays

Cysteine-less (Cys-less) FRQ was created in order to minimize disulfide cross-linking at the high protein concentrations needed for *in vitro* phase separation experiments. For this, all four of the native cysteine residues were substituted to serine using standard molecular biology techniques. For the phase separation experiments, this cys-less FRQ variant was purified and concentrated for dialysis into phase separation buffer. Cys-less FRQ was dialyzed overnight into 25 mM Tris pH 8.0, 150 mM NaCl or 25 mM Tris pH 8.0, 500 mM NaCl for p-FRQ and np-FRQ, respectively, at 4 °C. Phase separation was induced by dialyzing in either low or high salt and increasing temperature. The protein solution can be observed as a clear solution before dialysis and then turbid after dialysis indicating LLPS has occurred. Microscopy (See *DIC and Fluorescence Microscopy*) confirmed the turbid solution is due to LLPS and not aggregation. Turbidity as measured by absorbance at 300 – 600 nm was then recorded by an Agilent Cary 3500 UV-Vis Multicell Peltier using a temperature ramp rate of 5 °C/min increasing from 10 °C to 80 °C. A final protein concentration of 5-15 µM was used for all turbidity assays.

DIC and Fluorescence Microscopy

Phase separation was induced via overnight dialysis into 25 mM Tris pH 8.0, 150 mM NaCl or 25 mM Tris (pH 8.0), 500 mM NaCl for p-FRQ and np-FRQ, respectively. The final protein concentration used for microscopy were 5-15 µM. Images were taken on the Zeiss Axio Imager Z1 Upright Trinocular Fluorescence Microscope at 100x magnification. To test the interaction of FRQ with FRH and CK1, the phase separated FRQ was spiked with equimolar Cy5-labeled FRH or Cy5-labeled CK1 for final concentrations of 5 µM.

In vitro Phosphorylation Assay

22 µL samples containing 10-15 µM LFRQ were incubated in the presence or absence of 2 µM CK1 and 2 µM FRH in 50 mM Na₂HPO₄ pH 7.5, 300 mM NaCl, 200 mM arginine, 10 mM MgCl₂. For the LLPS assay, FRQ was dialyzed into a buffer containing 50 mM Na₂HPO₄ pH 7.5, 300 mM NaCl and 10 mM MgCl₂. 2 µL of 2.5 mM cold ATP and 0.1 µM of [γ-32P] ATP (3000 Ci/mmol, PerkinElmer) was added to produce a total volume of ~25 µL. After ~1h of incubation, the reaction was quenched with 25 µL of 4x SDS with 10 mM EDTA pH 8.0 and then subjected to gel electrophoresis on a 4–20% gradient Tris-glycine gel. The gel was dried with a GelAir dryer (Bio-Rad) and placed in an imaging cassette for at least 16 h and then imaged with a phosphorimager (GE, Typhoon FLA 7000). Band intensities were quantified using the Fiji software.

Mass Spectrometry and Phospho-Peptide Analysis

Phosphorylated FRQ was run on a 4-20% tris-glycine SDS-PAGE gel and the corresponding band was excised and trypsin digested. The digests were reconstituted in 0.5% formic acid (FA) for nanoLC-ESI-MS/MS analysis. The analysis was carried out using an Orbitrap Fusion™ Tribrid™ (Thermo-Fisher Scientific, San Jose, CA) mass spectrometer equipped with a nanospray Flex Ion Source, and coupled with a Dionex UltiMate 3000 RSLCnano system (Thermo, Sunnyvale, CA). The peptide samples (5 µL) were injected onto a PepMap C-18 RP nano trapping column (5 µm, 100 µm i.d x 20 mm) and then separated on a PepMap C-18 RP nano column (2 µm, 75 µm x 25 cm) at 35 °C. The Orbitrap Fusion was operated in positive ion mode with spray voltage set at 1.5 kV and source temperature at 275 °C. External calibration for FT, IT and quadrupole mass analyzers was performed. In data-dependent acquisition (DDA) analysis, the instrument was operated using FT

mass analyzer in MS scan to select precursor ions followed by 3 second “Top Speed” data-dependent CID ion trap MS/MS scans at 1.6 m/z quadrupole isolation for precursor peptides with multiple charged ions above a threshold ion count of 10,000 and normalized collision energy of 30%. MS survey scans at a resolving power of 120,000 (fwhm at m/z 200), for the mass range of m/z 375–1600. Dynamic exclusion parameters were set at 35 s of exclusion duration with ± 10 ppm exclusion mass width. All data were acquired under Xcalibur 4.4 operation software (Thermo-Fisher Scientific). The DDA raw files with MS and MS/MS were subjected to database searches using Proteome Discoverer (PD) 2.4 software (Thermo Fisher Scientific, Bremen, Germany) with the Sequest HT algorithm. The database search was conducted against a *Neurospora Crassa* database and *E. coli* database with adding LFRQ sequence into the database. The peptide precursor tolerance was set to 10 ppm and fragment ion tolerance was set to 0.6 Da with 2 miss-cleavage allowed for either trypsin or chymotrypsin. Variable modification of methionine oxidation, deamidation of asparagines/glutamine, phosphorylation of serine/threonine/tyrosine, acetylation, M-loss and M-loss+acetylation on protein N-terminus and fixed modification of cysteine carbamidomethylation, were set for the database search. Only high confidence peptides defined by Sequest HT with a 1% FDR by Percolator were considered for confident peptide identification. All the mass spectra of phosphorylated peptides were double checked by manually inspection.

Integrated Bioinformatic Analyses and Computational Biology

All canonical protein and UniRef50 sequences for FRQ, FRH and CK1, and SWISS-Prot obtained proteome sequences of *Neurospora crassa*, *Drosophila melanogaster* and *Homo sapiens* were downloaded from the UniProt database (Bateman, et al., 2021 [DOI](#)). A combination of (a) in-house written Python scripts to allow us to analyze large datasets, (b) localCIDER (Holehouse, et al., 2017 [DOI](#)), (c) IUPred2A (Mészáros, et al., 2018 [DOI](#)), (d) Agadir (Muñoz, et al., 1994), (e) AlphaFold 2.0 (Jumper, et al., 2021 [DOI](#)), (f) PScore (Vernon, et al., 2018 [DOI](#)) and (g) catGRANULE (Bolognesi, et al., 2016 [DOI](#)) were used to perform integrated bioinformatics analyses and computational biological studies of our proteins of interest. Unless mentioned, all programs were used by applying the default parameters. All scripts used to generate the data in this manuscript are provided as SI Datasets.

Neurospora crassa Strain Generation

Construction of the *frq*[mNeonGreen], *son-1*[mApple] strain used for imaging was previously described (Bartholomai, 2022 [DOI](#)). Briefly, a plasmid was constructed to target the endogenous *frq* locus with DNA encoding *N. crassa* codon-optimized mNeonGreen appended to the C-terminus of the *frq* ORF with a flexible linker. A gene encoding hygromycin B phosphotransferase (HPH) was introduced downstream from the 3' UTR of *frq* for selection. A plasmid was constructed containing the *N. crassa son-1* ORF with mApple appended to the C-terminus with a flexible linker for insertion at the *csr-1* locus. Transformation cassettes were amplified by PCR using NEB Q5® High-Fidelity 2X Master Mix (New England Bio Labs Cat. # M0492) and integrated into the genome by homologous recombination after electroporation of conidiophores. A homokaryotic strain containing the fluorescent fusion proteins was used for imaging.

Neurospora crassa Growth

N. crassa homokaryons containing *frq*[mNeonGreen] and *csr-1::son-1*[mApple], were grown overnight on 2% agar gel pads made with Vogel's minimal medium (Vogel, 1956 [DOI](#)) in constant light at 25 °C. Before imaging, a section near the edge of the growing mycelium was removed and inverted on a chambered cover glass (Thermo Scientific, Cat. # 155360).

Live Cell Image Acquisition and Processing

Images were acquired using a Zeiss AxioObserver laser scanning confocal microscope with a 100X/1.46 NA Oil immersion Plan-Apochromat objective. 561 nm and 488 nm lasers were used at 2% power with 2X averaging and a gain of 800 for detection of fluorescent signals by GaAsP PMTs. Images were processed for display using FIJI (Schindelin, et al., 2012 [DOI](#)). The cropped images displaying a single nucleus were enlarged 10X using bicubic interpolation and contrasted to the minimum and maximum gray level values. A multi-color lookup table was applied to the FRQ[mNeonGreen] channel to better observe heterogeneity of the signal, and the SON-1[mApple] channel was pseudo-colored cyan to better contrast with the lookup table applied to the FRQ[mNeonGreen] channel. The surface plot of FRQ[mNeonGreen] was generated from the raw image. Movie S1, and the image in **Figure 6E** [DOI](#) were acquired using a Nikon Ti-E based system, equipped with an Andor W1 spinning disk scan head, an Andor Zyla sCMOS camera, a 488 nm laser, and Chroma bandpass filters. **Figure 6E** [DOI](#) is a still image from a single focal plane from a z-stack that was deconvolved using Nikon Elements and was acquired using a 100X/NA1.45 plan apo oil immersion objective. **Figure 6E** [DOI](#) and Movie S1 were acquired using a 60X/NA1.45 plan apo oil immersion objective. The 1-minute movie was acquired with a 300 ms exposure per frame. Image processing for display was carried out using FIJI.

Results

	R_g [Guinier] (Å)	R_g [P(r)] (Å)	Dmax (Å)	MW [V_c] (kDa)
p-FRQ	123 ± 7	142 ± 5	530 ± 60	816
np-FRQ	100 ± 2	116 ± 4	446 ± 40	680
p-FFC	101 ± 2	136 ± 5	510 ± 20	314
np-FFC	80 ± 2	109 ± 7	480 ± 10	286

MW calculations for p-FRQ and np-FRQ are inaccurate because V_c (Volume of Correlation) plots (Integrated area of $I(q)q$ vs q ; $q = 2\pi/d$) did not converge for these species. Expected MW for LFRQ is 110 kDa. Expected FFC MW for the 1:1:1 stoichiometry of FRQ:FRH:CK1 is 273 kDa. Errors in Dmax were estimated from the range for which P(r) tailed to zero in the high distance range without constraint.

Table S1

SAXS parameters for p-FRQ, np-FRQ and their reconstitution into the FFC

Site	Domain	Binding Partner at this region	Reference
370C	N/A		
490C	FCD2	CK1	(Querfurth, Christina, et al., 2011)
658C	N/A		
746C	FFD	FRH	(Cheng, et al., 2005)

Table S2

FRQ Cysteine residue locations and context

Binding Partner	Region	Cysteine residue(s) present in this region	Reference
FRH	695-778	746C	(Cheng, et al., 2005)
CK1	319-326; 488-495	490C	(Querfurth, Christina, et al., 2011)
WC2		490C,658C,746C	(Guo, et al., 2010)

Table S3

FRQ regions that bind clock proteins

Site	np-FRQ (G)	p-FRQ (G)
NT	1.75	1.87
490C	1.97	2.47
658C	1.97	2.01
746C	1.76	1.82
CT	2.30	1.70

Table S4

o values from the cw-ESR central peak for FRQ variants of **Fig. 3C**

	370	490	658	746
370		0.99, 0.99	0.96, 0.99	NA, 0.99
490			0.94, 0.99	0.99, 0.99
658				0.99, 0.95
746				

Note: There are two values reported in each cell – the first value corresponds to the R-square value for FRQ only distance while the second value is for the same variant in the FFC complex (i.e., when the FRQ is bound to CK1a and FRH).

Table S5

R-square values for the quality of the two component gaussian fits to the distance distributions.

	5 μ M	10 μ M	15 μ M
Mean Initial Baseline Absorbance	0.02	0.08	0.13
Temperature of Initiation ($^{\circ}$ C)	56	23	21
Cloud Point (Inflection Point) ($^{\circ}$ C)	68	58	57
Absorbance at Inflection Point	0.07	0.21	0.33
Slope at Inflection Point	0.01	0.02	0.03
Mean Final Baseline Absorbance	0.14	0.47	0.64

Note: Initiation temperature was determined as baseline + 5X standard deviation of the baseline.

Table S6

Calculated values from the UV-Vis turbidity assays of np-FRQ shown in [Figure 6C](#).

	5 μ M	10 μ M	15 μ M
Mean Initial Baseline Absorbance	0.09	0.21	0.46
Temperature of Initiation ($^{\circ}$ C)	45	42	24
Cloud Point (Inflection Point) ($^{\circ}$ C)	-	-	-
Absorbance at Inflection Point	-	-	-
Slope	0.00	0.01	0.02
Mean Final Baseline Absorbance	-	-	-

Note: Initiation temperature was determined as baseline +5X standard deviation of the baseline. Also, because p-FRQ continues to undergo LLPS within the temperature range of the experiment, we cannot calculate a cloud point.

Table S7

Calculated values from the UV-Vis turbidity assays of p-FRQ shown in [Figure 6C](#).




Figure S1

(A) AlphaFold model of LFRQ colored by pLDDT score. (B) Predicted error alignment plot that shows the uncertainty in the relative positions of two residues in the structure. (Image source: <https://www.uniprot.org/uniprotkb/P19970/entry#structure>)




Figure S2

Domain map of FRQ with phospho-sites (as identified by mass-spec) highlighted. The sites in red were also identified in (Baker, et al., 2009; Tang, et al., 2009).




Figure S3

Pair-wise distance distributions calculated from SAXS data for p-FRQ (red), np-FRQ (dark blue), p-FFC (magenta), and np-FFC (light blue).




Figure S4

Enzymatic labeling of FRQ at its termini. X-band (9.8 GHz) RT cw-ESR measurements for the spin-labeled peptide by itself and when it has been grafted onto either the N or C-terminus of FRQ.




Figure S5

Time domain traces before (Raw) and after (Data) background correction and trace fits (Fit) from DEER experiments of different double-cysteine FRQ variants alone or within the FFC (i.e., when FRQ was bound to FRH and CK1a). The spins were present on the residue number of FRQ that is highlighted for example '370-490' refers to FRQ spin-labeled at the 370C and 490C site and the DEER experiment probed the dipolar coupling of the spins at these sites. The last graphs show the time domain data for the control single cysteine DEER experiments with p-FRQ 490C and p-FRQ 746C.




Figure S5 (continued)




Figure S6

Baseline-corrected time domain fit traces from Q-band DEER experiments of different double-cysteine FRQ mutants alone (blue) and within the FFC (orange). Original data are shown in **Supplementary Figures S5** and **S9**.




Figure S7

Full autoradiograph of np-FRQ phosphorylation by CK1 and γ - ^{32}P -ATP from **Figure 4A** [inset](#). Four technical replicates are shown for each np-FRQ concentration.




Figure S8

Background-corrected time domain trace and error analyses from DEER experiments targeting the ATP-binding pockets of FRH and CK1. The DEER experiment probed the dipolar coupling of the spins in the ADP- β -S-SL molecule that was present in ATP-binding pocket of FRH or CK1.




Figure S9

Error analysis of the distance distributions produced from DEER experiments of different double-cysteine FRQ variants. The spins were present on the amino acid number of FRQ that is highlighted for example '370-490' refers to FRQ spin-labeled at the 370C and 490C site and the DEER experiment probed the dipolar coupling of the spins at these sites.




Figure S10

Error analysis of the distance distributions produced from DEER experiments of different double-cysteine FRQ variants within the FFC (i.e., when FRQ was bound to FRH and CK1). The spins were present on the amino acid number of FRQ that is highlighted for example '370-490' refers to FRQ spin-labeled at the 370C and 490C site and the DEER experiment probed the dipolar coupling of the spins at these sites.




Figure S11

(Left) The two component Gaussian fitting of the distance distribution ($P(r)$) obtained from SVD for FFC 370-490 (i.e., spins were placed on residues 370 and 490 on FRQ, and this protein was bound to CK1 and FRH). No constraints were placed on the width, position, or amplitude of either component and the best fit values were obtained by minimizing the difference between the fits and the SVD distance distributions. Similar fits were carried out for all the different variants. The quality of the fit is reported by the R-square values and ranged between 0.94-0.99 for all the samples tested. They are reported in Table S5. (Right) The proportion of the closer distance peak in the distance distributions of FRQ (blue) or FRQ+FRH+CK1 (FFC) (red) samples. The error bars reflect the uncertainty in estimating the population of each peak and were derived from uncertainties in the distance distributions shown in *SI Appendix*, Fig. S9 and S10.




Figure S12

FRQ, but not FRH or CK1, is highly disordered and predicted to undergo LLPS. Shown are the disorder and LLPS propensity scores of the members for the FFC complex, FRQ, FRH and CK1. Columns from left to right are disorder and phase separation predictions for FRQ, FRH and CK1, respectively. Disorder graphs (red) were computed using the IUPred2A program (Mészáros, et al., 2018) where values above a 0.5 threshold (dashed line) are considered disordered. LLPS propensity predictions were computed using “Pi-Pi” (blue) (Vernon, et al., 2018) and catGRANULE (light blue) (Bolognesi, et al., 2016). Dashed lines represent neutral propensity. Shown on the top right corner of each graph are the percent disorder or LLPS propensity scores of each protein.




Figure S13

FRQ and its functional homologs are predicted to undergo LLPS. Shown are the disorder and LLPS propensity scores for FRQ, dPER1 and hPER1. Data for the disorder graphs (red) were computed using the IUPred2A program where values above a 0.5 threshold (dashed line) are considered disordered. Pi-pi predictions (blue) were computed using (Vernon, et al., 2018), while catGRANULE predictions (light blue) were computed using catGRANULE server (Bolognesi, et al., 2016). Shown on the top-right corner of each graph are the percent disorder, PScore or catGRANULE score for each protein. Dashed lines represent neutral propensity. FRQ, dPER1 and hPER1 are all predicted to be highly disordered (86.77%, 58.84% and 63.07%, respectively) and undergo LLPS.




Figure S14

FRQ and its homologs, dPER1 and hPER1, have similarly disparate physicochemical properties compared to their proteomes. (A) Shown are the distribution of fraction of charged residues (FCR), net charge per residue (NCPR), and hydropathy along the sequence of FRQ, dPER1 and hPER1. (B) Sequence characteristics of FRQ and its homologs, dPER1 and hPER1. Various protein sequence parameters are calculated using localCIDER (Holehouse, et al., 2017) and displayed including fraction of charged residues (FCR), the degree of charge mixing (Kappa), net charge per residue (NCPR), and hydropathy (the hydrophobic and hydrophilic character of protein chain as defined by (Kyte, et al., 1982)) for the UniProtRef50 clusters of FRQ (purple), dPER1 (blue) and hPER1 (orange). For comparison are shown protein sequences of similar length from the SWISS-Prot reviewed proteomes of *Neurospora crassa* (light purple), *Drosophila melanogaster* (light blue) and *Homo sapiens* (light orange). Hydropathy is consistently low for the clock proteins and FRQ orthologs have a high degree of charge separation, i.e., tracks of negative and positive residues (high kappa). High kappa values are common for proteins that phase separate (Somjee, et al., 2020). (C) Classification of the charge properties of the Ref50 clusters of FRQ and its functional homologs, dPER1 and hPER1. Das-Pappu phase plots showing the fraction of negatively vs positively charged residues for the UniProtRef50 clusters of FRQ (purple), *Neurospora crassa* proteome control (light purple), dPER1 (blue), *Drosophila melanogaster* proteome control (light blue), hPER1 (orange) and *Homo sapiens* proteome control (light orange). Shown are the fraction of negative residues versus fraction of positive residues for each cluster of homologs and their controls (similar size proteins from their respective proteomes). Despite the lack of amino acid conservation between the functional homologs of FRQ, dPER1 and hPER1, the ratio of fraction negative to fraction positive residues are similar among the clock proteins. dPER1 and hPER1 have a similarly low value of both fraction positive and fraction negative compared to their respective proteomes, whereas the fraction positive and fraction negative residues are both higher in FRQ. Note: Ref50 is a clustering method utilized by UniProt and is defined in (Steinegger, et al., 2018). While this work was in preparation, conceptually similar analyses were carried out by Jankowski et al., 2022 (Jankowski, et al., 2022).




Figure S14 (continued)




Figure S15

FRQ and its functional homologs dPER1 and hPER1 have similar amino acid composition compared to their respective proteomes. FRQ and its functional homologs dPER1 and hPER1 have similar amino acid compositions, which differs substantially from that of their respective proteomes. The amino acid composition for FRQ, dPER1 and hPER1 and their respective homolog clusters and proteomes are shown. Generally, G, S, T and P residues are enriched whereas hydrophobic (A, I, L, V) residues are relatively depleted for FRQ, dPER1, hPER1 and their homologs compared to their proteomes. Unlike dPER1 and hPER1 and their homologs, FRQ and its homologs are also highly enriched in R and D.




Figure S16

Phosphorylation modulates the physicochemical properties of FRQ. Shown are the distribution of fraction of charged residues (FCR), net charge per residue (NCPR), and hydropathy along the sequence of np and p-FRQ (top panel). In the bottom panel, shown are the disorder and LLPS propensity scores for np and p-FRQ. Data for the disorder graphs (red) were computed using the IUPred2A program (Mészáros, et al., 2018) where values above a 0.5 threshold (dashed line) are considered disordered. Pi-pi predictions (blue) were computed using, while catGRANULE predictions (light blue) were computed using catGRANULE server (Bolognesi, et al., 2016). Shown on the top-right corner of each graph are the percent disorder, PScore or catGRANULE score for each protein. Dashed lines represent neutral propensity. Note: Glu(E) residues were employed as phospho-mimetics in all analysis.




Figure S17

Exploring the phase behavior of FRQ. (A) UV-Vis absorption spectrum of p-FRQ (in 150 mM NaCl, 25 mM Tris pH 8.0) and np-FRQ (in 500 mM NaCl, 25 mM Tris pH 8.0) as a function of temperature at increasing concentrations. Note: the shading around each curve represents the standard deviation from the mean absorbance from 2 technical replicates. 15 μ M p-FRQ had the greatest standard deviation as seen by the largest shaded area. (B) UV-Vis absorption spectrum of p-FRQ and np-FRQ in a non-LLPS promoting buffer as a function of temperature. Note: p-FRQ appears to undergo a small amount of LLPS even in non-LLPS buffer as seen by its relatively high absorbance relative to np-FRQ. (C) (Left) UV-Vis absorption spectrum of p-FRQ (in 150 mM NaCl, 25 mM Tris pH 8.0) and (Right) np-FRQ (in 500 mM NaCl, 25 mM Tris pH 8.0) as a function of temperature to check for reversibility of LLPS transition. (D) DIC micrograph of FRQ showing droplets fusing and docking as indicated within the yellow circles. (E) UV-Vis absorption spectrum of p-FRQ (Left) and np-FRQ (Right) as a function of temperature in either low salt (150 mM) or high salt (500 mM).




Figure S18

Full radiograph of np-FRQ phosphorylation by CK1 and γ - ^{32}P -ATP from **Figure 7**.





Figure S19

Schematic diagram summarizing a potential mechanism for LLPS mediated circadian clock regulation. By reducing CK1 activity, especially at higher temperatures, LLPS provides a potential temperature compensation mechanism for the clock.



Movie S1

Visualization of FRQ [mNeonGreen] in living *Neurospora*, displaying heterogeneous, punctate intranuclear localization. Images of a single focal plane were acquired in a time series with 300 ms between exposures.

References

1. Babinchak WM, Haider R, Dumm BK, Sarkar P, Surewicz K, Choi JK, Surewicz WK (2019) **The role of liquid-liquid phase separation in aggregation of the TDP-43 low-complexity domain** *Journal of Biological Chemistry* **294**:6306–6317
2. Baker CL, Kettenbach AN, Loros JJ, Gerber SA, Dunlap JC (2009) **Quantitative Proteomics Reveals a Dynamic Interactome and Phase-Specific Phosphorylation in the Neurospora Circadian Clock** *Molecular Cell* **34**:354–363
3. Bartholomai BM, Gladfelter AS, Loros J. J., Dunlap JC (2022) **PRD-2 mediates clock-regulated perinuclear localization of clock gene RNAs within the circadian cycle of Neurospora** *Proc Natl. Acad. Sci. USA* **119**
4. Bellay J, Han SJ, Michaut M, Kim T, Costanzo M, Andrews BJ, Boone C, Bader GD, et al. (2011) **Bringing order to protein disorder through comparative genomics and genetic interactions** *Genome Biology* **12**
5. Berliner LJ, Grunwald J, Hankovszky HO, Hideg K (1982) **A novel reversible thiol-specific spin label – papain active-site labeling and inhibition** *Analytical Biochemistry* **119**:450–455
6. Bolognesi B, Gotor NL, Dhar R, Cirillo D, Baldrighi M, Tartaglia GG, Lehner B, Gaetano Tartaglia G (2016) **A Concentration-Dependent Liquid Phase Separation Can Cause Toxicity upon Increased Protein Expression** *CellReports* **16**:1–10
7. Bonucci A, Murrall MG, Banci L, Pierattelli R (2020) **A combined NMR and EPR investigation on the effect of the disordered RGG regions in the structure and the activity of the RRM domain of FUS** *Scientific Reports* **10**:1–13
8. Caragliano E, Bonazza S, Frascaroli G, Tang JJ, Soh TK, Grunewald K, Bosse JB, Brune W (2022) **Human cytomegalovirus forms phase-separated compartments at viral genomes to facilitate viral replication** *Cell Reports* **38**
9. Cha J, Zhou M, Liu Y (2014) **Mechanism of the Neurospora Circadian Clock, a FREQUENCY-centric View** *Biochemistry* **54**:150–156
10. Chandrasekaran S, Schneps CM, Dunleavy R, Lin C, DeOliveira CC, Ganguly A, Crane BR (2021) **Tuning flavin environment to detect and control light-induced conformational switching in Drosophila cryptochrome** *Communications Biology* **2021 4**:1 4:1–12
11. Cheng P, He Q, He Q, Wang L, Liu Y (2005) **Regulation of the Neurospora circadian clock by an RNA helicase** *Genes and Development* **19**:234–241
12. Cheng P, Yang Y, Heintzen C, Liu Y (2001) **Coiled-coil domain-mediated FRQ-FRQ interaction is essential for its circadian clock function in Neurospora** *EMBO J.* **20**:101–108
13. Conrad KS, Hurley JM, Widom J, Ringelberg CS, Loros JJ, Dunlap JC, Crane BR (2016) **Structure of the frequency-interacting RNA helicase: a protein interaction hub for the circadian clock** *EMBO J* **35**:1707–1719

14. Crane BR, Young MW (2014) **Interactive Features of Proteins Composing Eukaryotic Circadian Clocks** *Annual Review of Biochemistry* **83**:191–219
15. Diernfellner A, Colot HV, Dintzis O, Loros JJ, Dunlap JC, Brunner M (2007) **Long and short isoforms of Neurospora clock protein FRQ support temperature-compensated circadian rhythms** *Febs Letters* **581**:5759–5764
16. Dignon GL, Best RB, Mittal J (2020) **Biomolecular Phase Separation: From Molecular Driving Forces to Macroscopic Properties** *Annual Review of Physical Chemistry* **71**:53–75
17. Dunlap JC, Loros JJ (2017) **Making Time: Conservation of Biological Clocks from Fungi to Animals** *Microbiology Spectrum* **5**
18. Dunlap JC, Loros JJ (2018) **Just-So Stories and Origin Myths: Phosphorylation and Structural Disorder in Circadian Clock Proteins** *Molecular Cell* **69**:165–168
19. Dunleavy R, Chandrasekaran S, Crane BR (2023) **Enzymatic Spin-Labeling of Protein N- and C-Termini for Electron Paramagnetic Resonance Spectroscopy** *Bioconjugate Chemistry* **34**:686–695
20. Durand D, Vivès C, Cannella D, Pérez J, Pebay-Peyroula E, Vachette P, Fieschi F (2010) **NADPH oxidase activator p67phox behaves in solution as a multidomain protein with semi-flexible linkers** *Journal of Structural Biology* **169**:45–53
21. Emmanouilidis L, Esteban-Hofer L, Damberger FF, de Vries T, Nguyen CKX, Ibáñez LF, Mergenthal S, Klotzsch E, et al. (2021) **NMR and EPR reveal a compaction of the RNA-binding protein FUS upon droplet formation** *Nature Chemical Biology* **17**:608–614
22. Franck JM, Chandrasekaran S, Dzikovski B, Dunnam CR, Freed JH (2015) **Focus: Two-dimensional electron-electron double resonance and molecular motions: The challenge of higher frequencies** *Journal of Chemical Physics* **142**
23. Freed JH (2005) **ESR and Molecular Dynamics, Biomedical EPR, Part B: Methodology, Instrumentation, and Dynamics** :239–268
24. Froehlich AC, Liu Y, Loros JJ, Dunlap JC (2002) **White collar-1, a circadian blue light photoreceptor, binding to the frequency promoter** *Science* **297**:815–819
25. Gasset-Rosa F, Lu S, Yu H, Chen C, Melamed Ze, Guo L, Shorter J, Da Cruz S, et al. (2019) **Cytoplasmic TDP-43 De-mixing Independent of Stress Granules Drives Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death** *Neuron* **102**:339–357
26. Gonzalez A, Mannen T, Cagatay T, Fujiwara A, Matsumura H, Niesman AB, Brautigam CA, Chook YM, et al. (2021) **Mechanism of karyopherin-beta 2 binding and nuclear import of ALS variants FUS(P525L) and FUS(R495X)** *Scientific Reports* **11**
27. Gonzalez A, Mannen T, Çağatay T, Fujiwara A, Matsumura H, Niesman AB, Brautigam CA, Chook YM, et al. (2021) **Mechanism of karyopherin-β2 binding and nuclear import of ALS variants FUS(P525L) and FUS(R495X)** *Scientific Reports* **2021 11:1** **11**:1–15
28. Guo J, Cheng P, Liu Y (2010) **Functional significance of FRH in regulating the phosphorylation and stability of Neurospora circadian clock protein FRQ** *J Biol. Chem* **285**:11508–11515

29. Hubbell WL, Cafiso DS, Altenbach C (2000) **Identifying conformational changes with site-directed spin labeling** *Nature Structural Biology* **7**:735–739
30. Hurley JM, Larrondo LF, Loros JJ, Dunlap JC (2013) **Conserved RNA Helicase FRH Acts Nonenzymatically to Support the Intrinsically Disordered Neurospora Clock Protein FRQ** *Molecular Cell* **52**:832–843
31. Hurley JM, Loros JJ, Dunlap JC (2016) **Circadian Oscillators: Around the Transcription-Translation Feedback Loop and on to Output** *Trends in Biochemical Sciences* **41**:834–846
32. Jankowski MS, Griffith D, Shastru DG, Pelham JF, Ginell GM, Thomas J, Karande P, Holehouse AS, et al. (2022) **The formation of a fuzzy complex in the negative arm regulates the robustness of the circadian clock** *bioRxiv* <https://doi.org/10.1101/2022.01.04.474980>
33. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, et al. (2021) **Highly accurate protein structure prediction with AlphaFold** *Nature* **2021** 596:583–589
34. Jung JH, Barbosa AD, Hutin S, Kumita JR, Gao M, Derwort D, Silva CS, Lai X, et al. (2020) **A prion-like domain in ELF3 functions as a thermosensor in Arabidopsis** *Nature* **585**:256–260
35. Kataoka M, Kuwajima K, Tokunaga F, Goto Y (1997) **Structural characterization of the molten globule of alpha-lactalbumin by solution X-ray scattering** *Protein Science* **6**:422–430
36. Kikhney AG, Svergun DI (2015) **A practical guide to small angle X-ray scattering (SAXS) of flexible and intrinsically disordered proteins** *Febs Letters* **589**:2570–2577
37. Larrondo LF, Olivares-Yañez C, Baker CL, Loros JJ, Dunlap JC (2015) **Decoupling circadian clock protein turnover from circadian period determination** *Science* **347**
38. Larson AG, Elnatan D, Keenen MM, Trnka MJ, Ohnston JBJ, Burlingame AL, Agard DA, Redding S, et al. (2017) **Liquid droplet formation by HP1 alpha suggests a role for phase separation in heterochromatin** *Nature* **547**:236–241
39. Lauinger L, Diernfellner A, Falk S, Brunner M (2014) **The RNA helicase FRH is an ATP-dependent regulator of CK1a in the circadian clock of Neurospora crassa** *Nature communications*
40. Lin YX, Fichou Y, Longhini AP, Llanes LC, Yin PY, Bazan GC, Kosik KS, Han S (2021) **Liquid-Liquid Phase Separation of Tau Driven by Hydrophobic Interaction Facilitates Fibrillization of Tau** *Journal of Molecular Biology* **433**
41. Lin YX, McCarty J, Rauch JN, Delaney KT, Kosik KS, Fredrickson GH, Shea JE, Han S (2019) **Narrow equilibrium window for complex coacervation of tau and RNA under cellular conditions** *Elife* **8**
42. Liu BQ, Poolman B, Boersma AJ (2017) **Ionic Strength Sensing in Living Cells** *Acs Chemical Biology* **12**:2510–2514
43. Liu X, Chen A, Caicedo-Casso A, Cui GF, Du MJ, He Q, Lim S, Kim HJ, et al. (2019) **FRQ-CK1 interaction determines the period of circadian rhythms in Neurospora** *Nature Communications* **10**

44. Liu Y, Garceau NY, Loros JJ, Dunlap JC (1997) **Thermally regulated translational control of FRQ mediates aspects of temperature responses in the *Neurospora* circadian clock** *Cell* **89**:477–486
45. Martin EW, Holehouse AS (2020) **Intrinsically disordered protein regions and phase separation: sequence determinants of assembly or lack thereof** *Emerging Topics in Life Sciences* **4**:307–329
46. Martin EW, Holehouse AS, Peran I, Farag M, Incicco JJ, Bremer A, Grace CR, Soranno A, et al. (2020) **Valence and patterning of aromatic residues determine the phase behavior of prion-like domains** *Science* **367**:694–+
47. Martin EW, Hopkins JB, Mittag T (2021) **Small-angle X-ray scattering experiments of monodisperse intrinsically disordered protein samples close to the solubility limit** *Methods in Enzymology* **646**:185–222
48. Marzoll D, Serrano FE, Shostak A, Schunke C, Diernfellner ACR (2022) **Casein kinase 1 and disordered clock proteins form functionally equivalent, phospho-based circadian modules in fungi and mammals** *Proc. Natl. Acad. Sci. USA* **119**
49. Meyer P, Saez L, Young MW (2006) **PER-TIM interactions in living *Drosophila* cells: An interval timer for the circadian clock** *Science* **311**:226–229
50. Muñoz V, Serrano L (1994) **Elucidating the folding problem of helical peptides using empirical parameters** *Nature Structural Biology* **1**:399–409
51. Muok AR, Chua TK, Le H, Crane BR (2018) **Nucleotide Spin Labeling for ESR Spectroscopy of ATP-Binding Proteins** *Applied Magnetic Resonance* **49**:1385–1395
52. Nguyen GKT, Cao Y, Wang W, Liu CF, Tam JP (2015) **Site-Specific N-Terminal Labeling of Peptides and Proteins using Butelase 1 and Thiodipeptide** *Angewandte Chemie-International Edition* **54**:15694–15698
53. Pedersen CP, Seiffert P, Brakti I, Bugge K (2020) **Production of intrinsically disordered proteins for biophysical studies: Tips and tricks** *Methods in Molecular Biology* **2141**:195–209
54. Pelham JF, Dunlap JC, Hurley JM (2020) **Intrinsic disorder is an essential characteristic of components in the conserved circadian circuit** *Cell Communication and Signaling* **18**
55. Pelham JF, Mosier AE, Altshuler SC, Kirchoff CL, Fall WB, Baik LS, Chiu JC, Hurley JM (2021) **Conformational Changes in the Negative Arm of the Circadian Clock Correlate with Dynamic Interactomes Involved in Post-transcriptionally Regulated Processes** *bioRxiv* <https://doi.org/10.1101/2021.11.20.469315>
56. Posey AE, Holehouse AS, Pappu RV (2018) **Phase Separation of Intrinsically Disordered Proteins** *Methods in enzymology* **611**:1–30
57. Querfurth C, Diernfellner ACR, Gin E, Malzahn E, Höfer T, Brunner M (2011) **Circadian Conformational Change of the *Neurospora* Clock Protein FREQUENCY Triggered by Clustered Hyperphosphorylation of a Basic Domain** *Molecular Cell* **43**:713–722
58. Rambo RP, Tainer JA (2011) **Characterizing Flexible and Intrinsically Unstructured Biological Macromolecules by SAS Using the Porod-Debye Law** *Biopolymers* **95**:559–571

59. Rust MJ, Markson JS, Lane WS, Fisher DS, O'Shea EK (2007) **Ordered phosphorylation governs oscillation of a three-protein circadian clock** *Science* **318**:809–812
 60. Sagar A, Svergun D, Bernado P, Kragelund B.B., Skriver K. (2020) **Structural Analyses of Intrinsically Disordered Proteins by Small-Angle X-Ray Scattering** *INTRINSICALLY DISORDERED PROTEINS: Methods and Protocols* :249–269
 61. Seal M, Jash C, Jacob RS, Feintuch A, Harel YS, Albeck S, Unger T, Goldfarb D (2021) **Evolution of CPEB4 Dynamics Across its Liquid–Liquid Phase Separation Transition** *The Journal of Physical Chemistry B* **125**:12947–12957
 62. Shapiro DM, Ney M, Eghtesadi SA, Chilkoti A (2021) **Protein Phase Separation Arising from Intrinsic Disorder: First-Principles to Bespoke Applications** *Journal of Physical Chemistry B* **125**:6740–6759
 63. Somjee R, Mitrea DM, Kriwacki RW (2020) **Exploring Relationships between the Density of Charged Tracts within Disordered Regions and Phase Separation, Pacific Symposium on Biocomputing** *Hi* :207–218
 64. Szabó AL, Santa A, Pancsa R, Gáspári Z (2022) **Charged sequence motifs increase the propensity towards liquid–liquid phase separation** *FEBS Letters* **596**:1013–1028
 65. Tang CT, Li S, Long C, Cha J, Huang G, Li L (2009) **Setting the pace of the Neurospora circadian clock by multiple independent FRQ phosphorylation events** *Proc Natl. Acad. Sci. USA* **106**:10722–10727
 66. Tatavosian R, Kent S, Brown K, Yao TT, Huy ND, Huynh TN, Zhen CY, Ma B, et al. (2019) **Nuclear condensates of the Polycomb protein chromobox 2 (CBX2) assemble through phase separation** *Journal of Biological Chemistry* **294**:1451–1463
 67. Trewthella J, Duff AP, Durand D, Gabel F, Guss JM, Hendrickson WA, Hura GL, Jacques DA, et al. (2017) **2017 publication guidelines for structural modelling of small-angle scattering data from biomolecules in solution: An update** *Acta Crystallographica Section D: Structural Biology* **73**:710–728
 68. Vernon RM, Chong PA, Tsang B, Kim TH, Bah A, Farber P, Lin H, Forman-Kay JD (2018) **Pi-Pi contacts are an overlooked protein feature relevant to phase separation** *Elife* **7**
 69. Wang B, Kettenbach AN, Zhou X, Loros JJ, Dunlap JC (2019) **The Phospho-Code Determining Circadian Feedback Loop Closure and Output in Neurospora** *Molecular Cell* **74**:771–784
 70. Wright PE, Dyson HJ (2014) **Intrinsically disordered proteins in cellular signalling and regulation** *Nature Reviews Molecular Cell Biology* **16**:18–29
 71. Xiao YB, Yuan Y, Jimenez M, Soni N, Yadlapalli S (2021) **Clock proteins regulate spatiotemporal organization of clock genes to control circadian rhythms** *Proc Natl. Acad. Sci. USA* **118**
 72. Zhou M, Guo JH, Cha J, Chae M, Chen S, Barral JM, Sachs MS, Liu Y (2013) **Non-optimal codon usage affects expression, structure and function of clock protein FRQ** *Nature* **495**:111–115
- Baker CL, Kettenbach AN, Loros JJ, Gerber SA, Dunlap JC (2009) **Quantitative Proteomics Reveals a Dynamic Interactome and Phase-Specific Phosphorylation in the Neurospora Circadian Clock** *Molecular Cell* **34**:354–363

Bartholomai BM (2022) **Cell biological dissection of the *Neurospora* circadian oscillator: Tools, approaches and fundamental observations**, Biochemistry and Genetics, Dartmouth College, Hanover NH

Bateman A, Martin MJ, Orchard S, Magrane M, Agivetova R, Ahmad S, Alpi E, Bowler-Barnett EH, et al. (2021) **UniProt: the universal protein knowledgebase in 2021** *Nucleic Acids Research* **49**:D480–D489

Bolognesi B, Gotor NL, Dhar R, Cirillo D, Baldrighi M, Tartaglia GG, Lehner B, Gaetano Tartaglia G (2016) **A Concentration-Dependent Liquid Phase Separation Can Cause Toxicity upon Increased Protein Expression** *CellReports* **16**:1–10

Chandrasekaran S, Schneps CM, Dunleavy R, Lin C, DeOliveira CC, Ganguly A, Crane BR (2021) **Tuning flavin environment to detect and control light-induced conformational switching in *Drosophila* cryptochrome** *Communications Biology* 2021 4:1 4:1–12

Cheng P, He Q, He Q, Wang L, Liu Y (2005) **Regulation of the *Neurospora* circadian clock by an RNA helicase** *Genes and Development* **19**:234–241

Guo J, Cheng P, Liu Y (2010) **Functional significance of FRH in regulating the phosphorylation and stability of *Neurospora* circadian clock protein FRQ** *J. Biol. Chem.* **285**:11508–11515

Holehouse AS, Das RK, Ahad JN, Richardson MOG, Pappu RV (2017) **CIDER: Resources to Analyze Sequence-Ensemble Relationships of Intrinsically Disordered Proteins** *Biophysical Journal* **112**:16–21

Hopkins JB, Gillilan RE, Skou S (2017) **BioXTAS RAW: Improvements to a free open-source program for small-angle X-ray scattering data reduction and analysis** *Journal of Applied Crystallography* **50**:1545–1553

Jankowski MS, Griffith D, Shastru DG, Pelham JF, Ginell GM, Thomas J, Karande P, Holehouse AS, et al. (2022) **The formation of a fuzzy complex in the negative arm regulates the robustness of the circadian clock** *bioRxiv*

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, et al. (2021) **Highly accurate protein structure prediction with AlphaFold** *Nature* 2021 596 **596**:583–589

Kyte J, Doolittle RF (1982) **A simple method for displaying the hydropathic character of a protein** *Journal of Molecular Biology* **157**:105–132

Mészáros B, Erdős G, Dosztányi Z (2018) **IUPred2A: context-dependent prediction of protein disorder as a function of redox state and protein binding** *Nucleic Acids Research* **46**:W329–W337

Muñoz V, Serrano L (1994) **Elucidating the folding problem of helical peptides using empirical parameters** *Nature Structural Biology* **1**:399–409

Muok AR, Chua TK, Le H, Crane BR (2018) **Nucleotide Spin Labeling for ESR Spectroscopy of ATP-Binding Proteins** *Applied Magnetic Resonance* **49**:1385–1395

- Nguyen GKT, Cao Y, Wang W, Liu CF, Tam JP (2015) **Site-Specific N-Terminal Labeling of Peptides and Proteins using Butelase 1 and Thiodepsipeptide** *Chemie-International Edition* **54**:15694–15698
- Querfurth C, Diernfellner A, Heise F, Lauinger L, Neiss A, Tataroglu Ö, Brunner M, Schafmeier T (2007) **Posttranslational regulation of Neurospora circadian clock by CK1 α -dependent phosphorylation** *Cold Spring Harbor symposia on quantitative biology* **72**:177–183
- Querfurth C, Diernfellner ACR, Gin E, Malzahn E, Höfer T, Brunner M (2011) **Circadian Conformational Change of the Neurospora Clock Protein FREQUENCY Triggered by Clustered Hyperphosphorylation of a Basic Domain** *Molecular Cell* **43**:713–722
- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C (2012) **Fiji: an open-source platform for biological-image analysis** *Nature Methods* **9**:676–682
- Somjee R, Mitrea DM, Kriwacki RW (2020) **Exploring Relationships between the Density of Charged Tracts within Disordered Regions and Phase Separation, Pacific Symposium on Biocomputing** *Hi* :207–218
- Srivastava M, Freed JH (2017) **Singular Value Decomposition Method to Determine Distance Distributions in Pulsed Dipolar Electron Spin Resonance** *Journal of Physical Chemistry Letters* **8**:5648–5655
- Srivastava M, Freed JH (2019) **Singular Value Decomposition Method To Determine Distance Distributions in Pulsed Dipolar Electron Spin Resonance: II. Estimating Uncertainty** *Journal of Physical Chemistry A* **123**:359–370
- Srivastava M, Georgieva ER, Freed JH (2017) **A New Wavelet Denoising Method for Experimental Time-Domain Signals: Pulsed Dipolar Electron Spin Resonance** *Journal of Physical Chemistry A* **121**:2452–2465
- Steinegger M, Söding J (2018) **Clustering huge protein sequence sets in linear time** *Nature Communications* **9**
- Tang CT, Li S, Long C, Cha J, Huang G, Li L, Chen S, Liu Y (2009) **Setting the pace of the Neurospora circadian clock by multiple independent FRQ phosphorylation events** *Proc. Natl. Acad. Sci. USA* **106**:10722–10727
- Vernon RM, Chong PA, Tsang B, Kim TH, Bah A, Farber P, Lin H, Forman-Kay JD (2018) **Pi-Pi contacts are an overlooked protein feature relevant to phase separation** *Elife* **7**
- Vogel HJ (1956) **A convenient growth medium for Neurospora (Medium N)** *Microb. Genet. Bull* **13**:42–43

Article and author information

Daniyal Tariq

Department of Chemistry & Chemical Biology, Cornell University, Ithaca, NY, 14853

Nicole Maurici

Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University, Syracuse, NY, 13210;

Bradley M. Bartholomai

Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth,
Hanover, NH 03755

Siddarth Chandrasekaran

Department of Chemistry & Chemical Biology, Cornell University, Ithaca, NY, 14853

Jay C. Dunlap

Department of Molecular and Systems Biology, Geisel School of Medicine at Dartmouth,
Hanover, NH 03755

Alaji Bah

Department of Biochemistry and Molecular Biology, SUNY Upstate Medical University,
Syracuse, NY, 13210;

For correspondence: BahA@upstate.edu

Brian R. Crane

Department of Chemistry & Chemical Biology, Cornell University, Ithaca, NY, 14853

For correspondence: bc69@cornell.edu

ORCID iD: [0000-0001-8234-9991](https://orcid.org/0000-0001-8234-9991)

Copyright

© 2023, Tariq et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Volker Dötsch

Goethe University, Germany

Senior Editor

Volker Dötsch

Goethe University, Germany

Reviewer #1 (Public Review):

Summary:

"Phosphorylation, disorder, and phase separation govern the behavior of Frequency in the fungal circadian clock" is a convincing manuscript that delves into the structural and biochemical aspects of FRQ and the FFC under both LLPS and non-LLPS conditions. Circadian clocks serve as adaptations to the daily rhythms of sunlight, providing a reliable internal representation of local time.

All circadian clocks are composed of positive and negative components. The FFC contributes negative feedback to the *Neurospora* circadian oscillator. It consists of FRQ, CK1, and FRH. The FFC facilitates close interaction between CK1 and the WCC, with CK1-mediated phosphorylation disrupting WCC:c-box interactions necessary for restarting the circadian cycle.

Despite the significance of FRQ and the FFC, challenges associated with purifying and stabilizing FRQ have hindered in vitro studies. Here, researchers successfully developed a protocol for purifying recombinant FRQ expressed in *E. coli*.

Armed with full-length FRQ, they utilized spin-labeled FRQ, CK1, and FRH to gain structural insights into FRQ and the FFC using ESR. These studies revealed a somewhat ordered core and a disordered periphery in FRQ, consistent with prior investigations using limited proteolysis assays. Additionally, p-FRQ exhibited greater conformational flexibility than np-FRQ, and CK1 and FRH were found in close proximity within the FFC. The study further demonstrated that under LLPS conditions in vitro, FRQ undergoes phase separation, encapsulating FRH and CK1 within LLPS droplets, ultimately diminishing CK1 activity within the FFC. Intriguingly, higher temperatures enhanced LLPS formation, suggesting a potential role of LLPS in the fungal clock's temperature compensation mechanism.

Biological significance was supported by live imaging of *Neurospora*, revealing FRQ foci at the periphery of nuclei consistent with LLPS. The amino acid sequence of FRQ conferred LLPS properties, and a comparison of clock repressor protein sequences in other eukaryotes indicated that LLPS formation might be a conserved process within the negative arms of these circadian clocks.

In summary, this manuscript represents a valuable advancement with solid evidence in the understanding of a circadian clock system that has proven challenging to characterize structurally due to obstacles linked to FRQ purification and stability. The implications of LLPS formation in the negative arm of other eukaryotic clocks and its role in temperature compensation are highly intriguing.

Strengths:

The strengths of the manuscript include the scientific rigor of the experiments, the importance of the topic to the field of chronobiology, and new mechanistic insights obtained.

Weaknesses:

This reviewer had questions regarding some of the conclusions reached.

Reviewer #2 (Public Review):

Summary:

This study presents data from a broad range of methods (biochemical, EPR, SAXS, microscopy, etc.) on the large disordered protein FRQ relevant to circadian clocks and its interaction partners FRH and CK1, providing novel and fundamental insight into oligomerization state, local dynamics, and overall structure as a function of phosphorylation and association. Liquid-liquid phase separation is observed. These findings have bearings on the mechanistic understanding of circadian clocks, and on functional aspects of disordered proteins in general.

Strengths:

This is a thorough work that is well presented. The data are of overall high quality given the difficulty of working with an intrinsically disordered protein, and the conclusions are sufficiently circumspect and qualitative to not overinterpret the mostly low-resolution data.

Weaknesses:

None

Reviewer #3 (Public Review):

Summary:

The manuscript from Tariq and Maurici et al. presents important biochemical and biophysical data linking protein phosphorylation to phase separation behavior in the

repressive arm of the *Neurospora* circadian clock. This is an important topic that contributes to what is likely a conceptual shift in the field. While I find the connection to the *in vivo* physiology of the clock to be still unclear, this can be a topic handled in future studies.

Strengths: The ability to prepare purified versions of unphosphorylated FRQ and P-FRQ phosphorylated by CK-1 is a major advance that allowed the authors to characterize the role of phosphorylation in structural changes in FRQ and its impact on phase separation *in vitro*.

Weaknesses: The major question that remains unanswered from my perspective is whether phase separation plays a key role in the feedback loop that sustains oscillation (for example by creating a nonlinear dependence on overall FRQ phosphorylation) or whether it has a distinct physiological role that is not required for sustained oscillation.